

# RoWAN COUNTY



---

## EMERGENCY SERVICES EMS DIVISION

---

*Be an original.*

## EMS System Protocols

Released September 2025

Revised August 2025

## 2025 Rowan EMS System Protocol Index

### Medical Director Memo

#### Protocol Introduction PI

PI-1. Introduction

PI-2. Key to Protocol Utilization

#### Adult Cardiac Section [AC]

AC-1. Asystole / Pulseless Electrical Activity

AC-2. Bradycardia – Pulse Present

AC-3. Cardiac Arrest

AC-4. Chest Pain and STEMI

AC-5. CHF Pulmonary Edema

AC-6. Adult Tachycardia – Narrow Complex

AC-7. Adult Tachycardia – Monomorphic Wide Complex

AC-8. Adult Tachycardia – Polymorphic Wide Complex

AC-9. V-Fib / Pulseless V-Tach

AC-10. Post Resuscitation

AC-11. TEAM Focused CPR

AC-12. Termination of CPR

AC-13. Targeted Temperature Management

AC-14. Ventricular Assist Device

AC-15. Total Mechanical Circulation

AC-16. Wearable Cardioverter Defibrillator Vest

#### Adult Medical [AM]

AM-1. Allergic Reaction / Anaphylaxis

AM-2. Diabetic; Adult

AM-3. Dialysis – Renal Failure

AM-4. Hypertension

AM-5. Hypotension – Shock

AM-6. Suspected Stroke – Activae / t-PA Transfer

Adult Obstetrical [AO]

AO-1. Childbirth Labor

AO-2. Newly Born

AO-3. Obstetrical Emergency

Airway Respiratory [AR]

AR-1. Adult Airway

AR-2. Adult Failed Airway

AR-3. Airway, Drug Assisted

AR-4. COPD Asthma

AR-5. Pediatric Airway

AR-6. Pediatric Failed Airway

AR-7. Pediatric Respiratory Distress

AR-8. Post Intubation / BIAD Management

AR-9. Ventilator Emergencies

AR-10. Tracheostomy Tube Emergencies

AR-11. Mechanical Ventilation

Pediatric Cardiac [PC]

PC-1. Pediatric Asystole / PEA

PC-2. Pediatric Bradycardia – Poor Perfusion

PC-3. Pediatric CHF / Pulmonary Edema

PC-4. Pediatric Pulseless Arrest

PC-5. Pediatric Tachycardia – Narrow Complex

PC-6. Pediatric Tachycardia – Wide Complex

PC-7. Pediatric V-Fib / Pulseless V-Tach

PC-8. Pediatric Post Resuscitation

Pediatric Medical [PM]

PM-1. Pediatric Allergic Reaction

PM-2. Pediatric Diabetic

PM-3. Pediatric Hypotension – Shock

Special Circumstances [SC]

SC-1. Suspected Ebola

SC-2. Respiratory Disease

SC-3. Hospice or Palliative Care Patient

SC-4. Vaccination

SC-5. SARS CoV2 Monoclonal Antibody Infusion

Special Operations [SO]

SO-1. Scene Rehabilitation General

SO-2. Scene Rehabilitation – Responder

Toxins and Environmental [TE]

TE-1. Bites and Envenomations

TE-2. Carbon Monoxide / Cyanide

TE-3. Drowning / Submersion Injury

TE-4. Hyperthermia

TE-5. Hypothermia

TE-6. Marine Envenomation

TE-7. Overdose / Toxic Ingestion

TE-8. WMD / Nerve Agent Protocol

## Trauma and Burn [TB]

- TB-1. Blast Injury
- TB-2. Chemical and Electrical Burn
- TB-3. Crush Syndrome
- TB-4. Extremity Trauma
- TB-5. Head Trauma
- TB-6. Multiple Trauma
- TB-7. Radiation Incident
- TB-8. Selective Spinal Immobilization
- TB-9. Thermal Burn
- TB-10. Traumatic Arrest

## Universal Protocols [UP]

- UP-1. Universal Patient Care
- UP-2. Triage Protocol
- UP-3. Abdominal Pain / Vomiting / Diarrhea
- UP-4. Altered Mental Status
- UP-5. Back Pain
- UP-6. IV or IO Access
- UP-7. Dental Problems
- UP-8. Emergencies Involving Indwelling Catheters / Central Lines
- UP-9. Epistaxis
- UP-10. Fever
- UP-11. Pain Control – Adult
- UP-12. Police Custody
- UP-13. Seizure
- UP-14. Suspected Stroke
- UP-15. Suspected Sepsis

UP-16. Syncope

UP-17. Behavioral Health Crisis

UP-18. Behavioral Agitation Sedation Guide

UP-19. Behavioral – Hyperactive Delirium with Severe Agitation

UP-20. Sickle Cell Crisis

Triage Destination Plans

General Medical

Pediatric

STEMI

Stroke and LVO

Trauma and Burn

Drug List (Not Comprehensive of all medications)



*Be an original.*

2727 Old Concord Rd, Suite E, Salisbury, NC, 28146-8388  
[p] 704-216-8900 [f] 704-216-8921

30 July 2025

To: Rowan County EMS System Personnel  
  
From: Ronnie Barrier, MD – Medical Director  
Bertrand Fote, MD, FACEP – Associate Medical Director  
  
Re: 2025 – NC College of Emergency Physicians Protocols, Policies, and Procedures

The 2024-2025 NC College of Emergency Physicians Protocols, Policies, and Procedures have been reviewed, updated and approved. These are treatment guidelines for patients that you may encounter but should not replace common sense and excellent clinical judgement. Should you have any questions regarding patient treatment, or applicability regarding a treatment plan, contact Medical Control for immediate guidance. Policy questions needing immediate attention during a patient encounter not related to your treatment plan should be directed to the on-duty Rowan County EMS Division Shift Commander, or EMS Administrative staff.

These guidelines were approved and discussed during the March, and July 2025 EMS System Peer Review meeting. Each EMS provider should familiarize themselves with the layout of these guidelines so they can be quickly referenced when needed. Continuing Education sessions over the past few months, and in the upcoming months will review various Protocols, Policies, and Procedures. The planned implementation of these guidelines is September 1, 2025 for the Rowan County EMS System.

# Introduction

The following medical treatment protocols are developed for North Carolina EMS agencies. The process has evolved since 2007 and continues with input from Medical Directors, EMS Administration, North Carolina Chapter of Emergency Physicians Protocol Committee, North Carolina Office of EMS, EMS field personnel and the public at large through on-line surveys, public meetings across North Carolina and direct communication with stakeholders. The 2017 update expands on the 2012 and 2009 version and continues to incorporate evidence-based guidelines, expert opinion and historically proven practices meant to ensure that citizens and visitors of North Carolina will continue to be provided the highest quality pre-hospital patient care available. The North Carolina Chapter of Emergency Physicians develops and provides final approval.

The purpose of the protocol section is to provide treatment protocols outlining permissible and appropriate assessment, delivery of care, reassessment and procedures which may be rendered by pre-hospital providers. The protocols also outline which medical situations require direct voice communication with medical control. In general treatment protocols are specific orders which may and should be initiated prior to contact with Medical Control.

**Please note the medical protocols are divided into three (3) to four (4) sections.** The upper section includes three (3) boxes (History, Signs and Symptoms and Differential) which serves as a guide to assist in obtaining pertinent patient information and exam findings as well as considering multiple potential causes of the patients complaint. It is not expected that every historical element or sign / symptom be recorded for every patient. It is expected that those elements pertinent to your patient encounter will be included in the patient evaluation.

**The algorithm section describes the essentials of patient care. Virtually every patient should receive the care outlined in this section, usually in the order described. However each medical emergency must be dealt with individually and appropriate care determined accordingly. Professional judgment is mandatory in determining treatment modalities within the parameters of these protocols. Circumstances will arise where treatment may move ahead in the algorithm, move outside to another protocol and then re-enter later. While protocols are written based on body systems and primary complaints the patient should be treated as a whole and therefore the protocols should be considered as a whole in providing care.**

## Professional judgment hierarchy:

The pre-hospital provider may determine that no specific treatment is needed;

Or

The pre-hospital provider may follow the appropriate treatment protocols and then consult Medical Control;

Or

The pre-hospital provider may consult Medical Control before initiating any specific treatment.

**Some protocols will encompass two (2) pages.** Protocols which exist in a single page format may have page 2 added by the local medical director. The PEARLS section will either be located at the bottom of page 1 (single page protocol) or page 2 (double page protocol). The PEARLS section provides points regarding the main protocol based on evidence to date, common medical knowledge and expert medical opinion.

**Information boxes highlighted in purple.** These areas are editable at the local level. They will mainly involve specific medications and dosages utilized by the local EMS agency. Page 2 will have a large section highlighted in purple where the local Medical Director may edit as they see fit to provide expanded points and treatment not otherwise specified in the algorithm. If the box is not to be utilized – add “***This Space Left Blank Intentionally.***”

Finally these medical treatment protocols are established to ensure safe, efficient and effective interventions to relieve pain and suffering and improve patient outcomes without inflicting harm. They also serve to ensure a structure of accountability for Medical Directors, EMS agencies, pre-hospital providers and facilities to provide continual performance improvement. A recent report of the Institute of Medicine calls for the development of standardized, evidence-based pre-hospital care protocols for the triage, treatment and transport of patients. These protocols establish expectations of pre-hospital care in North Carolina.

# Key to Protocol Utilization

## History

- Important history items
- Circumstances of event
- SAMPLE
- Time of onset
- Duration

## Signs and Symptoms

- Important Signs and Symptoms specific to each protocol

## Differential

- A list of other disease or injury which should be considered

Black Box

Highlights  
Important  
Information



**Universal Patient Care Protocol**  
*Assumed all protocols utilize and will not appear on individual protocols*

Red Box

Highlights  
Critical  
Information

May direct to  
another  
protocol



Signals protocol within a protocol

Indicates  
Entry / Exit  
from / to  
to another protocol(s)



Information box

Decision Point  
Darker outline to highlight

Highlights medication after  
Contact Medical Control

May be added by Local Medical Director

## Purple Shading of Information Box

Indicates items changeable at local agency level, including medications / dosages on NCMB formulary

Local Medical Director may add / change at his / her discretion

Local medical director may add page 2 to any protocol where none exists for additional comments

## Algorithm Legend

	Emergency Medical Responder
B	Emergency Medical Technician
A	Advanced Emergency Medical Technician
P	Paramedic
	Notify Destination or Contact Medical Control

## Pearls

- Important information specific to each protocol will appear here.
- Will usually appear on page.
- Important exam items listed here specific to protocol.



# Adult Asystole / Pulseless Electrical Activity

## History

- SAMPLE
- Estimated downtime
- See Reversible Causes below
- DNR, MOST, or Living Will

## Signs and Symptoms

- Pulseless
- Apneic
- No electrical activity on ECG
- No heart tones on auscultation

## Differential

- See Reversible Causes below

### Cardiac Arrest Protocol AC 3

#### Criteria for Death / No Resuscitation Review DNR / MOST Form

NO

Decomposition  
Rigor mortis  
Dependent lividity  
Blunt force trauma  
Injury incompatible with life  
Extended downtime with asystole

Do not begin resuscitation

Follow  
Deceased Subjects  
Policy

## AT ANY TIME

Return of  
Spontaneous  
Circulation



Go to  
Post Resuscitation  
Protocol AC 10

		<b>Begin Continuous CPR Compressions</b> Push Hard ( $\geq$ 2 inches) Push Fast (100 - 120 / min) Change Compressors every 2 minutes (sooner if fatigued) (Limit changes / pulse checks $\leq$ 10 seconds)
		Ventilate 1 breath every 6 seconds 30:2 Compression:Ventilation if no Advanced Airway Monitor EtCO <sub>2</sub> if available
P		AED Procedure if available
	P	Cardiac Monitor
		IV or IO Access Protocol UP 6
A		Epinephrine (1:10,000) 1 mg IV / IO Repeat every 5 minutes
	A	Normal Saline Bolus 500 mL IV / IO May repeat as needed Maximum 2 L
		Search for Reversible Causes
		Blood Glucose Analysis Procedure if applicable
P		Available for Agency Medications
		This area intentionally left blank
		On Scene Resuscitation / Termination of Resuscitation Protocol(s) AC 12 as indicated

Notify Destination or  
Contact Medical Control

### Reversible Causes

Hypovolemia  
Hypoxia  
Hydrogen ion (acidosis)  
Hypothermia  
Hypo / Hyperkalemia

Tension pneumothorax  
Tamponade; cardiac  
Toxins  
Thrombosis; pulmonary (PE)  
Thrombosis; coronary (MI)

### Suspected Opioid Overdose

Administer Naloxone per  
Overdose / Toxic Ingestion  
Protocol TE 7



# Adult Asystole / Pulseless Electrical Activity

## Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional Team Focused CPR Protocol AC 11 or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT), compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- **Reassess and document BIAD and / or endotracheal tube placement and EtCO<sub>2</sub> frequently, after every move, and at transfer of care.**
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation: Follow manufacturer's recommendations concerning defibrillation / cardioversion energy when specified.**
- **End Tidal CO<sub>2</sub> (EtCO<sub>2</sub>):**
  - If EtCO<sub>2</sub> is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
  - If EtCO<sub>2</sub> spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Special Considerations**
  - Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
  - Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
  - Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
  - Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Transcutaneous Pacing:**
  - Pacing is NOT effective in cardiac arrest and pacing in cardiac arrest does NOT increase chance of survival
  - Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
  - Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.



# Bradycardia; Pulse Present

## History

- Past medical history
- Medications
  - Beta-Blockers
  - Calcium channel blockers
  - Clonidine
  - Digoxin
- Pacemaker

## Signs and Symptoms

- HR < 60/min with hypotension, acute altered mental status, chest pain, acute CHF, seizures, syncope, or shock secondary to bradycardia
- Chest pain
- Respiratory distress
- Hypotension or Shock
- Altered mental status
- Syncope

## Differential

- Acute myocardial infarction
- Hypoxia / Hypothermia
- Pacemaker failure
- Sinus bradycardia
- Head injury (elevated ICP) or Stroke
- Spinal cord lesion
- Sick sinus syndrome
- AV blocks (1°, 2°, or 3°)
- Overdose

Exit to  
Appropriate  
Protocol(s)



**Heart Rate < 60 / min and Symptomatic:**  
Hypotension, Acute AMS, Ischemic Chest Pain,  
Acute CHF, Seizures, Syncope, or Shock  
secondary to bradycardia  
Typically HR < 50 / min

YES

Airway Protocol(s) AR 1, 2, 3  
*if indicated*



Respiratory Distress  
Protocol AR 4  
*if indicated*



Chest Pain: Cardiac and STEMI  
Protocol AC 4  
*if indicated*



**B** Search for Reversible Causes



12 Lead ECG Procedure



IV / IO Protocol UP 6



**P** Cardiac Monitor



**A** Normal Saline Fluid Bolus  
500 mL – 2 L NS IV / IO  
(Unless Acute CHF)  
Maximum 2 L



Atropine 1 mg IV / IO  
May repeat every 3 – 5 minutes  
Maximum 3 mg



Epinephrine 1 - 10 mcg/min IV / IO  
Titrate to SBP  $\geq$  90 mmHg  
Or  
Dopamine 2 – 20 mcg/kg/min IV / IO  
Titrate to SBP  $\geq$  90 mmHg



Agency Vasopressor



If No Improvement  
Transcutaneous Pacing Procedure  
*(Consider earlier in 2<sup>nd</sup> or 3<sup>rd</sup> AVB)*



Notify Destination or  
Contact Medical Control

## Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypothermia
- Hypo / Hyperkalemia
- Tension pneumothorax
- Tamponade; cardiac
- Toxins
- Thrombosis; pulmonary (PE)
- Thrombosis; coronary (MI)

**Suspected Beta-Blocker or Calcium Channel Blocker**

Follow Overdose/  
Toxic Ingestion  
Protocol TE 7

Consider Sedation



**P** Midazolam 2 – 2.5 mg  
IV / IO / IM / IN

Maximum 10 mg




# Bradycardia; Pulse Present

## Pearls

- Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro
- Identifying signs and symptoms of poor perfusion caused by bradycardia are paramount.
- Rhythm should be interpreted in the context of symptoms and pharmacological treatment given only when symptomatic, otherwise monitor and reassess.
- Consider hyperkalemia with wide complex, bizarre appearance of QRS complex, and bradycardia. Give Calcium Chloride or Gluconate in addition to Sodium Bicarbonate if hyperkalemia suspected.
- 12-Lead ECG:
  - 12 Lead ECG not necessary to diagnose and treat
  - Obtain when patient is stable and/or following rhythm conversion.
- Hypoxemia is a common cause of bradycardia. Ensure oxygenation and support respiratory effort.
- Atropine:
  - Atropine is considered a first line agent in symptomatic bradycardia.
  - Ineffective and potentially harmful in cardiac transplantation. May cause paradoxical bradycardia.
- Symptomatic bradycardia causing shock or peri-arrest condition:
  - If no IV or IO access immediately available start Transcutaneous Pacing, establish IV / IO access, and then administer atropine and/or epinephrine.
  - Epinephrine or Dopamine may be considered if no response to Atropine.
- Symptomatic condition
  - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
  - Symptomatic bradycardia usually occurs at rates < 50 beats per minute.
  - Search for underlying causes such as hypoxia or impending respiratory failure.
- Serious Signs / Symptoms:
  - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.
- Transcutaneous Pacing Procedure (TCP)
  - Indicated with unstable bradycardia unresponsive to medical therapy.
  - If time allows transport to specialty center because transcutaneous pacing is a temporizing measure.
  - Transvenous / permanent pacemaker will probably be needed.
  - Immediate TCP with high-degree AV block (2d or 3d degree) with no IV / IO access.
- Consider treatable causes for bradycardia (Beta Blocker OD, Calcium Channel Blocker OD, etc.)



# Cardiac Arrest; Adult





# Cardiac Arrest; Adult

## Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assign responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT) compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- **Reassess and document BIAD and / or endotracheal tube placement and EtCO<sub>2</sub> frequently, after every move, and at transfer of care.**
- **IV / IO access and drug delivery is secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation:**  
Follow manufacturer's recommendations concerning defibrillation / cardioversion energy when specified.  
Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.  
Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO<sub>2</sub> (EtCO<sub>2</sub>)**  
If EtCO<sub>2</sub> is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.  
If EtCO<sub>2</sub> spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Special Considerations**  
**Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.  
**Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.  
**Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.  
**Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Transcutaneous Pacing:**  
Pacing is NOT effective in cardiac arrest and pacing in cardiac arrest does NOT increase chance of survival
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
- Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.

AC 3



# Chest Pain: Cardiac and STEMI

## History

- Age
- Medications (Viagra / sildenafil, Levitra / vardenafil, Cialis / tadalafil)
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Allergies
- Recent physical exertion
- **Onset / Palliation / Provocation**
- Quality (crampy, constant, sharp, dull, etc.)
- **Region / Radiation / Referred**
- **Severity (1-10)**
- **Time (onset /duration / repetition)**

## Signs and Symptoms

- CP (pain, pressure, aching, vice-like tightness)
- Location (substernal, epigastric, arm, jaw, neck, shoulder)
- Radiation of pain
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness
- **Time of Onset**
- Women:
  - More likely to have dyspnea, N/V, weakness, back or jaw pain

## Differential

- Trauma vs. Medical
- Angina vs. Myocardial infarction
- Pericarditis
- Pulmonary embolism
- Asthma / COPD
- Pneumothorax
- Aortic dissection or aneurysm
- GE reflux or Hiatal hernia
- Esophageal spasm
- Chest wall injury or pain
- Pleural pain
- Overdose: Cocaine or Methamphetamine

### 12 Lead ECG Procedure

**B** Aspirin 81 mg x 4 PO (chewed)  
Or 325 mg PO

Nitroglycerin 0.3 / 0.4 mg Sublingual  
Repeat every 5 minutes x 3  
*if prescribed to patient and (BP ≥ 100)*

**P** Cardiac Monitor

**Acute MI / STEMI**  
See box to right

YES

Transport based on:  
**STEMI**  
**EMS Triage and Destination Plan**  
**Immediate Notification of Facility**  
**Immediate Transmission of ECG if capable**  
**Keep Scene Time to ≤ 15 Minutes**

## Acute MI / STEMI

### STEMI Definition:

- $\geq 1$  mm ST Segment elevation in  $\geq 2$  contiguous leads
- $\geq 2$  mm ST/J point elevation in V2-V3 for men
- $\geq 1.5$  mm ST/J point elevation in V2-V3 for women
- ECG software diagnoses Acute MI (symptomatic)

	IV / IO Protocol UP 6
	Nitroglycerin 0.3 / 0.4 mg SL Repeat every 5 minutes as needed
	Nitroglycerin Paste
	Morphine 2 – 4 mg IV / IO Repeat every 5 minutes as needed <b>Maximum 10 mg</b> Or Fentanyl 50 – 75 mcg IV / IO Repeat 25 mcg every 20 minutes as needed <b>Maximum 200 mcg</b>
	Hypotension / Shock Protocol AM 5 <i>if indicated</i>
	CHF / Pulmonary Edema Protocol AC 5 <i>if indicated</i>

**B** If transporting to Non PCI Center  
**Reperfusion Checklist**

**P** Agency Specific Medications  
Thrombolytic  
Heparin  
Plavix

Notify Destination or  
Contact Medical Control

**AC 4**



# Chest Pain: Cardiac and STEMI

Patients who are abusing cocaine will show signs of agitation and have dilated pupils. First-line treatment for cocaine-induced arrhythmias and hypertensive episodes is usually benzodiazepine administration, which tempers the effects of cocaine on the central nervous system and cardiovascular system. Versed or Valium may be administered. If any question, contact Medical Control.

When encountering Non-ST change chest pain that is potentially Acute Coronary Syndrome (ACS) related you should consider the HEART Score for risk stratification. While troponin may not be immediately available in the field, the other score portions can show high/low risk for the patient that can be relayed to the receiving facility. This can be used in patients > 21 years old presenting with symptoms suggestive of ACS. Do not use if new ST-segment elevation > 1mm or other new ECG changes, hypotension, life expectancy less than 1 year, or non-cardiac medical/surgical/psychiatric illness determined.

- H – History

Slightly Suspicious = 0

Moderately suspicious = +1

Highly suspicious = +2

- E – ECG –

- 1 point: No ST depression but LBBB, LVH, repolarization changes (ex. Digoxin);
- 2 points: ST depression / elevation not due to LBBB, LVH, or digoxin
- Normal = 0
- Non-specific repolarization disturbance = +1
- Significant ST depression = +2

A - Age

< 45 = 0

45-64 = +1

> 65 = +2

- R – Risk Factors: HTN, hypercholesterolemia, DM, obesity (BMI >30), smoking (current or cessation less than 3 months), positive family history (parent or sibling with CVD before age 65); atherosclerotic disease; prior MI, PCI/CABG, CVA/TIA, or peripheral arterial disease

No known risk factors = 0

1-2 risk factors = +1

> 3 risk factors or history of atherosclerotic disease = +2

- T – Initial Troponin: Use local assays and corresponding cutoffs

< normal limit = 0

-2x normal limit = +1

> 2x normal limit = +2

Scores 0-3: 0.9-1.7% risk of adverse cardiac event.

Scores 4-6: 12-16.6% risk of adverse cardiac event.

Scores > 7: 50-65% risk of adverse cardiac event – may be candidates for early invasive measures.

Obtain serial 12 leads after medication administration to follow the potential evolution/changes with the ECG.

Notify the receiving facility of the STEMI – Speak with physician during encode to activate the cath-lab.

## Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Items in Red Text are the key performance indicators for the EMS Acute Cardiac (STEMI) Care Toolkit**
- **Nitroglycerin:**
  - Avoid Nitroglycerin in any patient who has used Viagra (sildenafil) or Levitra (vardenafil) in the past 24 hours or Cialis (tadalafil) in the past 36 hours due to potential severe hypotension.
  - Nitroglycerin may cause hypotension during any type myocardial infarction. It is NOT more likely to cause hypotension in an inferior MI and should NOT be avoided unless already hypotensive.
- **STEMI (ST-Elevation Myocardial Infarction)**
  - Positive Reperfusion Checklist should be transported to the appropriate facility based on STEMI EMS Triage and Destination Plan.**
  - Consider placing 2 IV sites in the left arm: Many PCI centers use the right radial artery for intervention.
  - Consider placing defibrillator pads on patient as a precaution.
  - Consider Normal Saline or Lactated Ringers bolus of 250 – 500 mL as pre-cath hydration.
  - Scene time goal is < 15 minutes.
  - Document and time-stamp facility STEMI notification and make notification as soon as possible.
  - Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).
- **Cardiac related symptoms in men and women:**
  - Pressure, squeezing, fullness, or pain in the chest.
  - Pain or discomfort in one or both arms, the back, neck, jaw, or stomach.
  - Shortness of breath with or without chest pain.
  - Sweating, nausea, weakness, and/or lightheadedness.

**Women, diabetic patients, and the elderly often experience only weakness, shortness of breath, nausea/vomiting, and back or jaw pain.**

## EMT:

- **EMT administration of Nitroglycerin is limited to patients currently prescribed the medication, unless approved by the Agency Medical Director and the NC office of EMS.**
- If patient has taken nitroglycerin without relief, consider potency of the medication.
- Monitor for hypotension after administration of nitroglycerin and opioids.
- Agency medical director may require Contact of Medical Control prior to administration.

**AC 4**



# CHF / Pulmonary Edema

## History

- Congestive heart failure
- Past medical history
- Medications (digoxin, Lasix, Viagra / sildenafil, Levitra / vardenafil, Cialis / tadalafil)
- Cardiac history --past myocardial infarction

## Signs and Symptoms

- Respiratory distress, bilateral rales
- Apprehension, orthopnea
- Jugular vein distention
- Pink, frothy sputum
- Peripheral edema, diaphoresis
- Hypotension, shock
- Chest pain

## Differential

- Myocardial infarction
- Congestive heart failure
- Asthma
- Anaphylaxis
- Aspiration
- COPD
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pericardial tamponade
- Toxic Exposure

	Airway Protocol(s) AR 1, 2, 3 <i>as indicated</i>
	Chest Pain and STEMI Protocol AC 4 <i>if indicated</i>
B	12 Lead ECG Procedure
B	<b>Nitroglycerin 0.3 / 0.4 mg Sublingual</b> Repeat every 5 minutes x 3 <i>if prescribed to patient and (BP &gt;100)</i>
P	Cardiac Monitor
	IV / IO Procedure

Assess Symptom Severity

### MILD

Normal Heart Rate  
Elevated or Normal BP

**Nitroglycerin 0.3 / 0.4 mg SL**  
Repeat every 5 minutes

A

Nitroglycerin Paste

Improving

NO

### MODERATE / SEVERE

Elevated Heart Rate  
Elevated BP

**B** Airway NIPPV Procedure

**Nitroglycerin 0.3 / 0.4 mg SL**  
Repeat every 5 minutes

A

Nitroglycerin Paste

ACE-Inhibitor

Consider **Furosemide 40 mg IV**  
ONLY IF  
Transport time > 30 minutes  
Known CHF / Daily Lasix  
Afebrile

YES

Notify Destination or  
Contact Medical Control

### CARDIOGENIC SHOCK

Tachycardia followed by bradycardia  
Hypertension followed by hypotension

B

Remove NIPPV  
*if in place*

Adult Hypotension / Shock  
Protocol AM 5  
*if indicated*

AC 5



# CHF / Pulmonary Edema

Management of heart failure is geared toward improving gas exchange and cardiac output. If the actual blood pressure is adequate (systolic blood pressure > 100 mm/Hg), help the patient get into a comfortable position. Many times this can be done with the patient sitting with legs dependent. Oxygen saturations above 90% are desired, so you should evaluate the patient for possible ventilatory assistance. If there are signs of respiratory failure along with altered mental status, intubation and invasive pulmonary ventilation will be necessary. If the patient is alert enough, noninvasive positive pressure ventilation (NIPPV) can be therapeutic in two ways: (1) decreasing venous return and preload, thereby reducing pulmonary edema, and (2) improving gas exchange.

Along with positive pressure ventilation, if the systolic blood pressure is above 100 mm/Hg, nitroglycerin has emerged as the primary treatment of pulmonary edema. Nitroglycerin acts to decrease preload through peripheral vasodilation. Caution must be exercised when employing nitroglycerin and positive pressure ventilation simultaneously; systemic blood pressure can drop quickly. Patients with subacute CHF who also feel volume overloaded may be given furosemide to initiate diuresis. Furosemide should also be used with caution because many patients with "crackles" on exam are later found to have pneumonia. Diuresis in this group of patients can be detrimental. In addition, many of those who do have CHF are not total-body-fluid overloaded; the fluid is just not distributed correctly. Diuresis can be detrimental in these patients because many have poor renal function to begin with.

The Rowan EMS System does not currently carry an ACE-inhibitor

## Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care**
- **Diuretics (furosemide) and opioids have NOT been shown to improve the outcomes of EMS patients with pulmonary edema. Even though this historically has been a mainstay of EMS treatment, it is no longer routinely recommended.**
- **Nitroglycerin:**
  - Avoid Nitroglycerin in any patient who has used Viagra (sildenafil) or Levitra (vardenafil) in the past 24 hours or Cialis (tadalafil) in the past 36 hours due to potential severe hypotension.
  - Nitroglycerin may cause hypotension during any type myocardial infarction. It is NOT more likely to cause hypotension in an inferior MI and should NOT be avoided unless already hypotensive.
- Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).
- Consider myocardial infarction in all these patients. Diabetics, geriatric and female patients often have atypical pain, or only generalized complaints.
- **Cardiac related symptoms in men and women:**
  - Pressure, squeezing, fullness, or pain in the chest.
  - Pain or discomfort in one or both arms, the back, neck, jaw, or stomach.
  - Shortness of breath with or without chest pain.
  - Sweating, nausea, weakness, and/or lightheadedness.
  - Women, diabetic patients, and the elderly often experience only weakness, shortness of breath, nausea/vomiting, and back or jaw pain.**
- If patient has taken nitroglycerin without relief, consider potency of the medication.
- Contraindications to opioids include severe COPD and respiratory distress. Monitor the patient closely.
- Monitor for hypotension after administration of nitroglycerin and opioids.
- Allow the patient to be in their position of comfort to maximize their breathing effort.
- **EMT may administer Nitroglycerin to patients already prescribed medication. May give from EMS supply.**
- Agency medical director may require Contact of Medical Control.



# Adult Tachycardia

## NARROW ( $\leq 0.11$ sec)

### History

- Age
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Recent physical exertion
- Palpitations, irregular heart beat
- Time (onset /duration / repetition)

### Signs and Symptoms

- Chest pain, heart failure, dyspnea
- AMS
- Shock, poor perfusion, hypotension
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness

### Differential

- Trauma vs. Medical
- Sinus Tachycardia vs. dysrhythmia
- Fever, sepsis, infection
- Pericarditis, pulmonary embolism
- Aortic dissection or aneurysm
- Overdose: Stimulants

Assess tachycardia in context of clinical condition  
Identify and treat underlying cause of tachycardia



#### Unstable / Serious Signs and Symptoms

HR Typically  $\geq 150$

Hypotension, Acute AMS, Ischemic Chest Pain, Acute CHF, Seizures, Syncope, or Shock secondary to tachycardia

YES → P

NO

#### Consider

Only if regular/narrow complex

Adenosine 6 mg IV / IO

Rapid push with flush

May repeat 12 mg IV / IO

Cardioversion Procedure

#### Consider Sedation Prior to Cardioversion

Midazolam 2 – 2.5 mg IV / IO

May repeat as needed

Maximum 10 mg

#### Synchronized

Narrow and Regular: 50 – 100J

Narrow and Irregular: 120 – 200J

May repeat and increase dose with subsequent cardioversion attempts

B	12 Lead ECG Procedure
P	Cardiac Monitor
IV	IV or IO Access Protocol UP 6



#### Regular Rhythm?

YES → P

NO

#### Attempt Vagal Maneuvers Procedure

Adenosine 6 mg IV / IO

Rapid push with flush

May repeat 12 mg IV / IO

May repeat 12 mg IV / IO

P	<p><b>Diltiazem 0.25 mg/kg IV / IO</b> Over 2 – 3 minutes <b>Maximum 25 mg</b></p> <p><u>If No Improvement</u> Diltiazem 5 – 15 mg/hr IV / IO</p> <p><u>If No Improvement in 15 minutes</u> Diltiazem 0.35 mg/kg IV / IO <b>Maximum 25 mg</b></p> <p>Or Amiodarone 150 mg in 100 mL of D5W IV / IO</p> <p>Or <b>Agency Specific Beta-blocker</b></p>
	Monitor and Reassess
TP	Notify Destination or Contact Medical Control



# Adult Tachycardia

## NARROW ( $\leq 0.11$ sec)

If treating with Diltiazem/Cardiazem for irregular rhythm and the patient is conscious and borderline hypotensive, consider treatment with calcium chloride prior to administration of Diltiazem / Cardiazem.

### Pearls

- **Recommended Exam:** Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro
- **Most important goal** is to differentiate the type of tachycardia and if STABLE or UNSTABLE and SYMPTOMATIC.
- **12-Lead ECG:**
  - 12 Lead ECG not necessary to diagnose and treat
  - Obtain when patient is stable and/or following rhythm conversion.
- **Unstable condition**
  - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
  - If at any point patient becomes unstable move to unstable arm in algorithm.
- **Search for underlying cause** of tachycardia such as fever, sepsis, dyspnea, etc.
- **Typical sinus tachycardia** is in the range of 100 to (200 - patient's age) beats per minute.
- **Symptomatic condition**
  - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
  - Symptomatic tachycardia usually occurs at rates  $\geq 150$  beats per minute.
  - Patients symptomatic with heart rates  $< 150$  likely have impaired cardiac function such as CHF.
- **Serious Signs / Symptoms:**
  - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.
- **If patient has history or 12 Lead ECG reveals Wolfe Parkinson White (WPW):**
  - DO NOT administer a Calcium Channel Blocker (e.g. Diltiazem) or Beta Blockers.
  - Use caution with Adenosine and give only with defibrillator available.
- **Regular Narrow-Complex Tachycardia:**
  - Vagal maneuvers and adenosine are preferred. Vagal maneuvers may convert 19% to 54 % of SVT.
  - Using passive leg raise with Valsalva is more effective.
  - Adenosine should be pushed rapidly via proximal IV site followed by 20 mL Normal Saline rapid flush.
  - Adenosine should not be used in the post-cardiac transplant patient without **Contact of Medical Control**.
  - Agencies using both calcium channel blockers and beta blockers should choose one primarily. Giving the agents sequentially requires **Contact of Medical Control**. This may lead to profound bradycardia / hypotension.
- **Irregular Narrow-Complex Tachycardia:**
  - Rate control is more important in pre-hospital setting rather than focus on rhythm conversion.
- **Synchronized Cardioversion:**
  - Recommended to treat UNSTABLE Atrial Fibrillation, Atrial Flutter and SVT.
  - Monitor for hypotension after administration of Calcium Channel Blockers or Beta Blockers.
  - Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.



# Adult Monomorphic Tachycardia

## Wide Complex ( $\geq 0.12$ sec)

### History

- Age
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Recent physical exertion
- Palpitations, irregular heart beat
- Time (onset /duration / repetition)

### Signs and Symptoms

- Chest pain, heart failure, dyspnea
- AMS
- Shock, poor perfusion, hypotension
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness

### Differential

- Trauma vs. Medical
- Sinus Tachycardia vs. dysrhythmia
- Fever, sepsis, infection
- Pericarditis, pulmonary embolism
- Aortic dissection or aneurysm
- Overdose: Stimulants

Assess tachycardia in context of clinical condition  
Identify and treat underlying cause of tachycardia



#### Unstable/ Serious Signs and Symptoms

HR Typically  $> 150$

Hypotension, Acute AMS, Ischemic Chest Pain, Acute CHF, Seizures, Syncope, or Shock secondary to tachycardia

NO

12 Lead ECG Procedure

Cardiac Monitor

IV or IO Access Protocol UP 6

**P** Consider consultation with medical control

#### Regular Rhythm?

NO

**P** Amiodarone 150 mg in 100 mL of D5W IV / IO Infuse over 10 minutes

May repeat if wide complex tachycardia recurs

Amiodarone 450 mg in 250 mL of D5W 1 mg/min (33 mL/hr) IV / IO

Or

Procainamide 20 – 50 mg / min IV / IO

Procainamide 1 – 4 mg / min

Maximum 17 mg / kg

Monitor and Reassess

**T** Notify Destination or Contact Medical Control

**P** Cardiac Monitor

Cardioversion Procedure

#### Consider Sedation Prior to Cardioversion

Midazolam 2 – 2.5 mg IV / IO

May repeat as needed

Maximum 10 mg

#### Wide: Regular and Irregular: 200 – 360J

- Monomorphic QRS (Synchronized)
- Polymorphic QRS (Not-Synchronized)

May repeat and increase dose with subsequent cardioversion attempts

**P** Attempt Vagal Maneuvers Procedure  
Only if regular monomorphic complex

#### Consider

#### Only if regular monomorphic complex

Adenosine 6 mg IV / IO

Rapid push with flush

May repeat 12 mg IV / IO

#### Monomorphic QRS:

- All QRS complexes in a single lead are similar in shape.



# Adult Monomorphic Tachycardia

## Wide Complex ( $\geq 0.12$ sec)

### Pearls

- **Recommended Exam:** Mental Status, Skin, Neck, Lung, Heart, Abdomen, Extremities, Neuro
- **Most important goal** is to differentiate the type of tachycardia and if STABLE or UNSTABLE and if SYMPTOMATIC.
- **12-Lead ECG:**
  - 12-Lead ECG is not necessary to diagnose and treat arrhythmia. A single lead ECG is often all that is needed.
  - Obtain 12-Lead when patient is stable and/ or following a rhythm conversion.
- **Monomorphic QRS:**
  - All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
  - QRS complexes in a single lead will change shape from complex to complex.
- **Rhythm** should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.
- **Unstable condition**
  - Condition which acutely impairs vital organ function and cardiac arrest may be impending.
  - If at any point patient becomes unstable move to unstable arm in algorithm.
- **Symptomatic condition**
  - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea but cardiac arrest is not impending.
  - Symptomatic tachycardia usually occurs at rates  $\geq 150$  beats per minute. Patients symptomatic with heart rates  $< 150$  likely have impaired cardiac function such as CHF.
- **Serious Signs/ Symptoms:**
  - Hypotension. Acutely altered mental status. Signs of shock/ poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute congestive heart failure.
- **Search for underlying cause of tachycardia** such as fever, sepsis, dyspnea, etc.
- **Typical sinus tachycardia** is in the range of 100 to (220 – patients age) beats per minute.
- **If patient has history or 12-Lead ECG reveals Wolfe Parkinson White (WPW),** DO NOT administer a Calcium Channel Blocker (e.g., Diltiazem) or Beta Blockers. Use caution with Adenosine and give only with defibrillator available.
- **Regular Wide-Complex Tachycardia:**
  - Unstable condition:**
    - Immediate defibrillation if pulseless and begin CPR.
  - Stable condition:**
    - Typically VT or SVT with aberrancy. Adenosine may be given if regular and monomorphic and if defibrillator available.
    - Verapamil contraindicated in wide-complex tachycardias.
    - Agencies using Amiodarone, Procainamide, and Lidocaine need to choose one agent primarily. Giving multiple anti-arrhythmics requires contact of Medical Control.
    - Atrial arrhythmias with WPW should be treated with Amiodarone or Procainamide
- **Irregular Tachycardia:**
  - Wide-complex, irregular tachycardia: Do not administer calcium channel, beta blockers, or adenosine as this may cause paradoxical increase in ventricular rate. This will usually require cardioversion. Contact Medical Control.
  - Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.



# Adult Polymorphic Tachycardia

## WIDE ( $\geq 0.12$ sec) Torsades de pointes

### History

- Age
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Recent physical exertion
- Palpitations, irregular heart beat
- Time (onset /duration / repetition)

### Signs and Symptoms

- Chest pain, heart failure, dyspnea
- AMS
- Shock, poor perfusion, hypotension
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness

### Differential

- Cardiac arrest
- Sinus Tachycardia vs. dysrhythmia
- Fever, sepsis, infection
- Pericarditis, pulmonary embolism
- Aortic dissection or aneurysm
- Overdose

Assess tachycardia in context of clinical condition  
Identify and treat underlying cause of tachycardia

Unstable / Serious Signs and Symptoms  
HR Typically  $\geq 150$   
Hypotension, Acute AMS, Ischemic Chest Pain, Acute CHF, Seizures, Syncope, or Shock secondary to tachycardia

Defibrillation Procedure

Consider Sedation Prior to Defibrillation

Midazolam 2 – 2.5 mg IV / IO

May repeat as needed

Maximum 10 mg

Wide and Irregular: 200 – 360J

Polymorphic QRS (Not-Synchronized)

*May repeat and increase dose with subsequent cardioversion attempts*

NO

B 12 Lead ECG Procedure

P Cardiac Monitor

IV or IO Access Protocol UP 6

Pulse Present?

NO

Exit to  
Cardiac Arrest  
Protocol AC 3

P Consider consultation with medical control

QT Interval  $< 500$  msec

Amiodarone 150 mg in 100 mL of D5W IV / IO  
Infuse over 10 minutes  
May repeat if tachycardia recurs or persists

Amiodarone 450 mg in 250 mL of D5W  
1 mg/min (33 mL/hr)

Or  
Lidocaine  
1 – 1.5 mg/kg IV / IO

May repeat if refractory  
Lidocaine  
0.75 mg/kg IV / IO  
Maximum 3 mg/kg

Monitor and Reassess

QT Interval  $> 500$  msec

Consider  
Magnesium 2 g IV / IO

May repeat

Maximum 4 g

Monitor and Reassess

### Polymorphic QRS:

- QRS complexes in a single lead will change shape from complex to complex.



Notify Destination or  
Contact Medical Control



AC 8



# Adult Polymorphic Tachycardia

## WIDE ( $\geq 0.12$ sec) Torsades de pointes

### Pearls

- **Recommended Exam:** Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro
- **Most important goal** is to differentiate the type of tachycardia and if STABLE or UNSTABLE and SYMPTOMATIC.
- **12-Lead ECG:**
  - 12 Lead ECG not necessary to diagnose and treat
  - Obtain when patient is stable and/or following rhythm conversion.
- **Monomorphic QRS:**
  - All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
  - QRS complexes in a single lead will change shape from complex to complex.
- **Rhythm** should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.
- **Unstable condition**
  - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
  - If at any point patient becomes unstable move to unstable arm in algorithm.
- **Symptomatic condition**
  - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
  - Symptomatic tachycardia usually occurs at rates  $\geq 150$  beats per minute. Patients symptomatic with heart rates  $< 150$  likely have impaired cardiac function such as CHF.
- **Serious Signs / Symptoms:**
  - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute congestive heart failure.
- **Search for underlying cause of tachycardia** such as fever, sepsis, dyspnea, etc.
- **Typical sinus tachycardia** is in the range of 100 to (220 – patients age) beats per minute.
- **If patient has history or 12 Lead ECG reveals Wolfe Parkinson White (WPW),** DO NOT administer a Calcium Channel Blocker (e.g., Diltiazem) or Beta Blockers. Use caution with Adenosine and give only with defibrillator available.
- **Polymorphic / Irregular Tachycardia:**
  - This situation is usually unstable and immediate defibrillation is warranted.
  - If QT length is known, use for decision-making. Prolonged QT length defined as  $> 500$  msec.
  - QT length  $< 500$  msec:**
    - Arrhythmia more likely related to ischemia or infarction and Magnesium not likely helpful.
    - May quickly deteriorate into Ventricular Fibrillation.
    - Even when terminated by defibrillation, may recur, so follow with medication therapy.
  - QT prolongation  $> 500$  msec:**
    - Magnesium more likely to be helpful.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.

**AC 8**



# Ventricular Fibrillation Pulseless Ventricular Tachycardia (Optional)

## Cardiac Arrest Protocol AC 3

**Begin Continuous CPR Compressions**  
Push Hard ( $\geq$  2 inches) Push Fast (100 - 120 / min)  
Change Compressors every 2 minutes  
(sooner if fatigued)  
(Limit changes / pulse checks  $\leq$  10 seconds)

Ventilate 1 breath every 6 seconds  
30:2 Compression:Ventilation if no Advanced Airway  
Monitor EtCO<sub>2</sub>  
if available

AED Procedure  
if available

### A Defibrillation Procedure

### IV / IO Access Protocol UP 6

**A Epinephrine (1:10,000) 1 mg IV / IO**  
Repeat every 3 to 5 minutes  
If VF / VT refractory to defibrillation, delay Epinephrine administration until after 2nd defibrillation

Search for Reversible Causes

**Continue CPR Compressions**  
Push Hard ( $\geq$  2 inches) Push Fast (100 - 120 / min)  
Change Compressors every 2 minutes  
(sooner if fatigued)  
(Limit changes / pulse checks  $\leq$  10 seconds)

#### If Rhythm Refractory

Continue CPR and give Agency specific Anti-arrhythmics and Epinephrine  
Continue CPR up to point where you are ready to defibrillate with device charged.  
Repeat pattern during resuscitation.

**P Amiodarone 300 mg IV / IO**

May repeat if refractory  
**Amiodarone 150 mg IV / IO**

Or  
**Lidocaine 1.0 – 1.5 mg/kg IV / IO**

May repeat if refractory  
**Lidocaine 0.75 mg/kg IV / IO**

**Maximum 3 mg/kg**

Refractory VF or VT without pulse  
**Magnesium 2 gm IV / IO**

**A Defibrillation Procedure**  
If VF / VT refractory after 3 shocks consider changing vector of defibrillation pads

### Reversible Causes


Hypovolemia  
Hypoxia  
Hydrogen ion (acidosis)  
Hypothermia  
Hypo / Hyperkalemia  
Tension pneumothorax  
Tamponade; cardiac  
Toxins  
Thrombosis; pulmonary (PE)  
Thrombosis; coronary (MI)

### AT ANY TIME

Return of Spontaneous Circulation



Go to Post Resuscitation Protocol AC 10



**T Notify Destination or Contact Medical Control**



# Ventricular Fibrillation Pulseless Ventricular Tachycardia

## Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional Team Focused CPR Protocol AC 11 or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT) compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- **Reassess and document BIAD and / or endotracheal tube placement and EtCO<sub>2</sub> frequently, after every move, and at transfer of care.**
- **IV / IO access and drug delivery is secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation:**  
Follow manufacturer's recommendations concerning defibrillation / cardioversion energy when specified.  
Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.  
Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.  
Manual Defibrillation at the AEMT level is permissible only during pulseless cardiac arrest with VF or VT.
- **End Tidal CO<sub>2</sub> (EtCO<sub>2</sub>)**  
If EtCO<sub>2</sub> is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.  
If EtCO<sub>2</sub> spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Special Considerations**  
**Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm.  
Defibrillation is safe at all energy levels.  
**Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.  
**Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.  
**Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Magnesium Sulfate is not routinely recommended during cardiac arrest, but may help with Torsades de points, prolonged QT, low Magnesium States (malnourished / alcoholic), and suspected digitalis toxicity**
- **Return of spontaneous circulation: Heart rate should be > 60 when initiating anti-arrhythmic infusions.**
- **Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.**
- **Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.**



# Post Resuscitation

## Transport Destination Decision

Post-resuscitation patient is medically complex.

### Consider facility capabilities:

- 24-hour cardiac catheterization laboratory
- Medical ICU service
- Cardiology service
- Neurology service
- Pulmonology service
- Targeted Temperature Management

Arrhythmias are common and usually self limiting after ROSC



If Arrhythmia Persists follow Rhythm Appropriate Protocol

## Return of Spontaneous Circulation

### Repeat Primary Assessment

#### Optimize Ventilation and Oxygenation

- Remove Impedance Threshold Device
- Respiratory Rate 10 / minute
- Maintain SpO<sub>2</sub> 92 – 98%
- **DO NOT HYPERVENTILATE**

**B** • ETCO<sub>2</sub> ideally 35 – 45 mm Hg

Airway  
Protocol(s) AR 1, 2, 3, 4  
*as indicated*

**B** 12 Lead ECG Procedure

IV or IO Access Protocol UP 6

**P** Cardiac Monitor

Monitor Vital Signs / Reassess

Search for reversible causes

### Reversible Causes

Hypovolemia  
Hypoxia  
Hydrogen ion (acidosis)  
Hypothermia  
Hypo / Hyperkalemia

Tension pneumothorax  
Tamponade; cardiac  
Toxins  
Thrombosis; pulmonary (PE)  
Thrombosis; coronary (MI)

### Chest Pain and STEMI

Protocol AC 4  
*if indicated*

Hypotension / Shock  
Protocol AM 5  
*as indicated*

**A** Optimize Systolic BP and Mean Arterial BP

- Systolic BP > 90 mmHg
- Mean Arterial BP > 65 mmHg

Appropriate Arrhythmia  
Protocol(s) AC 2, 6, 7  
*as indicated*

Seizure Protocol UP 13  
*as indicated*

Post Intubation BIAD Management  
Protocol AR 8

Targeted Temperature Management  
Protocol AC 13  
*if available*

**T** Notify Destination or Contact Medical Control **T**



# Post Resuscitation

The initial objectives of post-cardiac arrest care are to:

- Optimize cardiopulmonary function and vital organ perfusion.
- After resuscitation, transport patient to an appropriate hospital with a comprehensive post-cardiac arrest treatment system of care that includes acute coronary interventions, neurological care, goal-directed critical care, and hypothermia.
- Try to identify and treat the precipitating causes of the arrest and prevent recurrent arrest.

Subsequent objectives of post-cardiac arrest care are to:

- Control body temperature to optimize survival and neurological recovery.
- Identify and treat acute coronary syndromes (ACS)
- Optimize mechanical ventilation to minimize lung injury.
- Reduce the risk of multiorgan injury and support organ function if required.

Titrate oxygen to the lowest level required to achieve an arterial oxygen saturation of 94% - 99%, to avoid potential oxygen toxicity.

Hyperventilation or "overbagging" the patient is common after cardiac arrest and should be avoided because of potential adverse hemodynamic effects. Hyperventilation increases intrathoracic pressure and inversely lowers cardiac output. The decrease in PaCO<sub>2</sub> seen with hyperventilation can also potentially decrease cerebral blood flow directly. Ventilation may be started at 10 to 12 breaths per minute and titrated to achieve an EtCO<sub>2</sub> of 35 to 40 mmHg.

The goal of immediate post-cardiac arrest care is to optimize systemic perfusion, restore metabolic homeostasis, and support organ system function to increase the likelihood of intact neurological survival. The post-cardiac arrest period is often marked by hemodynamic instability as well as metabolic abnormalities. Support and treatment of acute myocardial dysfunction and acute myocardial ischemia can increase the probability of survival. Interventions to reduce secondary brain injury, such as therapeutic hypothermia, can improve survival and neurological recovery.

## Pearls

- **Recommended Exam:** Mental Status, Neck, Skin, Lungs, Heart, Abdomen, Extremities, Neuro
- **Continue to search for potential cause of cardiac arrest during post-resuscitation care.**
- **Hyperventilation is a significant cause of hypotension and recurrence of cardiac arrest in the post resuscitation phase and must be avoided. Titrate FiO<sub>2</sub> to maintain SpO<sub>2</sub> of 92 - 98%.**
- **Pain/sedation:**
  - Patients requiring advanced airways and ventilation commonly experience pain and anxiety.
  - Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.
  - Ventilated patients cannot communicate pain / anxiety and providers are poor at recognizing pain / anxiety.
  - Vital signs such as tachycardia and / or hypertension can provide clues to inadequate sedation, however they both are not always reliable indicators of patient's lack of adequate sedation.
  - Pain must be addressed first, before anxiety. Opioids are typically the first line agents before benzodiazepines. Ketamine is also a reasonable first choice agent.
- **Ventilator / Ventilation strategies:**
  - Tailored to individual patient presentations. Medical Control can indicate different strategies above.
  - In general ventilation with BVM should cause chest rise. With mechanical ventilation a reasonable tidal volume should be about 6 mL/kg and peak pressures should be < 30 cmH<sub>2</sub>O.
  - Continuous pulse oximetry and capnography should be maintained during transport for monitoring.
  - Head of bed should be maintained at least 10 – 20 degrees of elevation when possible to decrease aspiration risk.
- **EtCO<sub>2</sub> Monitoring:**
  - Initial End tidal CO<sub>2</sub> may be elevated immediately post-resuscitation, but will usually normalize.
  - Goal is 35 – 45 mmHg but avoid hyperventilation to achieve.
- **Titrate fluid resuscitation and vasopressor administration to maintain SBP of 90 – 100 mmHg or Mean Arterial Pressure (MAP) of 65 – 80 mmHg.**
- **STEMI (ST-Elevation Myocardial Infarction)**
  - Consider placing 2 IV sites in the left arm: Many PCI centers use the right radial artery for intervention.
  - Consider placing defibrillator pads on patient as a precaution.
  - Document and time-stamp facility STEMI notification and make notification as soon as possible.
  - Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).
- **Consider transport to facility capable of managing the post-arrest patient including hypothermia therapy, cardiology / cardiac catheterization, intensive care service, and neurology services.**
- **Targeted Temperature Management (optional):**
  - Maintain core temperature between 32 - 36°C.
  - Infusion of cold saline is NOT recommended in the prehospital setting.
  - No evidence suggests improved survival with prehospital cooling.
- The condition of post-resuscitation patients fluctuates rapidly and continuously, and they require close monitoring. Appropriate post-resuscitation management may best be planned in consultation with Medical Control.



# Team Focused CPR (Optional)

**AT ANY TIME**

Return of Spontaneous Circulation

Go to Post Resuscitation Protocol AC 10

Criteria for Death / No Resuscitation  
Review DNR / MOST Form

YES

NO

**Begin Continuous CPR Compressions**  
Push Hard ( $\geq$  2 inches)  
Push Fast (100 - 120 / min)  
Change Compressors every 2 minutes  
(sooner if fatigued)  
(Limit changes / pulse checks  $\leq$  10 seconds)

Ventilate 1 breath every 6 seconds  
30:2 Compression:Ventilation if no Advanced Airway  
Monitor EtCO<sub>2</sub>  
if available

**First Arriving BLS / ALS Responder**

Initiate Compressions Only CPR  
Initiate Defibrillation Automated Procedure  
if available  
Call for additional resources


**Second Arriving BLS / ALS Responder**

Assume Compressions or  
Initiate Defibrillation Automated / Manual Procedure  
Place BIAD  
DO NOT Interrupt Compressions  
Ventilate at 6 to 8 breaths per minute

Decomposition  
Rigor mortis  
Dependent lividity  
Blunt force trauma  
Injury incompatible with life  
Extended downtime with asystole

Do not begin resuscitation

Follow Deceased Subjects Policy



**Establish Team Leader**  
(Hierarchy)  
Fire Department or Squad Officer  
EMT

First Arriving Responder

**Rotate with Compressor**  
To prevent Fatigue and effect high quality compressions  
Take direction from Team Leader

**Fourth / Subsequent Arriving Responders**  
Take direction from Team Leader

Continue Cardiac Arrest Protocol AC 3

**Establish Team Leader**  
(Hierarchy)  
EMS ALS Personnel  
Fire Department or Squad Officer  
EMT

First Arriving Responder

Initiate Defibrillation Automated Procedure  
Establish IV / IO Protocol UP 6  
Administer Appropriate Medications  
Establish Airway with BIAD if not in place

Initiate Defibrillation Manual Procedure  
Continuous Cardiac Monitoring  
Establish IV / IO Protocol UP 6  
Administer Appropriate Medications  
Establish Airway with BIAD if not in place

Continue Cardiac Arrest Protocol AC 3

**Team Leader**  
ALS Personnel  
Responsible for patient care  
Responsible for briefing / counseling family

**Incident Commander**  
Fire Department / First Responder Officer  
Team Leader until ALS arrival  
Manages Scene / Bystanders  
Ensures high-quality compressions  
Ensures frequent compressor change  
Responsible for briefing family prior to ALS arrival



## Team Focused CPR (Optional)


### Pearls

- This protocol is optional and given only as an example. Agencies may and are encouraged to develop their own.
- Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.
- Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.
- DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT), compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.
- Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.
- Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.
- Reassess and document BIAD and / or endotracheal tube placement and EtCO<sub>2</sub> frequently, after every move, and at transfer of care.
- IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.
- IV access is preferred route. Follow IV or IO Access Protocol UP 6.
- **Defibrillation:** Follow manufacturer's recommendations concerning defibrillation / cardioversion energy when specified.  
    Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.  
    Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
- Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.

**AC 11**



# On Scene Resuscitation Termination of CPR (Optional)





# On Scene Resuscitation Termination of CPR

## Pearls

- **General approach:**
  1. Determine if a terminal disease is involved?
  2. Is there an advanced directive such as a DNR / MOST form?
  3. Did the patient express to your historian any desires regarding resuscitation and if so what measures?
  4. Remember a living will is not a DNR.
- Obtain a history while resuscitation efforts are ongoing. Determine the most legitimate person on scene as your information source such as a spouse, child, or sibling or Durable Health Care Power of Attorney.
- Basic and Advanced Life Support may use for treatment decisions.

## AC 12



# Target Temperature Management (Optional)

## History

- Non-traumatic cardiac arrests (drownings and hanging / asphyxiation are permissible in this protocol.)
- All presenting rhythms are permissible in this protocol
- Age 18 or greater

## Signs and Symptoms

- Cardiac arrest
- Return of Spontaneous Circulation post-cardiac arrest

## Differential

- Continue to address specific differentials associated with the arrhythmia

Return of Spontaneous Circulation ROSC

Criteria for Induced Hypothermia  
Initial rectal temperature  $\geq 93.2^{\circ}\text{F}$  ( $34^{\circ}\text{C}$ )

YES

Agencies utilizing cerebral cooling devices are unlikely to see a change in rectal temperature during transport.

Continued temperature assessment not warranted with these devices. Document initial temperature

NO

Exit to  
Post Resuscitation Protocol AC 10

<b>B</b>	<b>Advanced Airway (includes BIAD) in place with <math>\text{EtCO}_2 &gt; 20 \text{ mmHg}</math></b>
	Airway Protocol(s) AR 1, 2, 3 <b>as indicated</b>
	Post Resuscitation Protocol AC 910 <b>as indicated</b>
	IV / IO Access Protocol UP 6
	Hypotension / Shock Protocol AM 5 <b>as indicated</b>
	Perform Neurological Assessment
	Expose and apply ice packs to axilla and groin areas

Stop cooling measures  
Until temperature increases

Reassess temperature every 10 minutes

Continue Post Resuscitation Care

Agency Specific Cooling Device

$< 89.6^{\circ}\text{F}$  ( $32^{\circ}\text{C}$ )

Reassess Rectal Temperature  
**Target:  $89.6 - 96.8^{\circ}\text{F}$  (Range 32 - 36C)**

$\geq 89.6^{\circ}\text{F}$  ( $32^{\circ}\text{C}$ )

Continue Cooling

Exit to  
Post Resuscitation Protocol AC 9

Shivering noted

YES

**P**  
Fentanyl 50 - 75 mcg IV / IO  
Repeat every 5 minutes as needed  
**Maximum 200 mg**

Versed 2 - 2.5 mg IV / IO  
Repeat every 5 minutes as needed  
**Maximum 10 mg**

Vecuronium 10 mg IV / IO  
If shivering uncontrolled following Opioid and Benzodiazepine Administration

Notify Destination or Contact Medical Control



# Target Temperature Management (Optional)

## Pearls

- **Criteria for Targeted Temperature Management:**
  - Return of spontaneous circulation not related to blunt / penetrating trauma or hemorrhage with ventricular fibrillation / tachycardia and non-shockable arrhythmias.
  - Temperature greater than 93.2°F (34° C).
  - Advanced airway (including BIAD) in place with no purposeful response to verbal commands.
  - Infusion of cold saline is NOT recommended in the prehospital setting.
- **Hyperventilation is a significant cause of hypotension and recurrence of cardiac arrest in the post resuscitation phase and must be avoided. Titrate FiO<sub>2</sub> to maintain SpO<sub>2</sub> of 92 - 98%.**
- **Pain/sedation:**
  - Patients requiring advanced airways and ventilation commonly experience pain and anxiety.
  - Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.
  - Ventilated patients cannot communicate pain / anxiety and providers are poor at recognizing pain / anxiety.
  - Vital signs such as tachycardia and / or hypertension can provide clues to inadequate sedation, however they both are not always reliable indicators of patient's lack of adequate sedation.
  - Pain must be addressed first, before anxiety. Opioids are typically the first line agents before benzodiazepines. Ketamine is also a reasonable first choice agent.
- **EtCO<sub>2</sub> Monitoring:**
  - Initial End tidal CO<sub>2</sub> may be elevated immediately post-resuscitation, but will usually normalize.
  - Goal is 35 – 45 mmHg but avoid hyperventilation to achieve.
- **Titrate fluid resuscitation and vasopressor administration to maintain SBP of > 90 mmHg or Mean Arterial Pressure (MAP) of 65 mmHg.**
- **STEMI (ST-Elevation Myocardial Infarction)**
  - Consider placing 2 IV sites in the left arm: Many PCI centers use the right radial artery for intervention.
  - Consider placing defibrillator pads on patient as a precaution.
  - Document and time-stamp facility STEMI notification and make notification as soon as possible.
  - Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).
- **Consider transport to facility capable of managing the post-arrest patient including hypothermia therapy, cardiology / cardiac catheterization, intensive care service, and neurology services.**
- **Utilization of this protocol mandates transport to facility capable of managing the post-arrest patient and continuation of induced hypothermia therapy.**
- If no advanced airway in place obtained, cooling may only be initiated on order from medical control.
- No evidence suggests improved survival with prehospital cooling.
- The condition of post-resuscitation patients fluctuates rapidly and continuously, and they require close monitoring. Appropriate post-resuscitation management may best be planned in consultation with Medical Control.



# Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD

## History

- SAMPLE
- Bridge to transplant
- Destination therapy
- Estimated downtime
- LVAD, RVAD, Bi-Vad, TAH
- DNR, MOST, or Living Will
- Contact with LVAD coordinator

## Signs and Symptoms

- Unconsciousness
- Pulseless
- Apneic
- Poor capillary refill / skin color
- AMS or decreased mental status
- No electrical activity on ECG
- No heart tones on auscultation

## Differential

- See Reversible Causes below
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage

## Contact VAD coordinator:

- As quickly as possible for troubleshooting and treatment advice, but do not delay emergency treatment
- Follow patient specific emergency plan if present

## Rapid assessment

Check for signs of life  
Assess for adequate perfusion

## Criteria for Death / No Resuscitation

Review DNR / MOST Form

NO

Decomposition

Rigor mortis

Dependent lividity

Blunt force trauma

Injury incompatible with life

Extended downtime with asystole

Do not begin resuscitation

Follow  
Deceased Subjects  
Policy

Unresponsive and  
Not breathing normally

Responsive or Unresponsive and  
Breathing normally

Assess LVAD function  
Look and listen for alarms  
LVAD Alarming?

Assess LVAD function  
Look and listen for alarms  
LVAD Alarming?

NO

Place stethoscope over heart

Place stethoscope over heart

Humming sound present?

Humming sound present?

NO

Airway Protocol(s) AR 1, 2, 3  
if indicated

Respiratory Distress Protocol AR 4  
if indicated

Assume VAD failure  
Initiate age appropriate ventilation rate

Go to Page 2

Assume VAD failure  
Initiate age appropriate ventilation rate

Go to Page 2



# Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD

## History


- SAMPLE
- Bridge to transplant
- Destination therapy
- Estimated downtime
- LVAD, RVAD, Bi-Vad, TAH
- DNR, MOST, or Living Will
- Contact with LVAD coordinator

## Signs and Symptoms

- Unconsciousness
- Pulseless
- Apneic
- Poor capillary refill / skin color
- AMS or decreased mental status
- No electrical activity on ECG
- No heart tones on auscultation


## Differential

- See Reversible Causes below
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage





# Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD





# Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD

## Pearls

- **Recommended exam: Mental status, skin color, capillary refill, peripheral pulses, blood pressure.**
- **Assessment of blood flow and perfusion status:**
  - Optimal BP attained by manual BP and Doppler.
  - Automated BP devices can measure a BP in about 50% of attempts and is not reliable to assess perfusion
  - A MAP of  $\geq 60$  mmHg is adequate for most LVAD patients.
- **Skin color, skin temperature, capillary refill**
- **Mechanical Circulatory Support devices:**
  - LVAD – Left Ventricular Assist Device
  - RVAD – Right Ventricular Assist Device
  - BiVAD – Biventricular Ventricular Assist Device
  - TAH – Total Artificial Heart
- **Reasons for use:**
  - Bridge therapy – patients awaiting transplant or anticipated recovery.
  - Destination therapy – advanced heart failure, not candidate for transplant, and will live rest of life with device.
- **Pump type and assessing pulses:**
  - Pulsatile flow pumps – older units, not commonly in use now, but generate blood flow with a pulsatile flow and patient will have a palpable pulse.
  - Continuous flow pumps – majority of pumps now used and create blood flow in a continuous stream, no pulsatile flow, so patient will not have a palpable pulse.
  - Most devices are implanted inside the chest and have an internal pump, a driveline connected from the pump to the controller unit, and a power source consisting of batteries and electrical cord for receptacles.
- **Common complications:**
  - Disconnection of power supply, either battery disconnect, or electrical cord to receptacle disconnection.
  - Driveline failure or disconnection from controller unit.
  - Controller failure
  - Blood clot formation, acute stroke, and bleeding (mucosal and gastrointestinal most common sites)
  - Infection
- **Abnormal heart rhythm:**
  - Pseudo-PEA: Normal cardiac electrical activity in a patient who is alert and well perfused with no palpable pulse.
  - Tachyarrhythmias are usually well tolerated.
- **End Tidal CO<sub>2</sub> (EtCO<sub>2</sub>)**
  - If EtCO<sub>2</sub> is  $< 10$  mmHg, improve chest compressions. Goal is  $\geq 20$  mmHg.
  - If EtCO<sub>2</sub> spikes, typically  $> 40$  mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Transcutaneous Pacing:**
  - Pacing is NOT effective in cardiac arrest and pacing in cardiac arrest does NOT increase chance of survival



# Total Artificial Heart

## History

- SAMPLE
- Bridge to transplant
- Destination therapy
- Estimated downtime
- LVAD, RVAD, Bi-Vad, TAH
- DNR, MOST, or Living Will
- Contact with LVAD coordinator

## Signs and Symptoms

- Unconsciousness
- Pulseless
- Apneic
- Poor capillary refill / skin color
- AMS or decreased mental status
- No electrical activity on ECG
- No heart tones on auscultation

## Differential

- See Reversible Causes below
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage

Rapid assessment  
Check for signs of life

Assess for adequate perfusion

### DO NOT USE ECG MONITOR

- Total Artificial Heart does not generate ECG

### Contact transplant coordinator:

- As quickly as possible for troubleshooting and treatment advice, but do not delay emergency treatment
- Follow patient specific emergency plan if present

Criteria for Death / No Resuscitation  
Review DNR / MOST Form

Decomposition, Rigor mortis, Dependent lividity, Blunt force trauma  
Injury incompatible with life  
Extended downtime  
Do not begin resuscitation  
Follow Deceased Subjects Policy

NO

Pulse Present?

NO → Go to Page 2

YES

Airway Protocol(s) AR 1, 2, 3  
*if indicated*

Respiratory Distress Protocol AR 4  
*if indicated*

Altered Mental Status Protocol UP 4  
*if indicated*

Check Blood Pressure

Systolic BP  
≥ 150 mmHg

Systolic BP < 150 mmHg  
And  
≥ 90 mmHg

Systolic BP  
< 90 mmHg

**A** Nitroglycerin 0.3 / 0.4 mg SL  
Repeat every 5 minutes  
as needed

Maintain SBP ≥ 90 mmHg

**P** Furosemide 40 mg IV / IO  
*if available*

May assist patient taking their  
antihypertensive medication

Maintain SBP ≥ 90 mmHg

### DO NOT USE:

- Manual or mechanical chest compressions
- ECG/Defibrillation/Pacing/AED devices
- Vasopressor medications
- Antiarrhythmic medications

IV / IO Access Protocol  
UP 6

**A** Normal Saline Bolus 500 mL  
IV / IO  
May repeat as needed  
Maximum 1 L




Notify Destination or  
Contact Medical Control





# Total Artificial Heart





# Total Artificial Heart

## Pearls

- **Recommended exam:** Mental status, skin color, capillary refill, peripheral pulses, blood pressure.
- **Assessment of blood flow and perfusion status:**
  - Manual and automated BP devices can measure a BP.
  - Skin color, skin temperature, capillary refill
- **ECG and telemetry monitoring:**
  - The artificial heart does not produce an ECG wave form or tracing.
  - Do not use the 12-Lead ECG or ECG monitoring as it will only show asystole.
- **Total Artificial Heart:**
  - Different than Ventricular Assist Device (LVAD, RVAD, or Bi-VAD)
  - The patient's left and right ventricles are removed and the artificial heart is connected to the right and left atria.
  - The patient is totally dependent on the artificial heart for circulatory support – the native heart is removed.
  - There are both a right and left side pump, driven by air, and each side driven by a separate driveline.
  - The drivelines are not electric, they are driven by air, so kinking can disrupt the pumping action.
  - Artificial heart produces a pulsatile wave form so the patient will have a palpable pulse when operational.
- **Reasons for use:**
  - Bridge therapy – patients awaiting transplant or anticipated recovery.
  - Destination therapy – advanced heart failure, not candidate for transplant, and will live rest of life with device.
- **Common complications:**
  - Most common is kinking or bending of the driveline(s) which stops air from moving and stops pumping action.
  - Disconnection of power supply, either battery disconnect, or electrical cord to receptacle disconnection.
  - Driveline failure or disconnection from controller unit.
  - Controller failure
  - Blood clot formation, acute stroke, and bleeding (mucosal and gastrointestinal most common sites)
  - Infection
- **Blood pressure:**
  - Optimal SBP is < 130 mmHg and > 90 mmHg.
  - Hypertension puts great strain on the pump and can cause blood to back up into the lungs and cause pulmonary edema and respiratory failure.
  - Epinephrine and vasopressors are ineffective, can cause hypertension, and may worsen the patient's condition.
- **Manual or mechanical chest compressions:**
  - Do not use**
- **End Tidal CO<sub>2</sub> (EtCO<sub>2</sub>)**
  - Helpful in monitoring adequate perfusion status.
- **Defibrillation/Cardioversion:**
  - Do not use.
- **Transcutaneous Pacing:**
  - Do not use.



# Wearable Cardioverter Defibrillator Vest

## History


- SAMPLE
- Known risk for Sudden Cardiac Death
- Risk for life-threatening arrhythmia
- No implanted defibrillator
- Heart failure – cardiomyopathy
- Decreased ejection fraction

## Signs and Symptoms

- Chest pain, dyspnea
- Palpitations
- Received shock from vest
- Poor capillary refill / skin color
- AMS or decreased mental status

## Differential

- See Reversible Causes below
- Arrhythmia
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage





# Wearable Cardioverter Defibrillator Vest



## Pearls

- **Recommended exam:** Mental status, skin color, capillary refill, peripheral pulses, blood pressure.
- **Wearable Cardioverter-Defibrillator Vest:**
  - Device is preparing to delivery a shock to the patient:**  
Before device delivers a shock, it tests to see if patient is conscious – voice prompt instructs patient to press the “response” button (see diagram above).
  - Only the patient should press the “response” button.**
  - Once a treatable arrhythmia is detected it takes between 25 and 60 seconds to deliver the shock.**
- **Audible and tactile warning system:**  
The device will provide a vibration, a siren tone, and voice prompts to check if the patient is conscious and give them an opportunity to press the “response” button to abort a shock.  
See audible warning system above.
- **Reasons for use:**  
Currently only device on the market is the Zoll LifeVest.  
Worn by patients at risk of sudden cardiac arrest or risk of abnormal and/or lethal arrhythmia.
- **Blue gel on the patient's skin from the device:**  
Electrode pads release a blue gel prior to treatment to improve shock conduction and reduce burning.  
Do not remove the gel if the vest is left in place during treatment.  
Remove gel if vest is removed for prehospital care.
- **Shock to providers:**  
Do not touch the patient when the device is instructing you that a shock will be delivered.  
Providers can be shocked by the device during energy delivery if provider is touching the patient.
- **Removing the device for prehospital care:**  
The device should only be removed when ECG monitor and defibrillator is available.  
Continuous ECG monitoring and electrode pads should be in place when vest is removed.
- **Defibrillation/cardioversion with vest in place:**  
Disconnect the device from the vest before you deliver a cardioversion or defibrillation
- **Transcutaneous Pacing:**  
May be utilized with vest in place – disconnect the device from the vest before you perform transcutaneous pacing.



# Allergic Reaction/ Anaphylaxis

## History

- Onset and location
- Insect sting or bite
- Food allergy / exposure
- Medication allergy / exposure
- New clothing, soap, detergent
- Past history of reactions
- Past medical history
- Medication history

## Signs and Symptoms

- Itching or hives
- Coughing / wheezing or respiratory distress
- Chest or throat constriction
- Difficulty swallowing
- Hypotension or shock
- Edema
- N/V

## Differential

- Urticaria (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration / Airway obstruction
- Vasovagal event
- Asthma or COPD
- CHF

Assess Symptom Severity / Suspected Exposure to Allergen

### MILD Skin Only

### Moderate 2 + Body Systems

### SEVERE

2 + Body Systems + hypotension  
Or Isolated Hypotension

Diphenhydramine  
25 - 50 mg PO

IV or IO Access Protocol UP 6  
**if indicated**

Diphenhydramine  
25 - 50 mg  
PO / IV / IM / IO

Histamine (H2) Blocker  
*If available*

**B**  
Monitor and Reassess  
Monitor for Worsening  
Signs and Symptoms

Epinephrine 1:1000 IM  
0.3 – 0.5 mg

Repeat every 5 minutes  
*if no improvement*

Diphenhydramine  
25 - 50 mg PO  
*See Pearls*

Albuterol Nebulizer  
2.5 – 5 mg  
Repeat as needed x 3  
*if indicated*

Epinephrine 1:1000  
0.3 – 0.5 mg IM  
Repeat every 5 minutes  
*if no improvement*

Diphenhydramine  
25 - 50 mg IV / IM / IO  
*if not given PO (See Pearls)*

Epinephrine 1:1000 IM  
0.3 – 0.5 mg

Repeat every 5 minutes  
*if no improvement*

Albuterol 2.5 – 5 mg  
Nebulizer  
Repeat as needed x 3  
**if indicated**

Epinephrine 1:1000  
0.3 – 0.5 mg IM  
Repeat every 5 minutes  
*if no improvement*

Airway Protocol(s) AR 1 - 4  
**if indicated**

Hypotension/ Shock  
Protocol AM 5  
**if indicated**

IV or IO Access Protocol UP 6

Albuterol Nebulizer  
2.5 – 5 mg

+/- Ipratropium 0.5 mg (DuoNeb)  
Repeat as needed x 3  
*if indicated*

Histamine (H2) Blocker  
*If available*

Normal Saline Bolus  
500 mL IV / IO  
Repeat as needed  
Maximum 2 Liters

Methylprednisolone  
125 mg IV / IO

No improvement with IM Epinephrine  
Epinephrine IV / IO  
*See Pearls for dosing regimen*

Notify Destination or  
Contact Medical Control



# Allergic Reaction/ Anaphylaxis

Epinephrine 1:100,000 infusion may be needed for severe allergic reaction unresolved by initial doses of epi IM/IV. Mix 1mg Epi in 1000ml of NS – Administer 1 mL per minute (severe unresolved after initial treatment or infusion – may need 1mg Epi in 1000mL of NS over 10 minutes).

Cardiac monitoring is indicated for moderate and severe reactions

## Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdominal**
- **Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.**
- **Epinephrine and administration:**
  - Drug of choice and the FIRST drug that should be administered in acute anaphylaxis (Moderate / Severe Symptoms.)
  - IM Epinephrine should be administered in priority before or during attempts at IV or IO access.
- **Diphenhydramine and steroid administration:**
  - Diphenhydramine/ steroids have no proven benefit in Moderate/ Severe anaphylaxis.
  - Diphenhydramine/ steroids should NOT delay initial or repeat Epinephrine administration.
  - In Moderate and Severe anaphylaxis, Diphenhydramine may decrease mental status.
  - Diphenhydramine should NOT be given to a patient with decreased mental status and/ or a hypotensive patient as this may cause nausea, vomiting, and/ or worsening mental status.
- **Anaphylaxis unresponsive to repeat doses of IM epinephrine may require IV epinephrine administration by IV push or epinephrine infusion. Contact Medical Control for appropriate dosing.**
- **Symptom Severity Classification:**
  - Mild symptoms:**
    - Flushing, hives, itching, erythema with normal blood pressure and perfusion.
  - Moderate symptoms:**
    - Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with normal blood pressure and perfusion.
  - Severe symptoms:**
    - Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with hypotension/ poor perfusion or isolated hypotension.
- **Allergic reactions may occur with only respiratory and gastrointestinal symptoms and have no rash/ skin involvement.**
- **Angioedema** is seen in moderate to severe reactions and is swelling involving the face, lips or airway structures. This can also be seen in patients taking blood pressure medications like Prinivil / Zestril (lisinopril)-typically end in -il.
- **Hereditary Angioedema** involves swelling of the face, lips, airway structures, extremities, and may cause moderate to severe abdominal pain. Some patients are prescribed specific medications to aid in reversal of swelling.
  - Paramedic may assist or administer this medication per patient/ package instructions.**
- **Patients with moderate and severe reactions should receive a 12 lead ECG and should be continually monitored, but this should NOT delay administration of epinephrine.**
- **EMR/ EMT:**
  - The use of Epinephrine IM is limited to the treatment of anaphylaxis and may be given only by autoinjector, unless manual draw-up is approved by the Agency Medical Director and the NC office of EMS.**
  - Administration of diphenhydramine is limited to the oral route only.**
- **EMT administration of beta-agonist is limited to only patients currently prescribed the medication, unless approved by the Agency Medical Director and the NC office of EMS.**
- **Agency Medical Director may require contact of medical control prior to EMT/ EMR administering any medication(s).**
- **The shorter the onset from exposure to symptoms the more severe the reaction.**



# Diabetic; Adult

## History

- Past medical history
- Medications
- Recent blood glucose check
- Last meal

## Signs and Symptoms

- Altered mental status
- Combative / irritable
- Diaphoresis
- Seizures
- Abdominal pain
- Nausea / vomiting
- Weakness
- Dehydration
- Deep / rapid breathing

## Differential

- Alcohol / drug use
- Toxic ingestion
- Trauma; head injury
- Seizure
- CVA
- Altered baseline mental status

	Blood Glucose Analysis Procedure
<b>B</b>	12 Lead ECG Procedure <i>if indicated</i>
	IV or IO Access Protocol UP 6
<b>P</b>	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Hypotension/ Shock Protocol AM 5 <i>if indicated</i>
	Suspected Stroke Protocol AM 7 <i>if indicated</i>
	Seizure Protocol UP 13 <i>if indicated</i>

<b>B</b>	Blood Glucose $\leq$ 69 mg / dl and symptomatic No venous access <b>Glucagon 1 – 2 mg IM</b> Repeat in 15 minutes if needed
----------	--

Blood Sugar  
 $\leq$  69 mg / dl

Blood Sugar  
70 – 249 mg / dl

Blood Sugar  
 $\geq$  250 mg / dl

Consider Oral Glucose Solution  
*if available*

Consider Oral Solution (Juices / Food)

If no improvement

**A** D10 125 mL – 250 mL IV / IO  
Or  
D50 12.5 - 25 gm IV / IO  
May repeat until Blood Glucose  $\geq$  80 mg/dL

Blood Glucose Analysis Procedure  
*if condition changes*

**A** Normal Saline Bolus  
500 mL IV / IO  
May repeat as needed  
Then infuse 150 mL / hr

Exit to  
Appropriate  
Protocol(s)

Monitor and Reassess  
Every 5 minutes  
Until Blood Glucose  $\geq$  80 mg / dl

Notify Destination or  
Contact Medical Control



# Diabetic; Adult

If using D50% IV or IO – consider placing the D50% in to 500ml of fluid for administration of the first 12.5g, and repeat if necessary.

Because diabetics tend to incur a disproportionate incidence of cardiac problems, consider ECG evaluation. If correction of dehydration, hypoglycemia, hyperglycemia, or acidosis fails to abolish a cardiac dysrhythmia, refer to appropriate protocol.

Supplemental dextrose provides the best means for reversing hypoglycemia. For the hypoglycemic who merely displays confusion and has the ability to swallow and obey commands, oral ingestion of food, drink, or instant glucose is the preferred treatment. For any lethargic, stuporous, or comatose hypoglycemic patient or for one who is unable to swallow and to understand or obey your commands, who would possibly aspirate oral glucose, intravenous/intraosseous or rectal administration of dextrose is indicated.

## Pearls

- **Recommended exam: Mental Status, Skin, Respirations and effort, Neuro.**
- **Patients with prolonged hypoglycemia or those who are malnourished may not respond to glucagon.**
- **Do not administer oral glucose to patients who are not able to swallow or protect their airway.**
- **Quality control checks should be maintained per manufacturers recommendation for all glucometers.**
- **Patient's refusing transport to medical facility after treatment of hypoglycemia:**

Blood sugar must be  $\geq 80$ , patient has ability to eat and availability of food with responders on scene.

Blood sugar trending up is necessary with 2 or more readings above 80, obtained over 15-minute period.

Patient must have known history of diabetes and not taking any oral diabetic agents.

Patient returns to normal mental status and has a normal neurological exam with no new neurological deficits.

Must demonstrate capacity to make informed health care decisions. See Universal Patient Care Protocol UP-1.

Otherwise contact medical control.

- **Hypoglycemia with Oral Agents:**

Patient's taking oral diabetic medications should be encouraged to allow transportation to a medical facility.

They are at risk of recurrent hypoglycemia that can be delayed for hours and require close monitoring even after normal blood glucose is established.

Not all oral agents have prolonged action so Contact Medical Control or NC Poison Control Center for advice.

Patient's who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.

- **Hypoglycemia with Insulin Agents:**

Many forms of insulin now exist. Longer acting insulin places the patient at risk of recurrent hypoglycemia even after a normal blood glucose is established.

Not all insulins have prolonged action so Contact Medical Control for advice.

Patient's who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.

- **Congestive Heart Failure patients who have Blood Glucose  $> 250$ :**

Limit fluid boluses unless patient has signs of volume depletion such as, dehydration, poor perfusion, hypotension, and/ or shock.

- In extreme circumstances with no IV / IO access and no response to glucagon, D50 can be administered rectally, Contact Medical Control for advice.



# Dialysis/ Renal Failure

## History


- Peritoneal or Hemodialysis
- Anemia
- Catheter access noted
- Shunt access noted
- Hyperkalemia

## Signs and Symptoms

- Hypotension
- Bleeding
- Fever
- Electrolyte imbalance
- Nausea and / or vomiting
- Altered Mental Status
- Seizure
- Arrhythmia

## Differential

- Congestive heart failure
- Pericarditis
- Diabetic emergency
- Sepsis
- Cardiac tamponade





# Dialysis/ Renal Failure

## Pearls

- **Recommended exam: Mental status. Neurological. Lungs. Heart. Skin.**
- **Preferably transport to a medical facility capable of providing dialysis treatment.**
- **Do not take Blood Pressure or start IV / IO in extremity which has a shunt/ fistula in place.**
- **Access of shunt indicated in the dead or near-dead patient only with no IV or IO access.**
- **If hemorrhage cannot be controlled with firm, uninterrupted direct pressure, application of tourniquet with uncontrolled dialysis fistula bleeding is indicated.**
- **Hemodialysis:**  
Process which removes waste from the blood stream and occurs about three times each week.  
Some patients do perform hemodialysis at home.
- **Peritoneal dialysis:**  
If patient complains of fever, abdominal pain, and/ or back pain, bring the Peritoneal Dialysis fluid bag, which has drained from the abdomen, to the hospital.

## Complications of Dialysis Treatment:

### Hypotension:

Typically responds to small fluid bolus of 250 mL Normal Saline.

May result in angina, AMS, seizure or arrhythmia.

### Filtration and decreased blood levels of some medications like some seizure medications:

### Disequilibrium syndrome:

Shift of metabolic waste and electrolytes causing weakness, dizziness, nausea and/ or vomiting and seizures.

### Equipment malfunction:

Air embolism.

Bleeding.

Electrolyte imbalance.

Fever.

- **Fever:**

Consider sepsis in a dialysis patient with any catheter extending outside the body.

- Always consider Hyperkalemia in all dialysis or renal failure patients.
- Sodium Bicarbonate and Calcium Chloride/ Gluconate should not be mixed. Ideally give in separate lines.
- Renal dialysis patients have numerous medical problems typically. Hypertension and cardiac disease are prevalent.



# Hypertension

## History

- Documented Hypertension
- Related diseases: Diabetes; CVA; Renal Failure; Cardiac Problems
- Medications for Hypertension
- Compliance with Hypertensive Medications
- Erectile Dysfunction medications
- Pregnancy

## Signs and Symptoms

### One of these

- Systolic BP 220 or greater
- Diastolic BP 120 or greater


### AND at least one of these

- Headache
- Chest Pain
- Dyspnea
- Altered Mental Status
- Seizure

## Differential

- Hypertensive encephalopathy
- Primary CNS Injury
- Cushing's Response with Bradycardia and Hypertension
- Myocardial Infarction
- Aortic Dissection / Aneurysm
- Pre-eclampsia / Eclampsia

Hypertension is not uncommon especially in an emergency setting. Hypertension is usually transient and in response to stress and/ or pain. A hypertensive emergency is based on blood pressure along with symptoms which suggest an organ is suffering damage such as MI, CVA or renal failure. This is very difficult to determine in the pre-hospital setting in most cases. Aggressive treatment of hypertension can result in harm. Most patients, even with significant elevation in blood pressure, need only supportive care. Specific complaints such as chest pain, dyspnea, pulmonary edema or altered mental status should be treated based on specific protocols and consultation with Medical Control.



## Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- Elevated blood pressure is based on two to three sets of vital signs.
- Symptomatic hypertension is typically revealed through end organ dysfunction to the cardiac, CNS, or renal systems.
- All symptomatic patients with hypertension should be transported with their head elevated at 30 degrees.
- Ensure appropriate size blood pressure cuff utilized for body habitus.



# Hypotension/ Shock

## History

- Blood loss - vaginal or gastrointestinal bleeding, AAA, ectopic
- Fluid loss - vomiting, diarrhea, fever
- Infection
- Cardiac ischemia (MI, CHF)
- Medications
- Allergic reaction
- Pregnancy
- History of poor oral intake

## Signs and Symptoms

- Restlessness, confusion
- Weakness, dizziness
- Weak, rapid pulse
- Pale, cool, clammy skin
- Delayed capillary refill
- Hypotension
- Coffee-ground emesis
- Tarry stools

## Differential

- Ectopic pregnancy
- Dysrhythmias
- Pulmonary embolus
- Tension pneumothorax
- Medication effect / overdose
- Vasovagal
- Physiologic (pregnancy)
- Sepsis

	Blood Glucose Analysis Procedure
<b>B</b>	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
<b>P</b>	Cardiac Monitor
	Airway Protocol(s) <i>if indicated</i>
	Diabetic Protocol AM 2 <i>if indicated</i>

History and Exam Suggest Type of Shock

Cardiogenic

Hypovolemic

Distributive

Obstructive

Chest Pain: Cardiac and STEMI  
Protocol AC 4  
 Appropriate Cardiac Protocol(s)  
*if indicated*

Allergy Protocol AM 1  
*if indicated*

Suspected Sepsis  
Protocol UP 15  
*if indicated*

Multiple Trauma  
Protocol TB 6  
*if indicated*

**P** Chest Decompression-Needle Procedure  
*if indicated*

**A** Normal Saline  
Bolus 500 mL IV  
Repeat to effect  
SBP > 90  
2 L Maximum

**P** Vasopressor(s)

Notify Destination or Contact Medical Control

**AM 5**



# Hypotension/ Shock

## Septic Shock

Skin – Varies from flushed pink (if fever is present) to pale and cyanotic, possible petechia, possible purple blotches, possible peeling (general or at palms and soles), red streaks progressing proximally

Blood Pressure – Early: cardiac output increases but toxins may cause loss of peripheral vascular resistance

Blood Pressure – Late: Hypotension; precipitous fall in blood pressure

Respiration – Dyspnea with altered lung sounds

Other – Possible high fever (except some elderly and very young patients),

Other – Late: frank pulmonary edema

Septic shock is the result of an overwhelming infection. Sepsis may not be noticed for some time and may be confused with a wide variety of other conditions. Sepsis begins with an infection that sets in motion an overwhelming systemic response from the immune system, the end results of which are hypotension, hypoperfusion, and end organ dysfunction. The infection can be caused by bacteria, fungus, and some viruses. Gramnegative bacteremia is more likely to cause sepsis (50 percent of infections) compared to gram-positive bacteremia (25 percent of infections). The inflammatory and cellular events are complex and significant. Organisms invade the body through the bloodstream or locally. In either case, the organisms release structural components (commonly referred to as toxins, either endotoxins or exotoxins) that trigger our natural immune system to release its own endogenous mediators (e.g., cytokines from monocytes and prostaglandins from neutrophils, along with histamine, heparin, tumor necrosis factor [TNF], and others).

Adrenal crisis, also known as Addisonian crisis and acute adrenal insufficiency, is a medical emergency and potentially life-threatening situation requiring immediate emergency treatment. It is a constellation of symptoms that indicate severe adrenal insufficiency caused by insufficient levels of the hormone cortisol. The signs and symptoms may include:

- Pain in the lower back, flank, abdomen, or legs
- Severe vomiting and diarrhea, leading to dehydration
- Low blood pressure
- Loss of consciousness
- High potassium (hyperkalemia) and low sodium (hyponatremia)

## Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Hypotension is defined as a systolic blood pressure less than 90. This is not always reliable and should be interpreted in context and consider patient's typical BP if known.**
- **Shock may be present with a normal blood pressure initially or even elevated blood pressure.**
- **Shock is often present with normal vital signs and may develop insidiously. Tachycardia may be the first and only sign.**
- **Consider all possible causes of shock and treat per appropriate protocol.**
- **Hypovolemic Shock:**

Hemorrhage, trauma, GI bleeding, ruptured aortic aneurysm or pregnancy-related bleeding.

### Tranexamic Acid (TXA):

Agencies utilizing TXA must submit letters from their receiving trauma centers for approval by the OEMS Medical Director.

Receiving trauma centers must agree to continue TXA therapy with repeat dosing.

TXA is NOT indicated and should NOT be administered where trauma occurred > 3 hours prior to EMS arrival.

### Cardiogenic Shock:

Heart failure: MI, Cardiomyopathy, Myocardial contusion, Ruptured ventricle / septum / valve / toxins.

### Distributive Shock:

#### Sepsis/ Anaphylactic/ Neurogenic/ Toxins

Hallmark is warm, dry, pink skin with normal capillary refill time and typically alert.

### Obstructive Shock:

Pericardial tamponade. Pulmonary embolus. Tension pneumothorax.

Signs may include hypotension with distended neck veins, tachycardia, unilateral decreased breath sounds or muffled heart sounds.

### Acute Adrenal Insufficiency or Congenital Adrenal Hyperplasia:

Body cannot produce enough steroids (glucocorticoids/ mineralocorticoids.)

May have primary or secondary adrenal disease, congenital adrenal hyperplasia, or more commonly have stopped a steroid like prednisone. Injury or illness may precipitate.


Usually hypotensive with nausea, vomiting, dehydration and/ or abdominal pain.

If suspected, AEMT or Paramedic should give Methylprednisolone 125 mg IM / IV / IO or Dexamethasone 10 mg IM / IV / IO. Use steroid agent specific to your drug list.

May administer prescribed steroid carried by patient IM / IV / IO. Patient may have Hydrocortisone (Cortef or Solu-Cortef). Dose: < 1y.o. give 25 mg, 1-12 y.o. give 50 mg, and > 12 y.o. give 100 mg or dose specified by patient's physician.



# Suspected Stroke: Activase/ t-PA or Tenecteplase/ TNK Interfacility Transfer (Optional)





# Suspected Stroke: Activase/ t-PA or Tenecteplase/ TNK Interfacility Transfer (Optional)

## Pearls

- **Recommended Exam:** Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro
- **This protocol is optional. Agencies may develop their own in conjunction with their regional stroke center(s) guidance.**
- **This protocol is intended for interfacility transfer patients only. Medication must be started at initial treating hospital.**
- **Items in Red Text are key performance measures used in protocol compliance.**
- **The Reperfusion Checklist** should be completed for any suspected stroke patient.
- **Time of Onset or Last Seen Normal:**

One of the most important items the pre-hospital provider can obtain, of which all treatment decisions are based.

Be very precise in gathering data to establish the time of onset and report as an actual time (i.e. 13:47 NOT "about 45 minutes ago.")

Without this information patient may not be able to receive thrombolytics at facility.

Wake up stroke: Time starts when patient last awake or symptom free.

- **Time of Symptom Discovery:**

Time when symptoms of stroke are first noticed by patient, bystanders, witnesses, or family/ caregivers.

- The differential listed on the Altered Mental Status Protocol should also be considered.
- Be alert for airway problems (swallowing difficulty, vomiting/aspiration).
- Hypoglycemia can present as a localized neurologic deficit, especially in the elderly.

### **Infusion Pump Alarm / No Flow:**

Remove drip chamber from Activase / t-PA bag.

Spike Activase/ t-PA drip chamber to NS bag.

Restart infusion to complete medication remaining in IV tubing.

- **Medication dosing safety:**

When IV Activase/ t-PA dose administration will continue en route, verify estimated time of completion.

Verify with sending hospital that excess Activase/ t-PA has been withdrawn from the bottle and wasted.

This ensures the bottle will be empty when the full dose is finished. *For example, if the total dose is 70 mg, then 30 cc should be withdrawn and wasted since a 100 mg bottle of Activase/ t-PA contains 100 mL of fluid when reconstituted.*

Sending hospital should apply a label to Activase/ t-PA bottle with the number of mL of fluid that should be in the bottle in case of pump failure during transit.

**Tenecteplase/ TNK** is given as a single weight-based bolus dose and does not involve an infusion. Medication should be administered prior to patient transport.

- **Allergy Anaphylaxis:**

Activase/ t-PA, is structurally identical to endogenous t-PA and therefore should not induce allergy, single cases of acute hypersensitivity reactions have been reported.

### **Angioedema:**

Rapid swelling (edema) of the dermis, subcutaneous tissue, mucosa and submucosal tissues. Typically involves the face, lips, tongue and neck.

Almost always self limiting but may progress to interfere with airway / breathing so close monitoring is warranted.

**Tenecteplase/ TNK** can also cause hypersensitivity reactions and should be treated based on AM 1 and PM 1.

Utilize the Allergy / Anaphylaxis Protocol as indicated and also for angioedema. Infusion should be stopped.

Give all medications related to the Allergy/ Anaphylaxis Protocol by IV route only as patient should remain NPO.



# Childbirth/ Labor

## History


- Due date
- Time contractions started / how often
- Rupture of membranes
- Time / amount of any vaginal bleeding
- Sensation of fetal activity
- Past medical and delivery history
- Medications
- Gravida / Para Status
- High Risk pregnancy

## Signs and Symptoms

- Spasmodic pain
- Vaginal discharge or bleeding
- Crowning or urge to push
- Meconium

## Differential

- Abnormal presentation  
Buttock  
Foot  
Hand
- Prolapsed cord
- Placenta previa
- Abruptio placenta





# Childbirth/ Labor

Apgar score			
	Score 2	Score 1	Score 0
<b>A</b> pppearance			
<b>P</b> ulse	>100 bpm	<100 bpm	No pulse
<b>G</b> rimace	Cries and pulls away	Grimaces or weak cry	No response to stimulation
<b>A</b> ctivity			
<b>R</b> espiration	Strong cry	Slow, irregular	No breathing

## Pearls

- **Recommended Exam (of Mother): Mental Status, Heart, Lungs, Abdomen, Neuro**
- **Record APGAR at 1 minute and 5 minutes after birth. Do not delay resuscitation to obtain APGAR.**
- **If neonate requiring resuscitation, move quickly to AO 2 Newly Born Protocol**
- **After delivery, massaging the uterus (lower abdomen) will promote uterine contraction and help to control post-partum bleeding (apply uterine massage only after placenta delivery).**
- **Postpartum hemorrhage:**

### Pitocin (Oxytocin):

Following field delivery, where available, administer 10 IU IM to promote uterine contraction and decrease postpartum hemorrhage.

Agencies may administer via IV or IO route per local agency medical director.

### Tranexamic Acid (TXA):

Administer when postpartum hemorrhage is associated with signs and symptoms of shock. CONTRAINDICATED where birth occurs > 3 hours prior to EMS arrival.

- **Transport or Delivery?**

Decision to transport versus remain and deliver is multifactorial and difficult. Generally it is preferable to transport.

Factors that will impact decision include: number of previous deliveries; length of previous labors; frequency of contractions; urge to push; and presence of crowning.

- **Maternal positioning for labor:**

Supine with head flat or elevated per mother's choice. Maintain flexion of both knees and hips. Elevated buttocks slightly with towel. If delivery not imminent, place mother in the left, lateral recumbent position with right side up about 10 – 20°.

- **Umbilical cord clamping and cutting:**

Place first clamp about 10 cm from infant's abdomen and second clamp about 5 cm away from first clamp.

- **Multiple Births:**

Twins occur about 1/90 births. Typically manage the same as single gestation. If imminent delivery call for additional resources, if needed. Most twins deliver at about 34 weeks so lower birth weight and hypothermia are common. Twins may share a placenta so clamp and cut umbilical cord after first delivery. Notify receiving facility immediately.

- Document all times (Contraction onset, contraction duration and frequency, delivery, APGAR 1 and 2, and placenta delivery).
- If maternal seizures occur, refer to the Obstetrical Emergencies Protocol.
- Some perineal bleeding is normal with any childbirth. Large quantities of blood or free bleeding are abnormal.

## AO 1



# Newly Born

## History

- Due date and gestational age
- Multiple gestation (twins etc.)
- Meconium / Delivery difficulties
- Congenital disease
- Medications (maternal)
- Maternal risk factors such as substance abuse or smoking


## Signs and Symptoms

- Respiratory distress
- Peripheral cyanosis or mottling (normal)
- Central cyanosis (abnormal)
- Altered level of responsiveness
- Bradycardia

## Differential

- Airway failure, Secretions, or Respiratory drive
- Infection
- Maternal medication effect
- Hypovolemia, Hypoglycemia, Hypothermia
- Congenital heart disease

In a non-vigorous infant whose respirations are not improving after warming, drying, and stimulating within 30 seconds, move quickly to Positive Pressure Ventilation with BVM





# Newly Born

## Pearls

- **Recommended Exam: Quality of Cry, Muscle tone, Respirations, Heart Rate, Pulse Oximetry, and Gestational Age**
- **Majority of newborns do not require resuscitation, only warming, drying, stimulating, and cord clamping.**
  - With term gestation, strong cry/ breathing, and good muscle tone, generally will not need resuscitation.
  - If no resuscitation needed, skin-to-skin contact with the mother is best way to maintain warmth of infant.
  - Maintain warmth of infant following delivery adjuncts; cap/ hat, plastic wrap, thermal mattress, radiant heat.
  - Most important vital signs in the newly born are heart rate, respirations, and respiratory effort.
  - About 10% of newborns need assistance to help them start breathing after birth.
  - About 1% of newborns require intensive resuscitation to restore/ support cardiorespiratory functions.
- **Airway:**
  - Positive Pressure Ventilations with BVM is the most important treatment in a newborn with poor respirations and/ or persistent bradycardia (HR < 100 BPM).
  - When BVM is needed, ventilation rate is 40 – 60 breaths per minute.
  - Adequacy of ventilation/ is measured mainly by increase in heart rate as well as chest rise.
  - If heart rate or respirations are not improving after 30 to 60 seconds of resuscitation, place BIAD or endotracheal tube.
  - Routine suctioning is no longer recommended, bulb suction only if needed.
- **Breathing:**
  - Oxygen is not necessary initially, but if infant is not responding with increased heart rate or adequate breathing, add oxygen to the BVM.
- **Circulation/ Compressions:**
  - Heart rate is critical during first few moments of life and is best monitored by 3 or 4 lead ECG, as pulse assessment is difficult in the neonate. Heart Rate is best tool for gauging resuscitation success.
  - If heart rate remains < 60 BPM after 30 to 60 seconds of BVM/ resuscitation, begin compressions.
  - With BIAD or ETT in place, compressions and ventilation should be coordinated with compression, compression, compression, then ventilation. (3:1 ratio with all events totaling 120 per minute)
  - 2-thumbs encircling chest and supporting the back is recommended. Limit interruptions of chest compressions.
- **If infant not responding to BVM, compressions, and/ or epinephrine, consider hypovolemia, pneumothorax, and/ or hypoglycemia (< 40 mg/dL).**
- **Document 1 and 5 minute APGAR in PCR or ePCR. DO NOT delay or interrupt resuscitation to obtain an APGAR score.**
- **Meconium staining:**
  - Infant born through meconium staining who is NOT vigorous:**
    - Bulb suction mouth and nose and provide positive pressure ventilation.
    - Direct endotracheal suctioning is no longer recommended.
- **Expected Pulse Oximetry readings following birth:**
  - (Accurate only in infant NOT requiring resuscitation)**

1 minute	60 – 65%
2 minutes	65 – 70%
3 minutes	70 – 75%
4 minutes	75 – 80%
5 minutes	80 – 85%
10 minutes	85 – 95%
- Pulse oximetry should be applied to the right upper arm, wrist, or palm.
- **Cord clamping:**
  - Recommended to delay for 1 minute, unless infant requires resuscitation.
- Maternal sedation or narcotics will sedate infant (Naloxone NO LONGER recommended, use supportive care only).
- D10 = D50 diluted (1 ml of D50 with 4 ml of Normal Saline) or **D10 solution at 2 mL/kg IV / IO.**
- **In the NEONATE, D10 is administered at 2 mL/kg. (NOT 5 mL/kg in the pediatric patient after the first month of life.)**

Apgar score			
	Score 2	Score 1	Score 0
<b>A</b> pppearance			
<b>P</b> ulse	> 100 bpm	< 100 bpm	No pulse
<b>G</b> rimace	Cries and pulls away	Grimaces or weak cry	No response to stimulation
<b>A</b> ctivity			
<b>R</b> espiration	Strong cry	Slow, irregular	No breathing



# OB-GYN Emergency

## History


- Past medical history
- Hypertension meds
- Prenatal care
- Prior pregnancies / births
- Gravida / Para

## Signs and Symptoms

- Vaginal bleeding
- Abdominal pain
- Seizures
- Hypertension
- Severe headache
- Visual changes
- Edema of hands and face

## Differential

- Pre-eclampsia / Eclampsia
- Placenta previa
- Placenta abruptio
- Spontaneous abortion





# OB-GYN Emergency

## Pearls

- **Recommended Exam: Mental Status, Abdomen, Heart, Lungs, Neuro**
- **With active seizure activity, benzodiazepine is a priority over magnesium sulfate.**
- **Midazolam 5 – 10 mg IM is effective in termination of seizures. Do not delay IM administration with difficult or no IV or IO access.**
- **Magnesium Sulfate should be administered as quickly as possible. May cause hypotension and decreased respiratory drive, but more likely in doses higher than 6 gm.**
- Any pregnant patient involved in a MVC should be seen immediately by a physician for evaluation. Greater than 20 weeks generally require 4 to 6 hours of fetal monitoring. DO NOT suggest the patient needs an ultrasound but emphasize patient needs 4 to 6 hours of fetal monitoring.
- After delivery, massaging the uterus (lower abdomen) will promote uterine contraction and help to control post-partum bleeding (apply uterine massage only after placenta delivery).
- **Postpartum or Vaginal hemorrhage:**

### **Pitocin (Oxytocin):**

Following field delivery, where available, administer 10 IU IM to promote uterine contraction and decrease postpartum hemorrhage.

Agencies may administer via IV or IO route per local agency medical director.

### **Tranexamic Acid (TXA):**

Administer when postpartum hemorrhage is associated with signs and symptoms of shock.

CONTRAINdICATED where birth occurs > 3 hours prior to EMS arrival.

Vaginal hemorrhage unrelated to pregnancy, administer with signs and symptoms of shock.

- **Ectopic pregnancy:**

Implantation of fertilized egg outside the uterus, commonly in or on the fallopian tube. As fetus grows, rupture may occur. Vaginal bleeding may or may not be present. Many women with ectopic pregnancy do not know they are pregnant. Usually occurs within 5 to 10 weeks of implantation. Maintain high index of suspicion with women of childbearing age experiencing abdominal pain.

- **Preeclampsia:**

Occurs in about 6% of pregnancies. Defined by hypertension and protein in the urine. RUQ pain, epigastric pain, N/V, visual disturbances, headache, and hyperreflexia are common symptoms.

In the setting of pregnancy, hypertension is defined as a BP > 140 systolic or > 90 diastolic mmHg, or a relative increase of 30 systolic and 20 diastolic from the patient's normal (pre-pregnancy) blood pressure.

Risk factors: < 20 years of age, first pregnancy, multi-gestational pregnancy, gestational diabetes, obesity, personal or family history of gestational hypertension.

- **Eclampsia:**


Seizures occurring in the context of preeclampsia. Remember, women may not have been diagnosed with preeclampsia.

- Maintain patient in a left lateral position, right side up 10 - 20° to minimize risk of supine hypotensive syndrome.
- Ask patient to quantify bleeding - number of pads used per hour.

## AO 3



# Adult Airway





# Adult Airway

## Pearls

- See Pearls section of protocols AR 2 and 3.
- For the purposes of this protocol a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of  $\geq 90\%$ , it is acceptable to continue with basic airway measures.
- Ventilation rate should be 10 - 12 per minute to maintain a EtCO<sub>2</sub> of 35 – 45 and avoid hyperventilation.
- Anticipating the Difficult Airway and Airway Assessment

**Difficult BVM Ventilation (ROMAN):** Radiation treatment/ Restriction; **Obese/ Obstruction/ OB** – 2d and 3d trimesters/ Obstructive sleep apnea; Mask seal difficulty (hair, secretions, trauma); **Age  $\geq 55$ ; No teeth.**

**Difficult Laryngoscopy (LEON):** Look externally for anatomical problems; Evaluate 3-3-2 (Mouth opening should equal 3 of patients finger's width, mental area to neck should equal 3 of patient's finger's width, base of chin to thyroid prominence should equal 2 of patients finger's width); **Obese, obstruction, OB – 2d and 3d trimesters; Neck mobility limited.**

**Difficulty BIAD (RODS):** Radiation treatment/ Restriction; **Obese/ Obstruction/ OB** – 2d and 3d trimesters/ Obstructive sleep apnea; Distorted or disrupted airway; **Short thyromental distance/ Small mandible.**

**Difficulty Cricothyrotomy / Surgical Airway (SMART):** Surgery scars; **Mass or hematoma, Access or anatomical problems; Radiation treatment to face, neck, or chest; Tumor.**

- Complete an Airway Evaluation Form with any BIAD or Intubation procedure where medications are used to facilitate.
- Nasotracheal intubation:

Procedure requires spontaneous breathing and may require considerable time, exposing patient to critical desaturation.

Contraindicated in combative, anatomically disrupted or distorted airways, increased ICP, severe facial trauma, basal skull fracture, and head injury. Orotracheal route is preferred.

- Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
- If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment).
- AEMT and Paramedics should consider using a BIAD if oral-tracheal intubation is unsuccessful.
- During intubation attempts use External Laryngeal Manipulation to improve view of glottis.
- Gastric tube placement should be considered in all intubated patients if available or time allows.
- It is important to secure the endotracheal tube well to better maintain ETT placement. Manual stabilization of endotracheal tube should be used during all patient moves / transfers.
- **DOPE:** Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.



# Adult, Failed Airway

## Definition of Failed Airway:

Unable to Ventilate and Oxygenate  $\geq 90\%$  during or after one (1) or more unsuccessful intubation attempts

and/ or

Anatomy inconsistent with continued attempts

and/ or

Three (3) unsuccessful attempts by most experienced Paramedic/AEMT.


*Each attempt should include change in approach or equipment*

NO MORE THAN THREE (3) ATTEMPTS TOTAL

### Capnography Monitoring

- End-tidal (EtCO<sub>2</sub>) monitoring is mandatory following placement of an endotracheal tube.
- EtCO<sub>2</sub> monitoring is mandatory following placement of a BIAD once available on scene.

Protocols AR 1, 2, and 3 should be utilized together (even if agency is not using Drug Assisted Airway as they contain useful information for airway management.





# Adult, Failed Airway

A failed airway occurs when a provider begins a course of airway management by endotracheal intubation and identifies that intubation by that means will not succeed.

## Conditions which define a Failed Airway:

1. Failure to maintain adequate oxygen saturation 90% or greater after 2 or more failed intubation attempts.
2. Three (3) failed attempts at intubation by the most experienced prehospital provider on scene even when adequate oxygen saturation of 90% or greater can be maintained.
3. Unable to maintain adequate oxygen saturation 90% or greater with BM ventilation techniques and insufficient time to attempt alternative maneuvers. A patient near death or dying.

**The most important way to avoid a failed airway is to identify patients with expected difficult airway, difficult BM ventilations, difficult BIAD, difficult laryngoscopy and / or difficult cricothyrotomy.**

Please refer to Adult Airway Protocol for more information on how to identify the patient with the potential difficult airway.

## Position of patient:

In the field setting improper position of the patient and rescuer are responsible for many failed and difficult intubations. Often this is dictated by uncontrolled conditions present at the scene and we must adapt. However many times the rescuer does not optimize the patient and rescuer position. The sniffing position or the head simply extended upon the neck are probably the best positions. The goal is to align the ear canal with the suprasternal notch in a straight line.

In the obese or late pregnant patient, elevating the torso by placing blankets, pillows or towels will optimize the position. This can be facilitated by raising the head of the cot.

## Use of cot in optimal patient / rescuer position:

The cot can be elevated and lowered to facilitate intubation. With the patient on the cot raise until the patient's nose is at the level of your umbilicus which will place you at the optimal position

## Pearls

- **For the purposes of this protocol a secure airway is when the patient is receiving appropriate oxygenation and ventilation.**
- **If an effective airway is being maintained by BVM with continuous pulse oximetry values of  $\geq 90\%$ , it is acceptable to continue with basic airway measures.**
- **Ventilation rate should be 10 - 12 per minute to maintain a EtCO<sub>2</sub> of 35-45 and avoid hyperventilation.**
- **Anticipating the Difficult Airway and Airway Assessment**

**Difficult BVM Ventilation (ROMAN):** Radiation treatment/ Restriction; **Obese/ Obstruction/ OB** – 2d and 3d trimesters/ Obstructive sleep apnea; Mask seal difficulty (hair, secretions, trauma); **Age  $\geq 55$ ; No teeth.**

**Difficult Laryngoscopy (LEON):** Look externally for anatomical problems; Evaluate 3-3-2 (Mouth opening should equal 3 of patients finger's width, mental area to neck should equal 3 of patient's finger's width, base of chin to thyroid prominence should equal 2 of patients finger's width); **Obese, obstruction, OB – 2d and 3d trimesters; Neck mobility limited.**

**Difficulty BIAD (RODS):** Radiation treatment/ Restriction; **Obese/ Obstruction/ OB** – 2d and 3d trimesters/ Obstructive sleep apnea; Distorted or disrupted airway; **Short thyromental distance/ Small mandible.**

**Difficulty Cricothyrotomy / Surgical Airway (SMART):** Surgery scars; **Mass or hematoma, Access or anatomical problems; Radiation treatment to face, neck, or chest; Tumor**

- **Complete an Airway Evaluation Form with any BIAD or Intubation procedure where medications are used to facilitate.**
- **Nasotracheal intubation:**

Procedure requires spontaneous breathing and may require considerable time, exposing patient to critical desaturation.

Contraindicated in combative, anatomically disrupted or distorted airways, increased ICP, severe facial trauma, basal skull fracture, and head injury. Orotracheal route is preferred.

- **Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.**
- **If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)**
- AEMT and Paramedics should consider using a BIAD if oral-tracheal intubation is unsuccessful.
- During intubation attempts use External Laryngeal Manipulation to improve view of glottis.
- Gastric tube placement should be considered in all intubated patients if available or time allows.
- It is important to secure the endotracheal tube well to better maintain ETT placement. Manual stabilization of endotracheal tube should be used during all patient moves/ transfers.
- **DOPE:** Displaced tracheostomy tube/ ETT, **Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.**



# Airway, Drug Assisted (OPTIONAL)


**Indications for Drug Assisted Airway**

- Failure to protect the airway and/or
- Unable to oxygenate and/or
- Unable to ventilate and/or
- Impending airway compromise

## Capnography Monitoring

- End-tidal (EtCO<sub>2</sub>) monitoring is mandatory following placement of an endotracheal tube.
- EtCO<sub>2</sub> monitoring is mandatory following placement of a BIAD once available on scene.

Protocols AR 1, 2, 3, 5, and 6 should be utilized together (even if agency is not using Drug Assisted Airway Protocol) as they contain useful information for airway management.





# Airway, Drug Assisted (OPTIONAL)

## Pearls

- Agencies must maintain a separate Performance Improvement Program specific to Drug Assisted Airway.
- This procedure requires at least 2 Paramedics. See Pearls section of protocols AR 1 and 2.
- For the purposes of this protocol, a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of  $\geq 90\%$ , it is acceptable to continue with basic airway measures.
- **Ventilation rate:**  
30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 10 - 12 per minute.  
Maintain EtCO<sub>2</sub> between 35 - 45 and avoid hyperventilation.
- **Hypoxia and/ or Hypotension:**  
Increased risk of cardiac arrest when a sedative with paralytic medications are administered while hypoxic and/ or hypotensive.  
Resuscitation and correction of hypoxia and/ or hypotension are paramount prior to use of these combined agents.  
Ketamine administration allows time for appropriate resuscitation of hypoxia and/ or hypotension while managing the airway.
- **PEDIATRIC POPULATION (fit within a Pediatric Medication/ Skill Resuscitation System product,  $\leq 15$  years of age, or  $\leq 49$  kg):**  
This protocol may be used in the pediatric population ONLY with real time, direct online medical control by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR.  
Agencies using Ketamine in the pediatric population must also be using in their adult population.
- **KETAMINE:**  
Ketamine may be used with or without a paralytic agent in conjunction with either an OPA, NPA, BIAD, or endotracheal tube.  
(BIAD is preferred over endotracheal tube until hypoxia and/ or hypotension are corrected).  
Ketamine may be used during the resuscitation of hypoxia or hypotension in conjunction with airway management. Once hypoxia and hypotension are corrected, use of a sedative and paralytic can proceed if indicated.  
Ketamine may be used in the dangerously combative patient requiring airway management IM, IV, IO should be established as soon as possible.  
Ketamine may be used for sedation once a BIAD or endotracheal tube are established and confirmed.  
Agencies using Ketamine must follow Standards Policy: Medical Policy Section Ketamine Program Requirements. Medical Policy 2.
- Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
- If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)
- **NC EMS Airway Evaluation Form:**  
Fully complete and have receiving healthcare provider sign confirming BIAD or endotracheal tube placement.  
Complete online in region specific ReadyOp and upload completed form.  
Complete when Ketamine, Etomidate, Succinylcholine and/ or Rocuronium or used to facilitate use of a BIAD and/ or endotracheal intubation.
- Paramedics/ AEMT should consider using a BIAD if endotracheal intubation is unsuccessful.
- Drug Assisted Airway is not recommended in an urban setting (short transport) when able to maintain oxygen saturation  $\geq 90\%$ .
- **DOPE:** Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.



# Adult COPD/ Asthma Respiratory Distress

## History


- Asthma; COPD -- chronic bronchitis, emphysema, congestive heart failure
- Home treatment (oxygen, nebulizer)
- Medications (theophylline, steroids, inhalers)
- Toxic exposure, smoke inhalation

## Signs and Symptoms

- Shortness of breath
- Pursed lip breathing
- Decreased ability to speak
- Increased respiratory rate and effort
- Wheezing, rhonchi
- Use of accessory muscles
- Fever, cough
- Tachycardia

## Differential

- Asthma
- Anaphylaxis
- Aspiration
- COPD (Emphysema, Bronchitis)
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pneumothorax
- Cardiac (MI or CHF)
- Pericardial tamponade
- Hyperventilation
- Inhaled toxin (Carbon monoxide, etc.)





# Adult COPD/ Asthma Respiratory Distress


## Pearls

- **Recommended Exam:** Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro
- **Items in Red Text** are key performance measures used to evaluate protocol compliance and care.
- **This protocol includes all patients with respiratory distress, COPD, Asthma, Reactive Airway Disease, or bronchospasm.**
- **Patients may also have wheezing and respiratory distress with viral upper respiratory tract infections and pneumonia.**
- **Pulse oximetry should be monitored continuously and consider End-tidal CO<sub>2</sub> monitoring if available.**
- **Combination nebulizers containing albuterol and ipratropium (DuoNeb):**
  - Patients may require more than 3 nebulizer treatments, treatments should continue until improvement.
  - Following 3 combination nebulizers (DuoNeb), it is preferable to continue albuterol solely with subsequent treatments as there is no proven benefit to continual use of ipratropium.
- **Epinephrine:**
  - If allergic reaction or anaphylaxis is suspected, give immediately and repeat until improvement.
  - If allergic reaction is not suspected, administer with failure to improve and/ or impending respiratory failure.
- **Consider Magnesium Sulfate with no improvement and/ or impending respiratory failure. Likely more effective with asthmatic exacerbation and less so with COPD exacerbation.**
- **Non-Invasive Positive Pressure Ventilation (NIPPV: CPAP or Bi-Level/ BiPap):**
  - May be used with COPD, Asthma, Allergic reactions, and/ or CHF.
  - Consider early in treatment course.
  - Consider removal if SBP remains < 100 mmHg and not responding to other treatments.
- In patients using levalbuterol (Xopenex) you may use Albuterol for the first treatment then use the patient's supply for repeat nebulizers or agency's supply.
- A silent chest in respiratory distress is a pre-respiratory arrest sign.
- **EMR/ EMT:**
  - The use of Epinephrine IM is limited to the treatment of anaphylaxis and may be given only by autoinjector, unless manual draw-up is approved by the Agency Medical Director and the NC office of EMS.
  - Administration of diphenhydramine is limited to the oral route only.
- **EMT administration of beta-agonist is limited to only patients currently prescribed the medication**, unless approved by the Agency Medical Director and the NC office of EMS.
- Agency Medical Director may require contact of medical control prior to EMT/ EMR administering any medication(s).

**AR 4**



# Pediatric Airway





# Pediatric Airway

## Pearls

**This protocol is for use in patients who FIT within a Pediatric Medication/ Skill Resuscitation System Product.**

- For the purposes of this protocol, a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of  $\geq 90\%$ , it is acceptable to continue with basic airway measures.
- **Ventilation rate:**  
30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 10 - 12 per minute. Maintain EtCO<sub>2</sub> between 35 - 45 and avoid hyperventilation.
- **Ketamine for airway intervention and/ or sedation purposes:**  
Ketamine may be used in pediatric patients (fit within a Pediatric Medication/Skill Resuscitation System product,  $\leq 15$  years of age, or  $\leq 49$  kg) with DIRECT ONLINE MEDICAL ORDER by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR only.  
Agencies using Ketamine in the pediatric population must also be using in their adult population.
- **KETAMINE:**  
Ketamine may be used with or without a paralytic agent in conjunction with either an OPA, NPA, BIAD or endotracheal tube. BIAD is preferred over endotracheal tube until hypoxia and/ or hypotension are corrected.  
Ketamine may be used during the resuscitation of hypoxia or hypotension in conjunction with airway management. Once hypoxia and hypotension are corrected, use of a sedative and paralytic can proceed if indicated.  
Ketamine may be used in the dangerously combative patient requiring airway management IM, IV, IO should be established as soon as possible.  
Ketamine may be used for sedation once a BIAD or endotracheal tube are established and confirmed.  
Agencies using Ketamine must follow Standards Policy: Medical Policy Section Ketamine Program Requirements. Medical Policy 2.
- **Intubation:**  
Attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.  
Use of a stylet is recommended in all pediatric intubations.  
Endotracheal tube: Depth = 3 x the diameter of the ETT. Estimated Size =  $16 + \text{age (years)} / 4$ . Term newborn = 3.5 mm.  
If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)
- **NC EMS Airway Evaluation Form:**  
Fully complete and have receiving healthcare provider sign confirming BIAD or endotracheal tube placement.  
Complete online in region specific ReadyOp and upload completed form.  
Complete when Ketamine, Etomidate, Succinylcholine and/ or Rocuronium or used to facilitate use of a BIAD and/ or endotracheal intubation. Paramedics/ AEMT should consider using a BIAD if endotracheal intubation is unsuccessful.
- Secure the endotracheal tube well and consider c-collar in pediatric patients (even in absence of trauma) to better maintain ETT placement.  
Manual stabilization of endotracheal tube should be used during all patient moves / transfers.
- **Airway Cricothyrotomy Percutaneous Needle Procedure:**  
Indicated as a lifesaving / last resort procedure in pediatric patients  $< 10$  years of age.  
Very little evidence to support its use and safety.  
A variety of alternative pediatric airway devices now available make the use of this procedure rare.  
Agencies who utilize this procedure must develop a written procedure, establish a training program, maintain equipment and submit procedure and training plan to the State Medical Director/ Regional EMS Office.  
 $\geq 10$  years: Surgical cricothyrotomy or commercial kits based on agency preference recommended.
- **DOPE:** Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.




# Pediatric Failed Airway

## Definition of Failed Airway:

- Unable to Ventilate and Oxygenate  $\geq 90\%$  during or after one (1) or more unsuccessful intubation attempts.  
and/or
- Anatomy inconsistent with continued attempts.  
and/or
- Three (3) unsuccessful attempts by most experienced Paramedic/AEMT.

*Each attempt should include change in approach or equipment*

NO MORE THAN THREE (3) ATTEMPTS TOTAL





# Pediatric Failed Airway

## Pearls

**This protocol is for use in patients who FIT within a Pediatric Medication/ Skill Resuscitation System Product.**

- For the purposes of this protocol, a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of  $\geq 90\%$ , it is acceptable to continue with basic airway measures.
- **Ventilation rate:**  
30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 10 - 12 per minute. Maintain EtCO<sub>2</sub> between 35 - 45 and avoid hyperventilation.
- **Ketamine for airway intervention and/ or sedation purposes:**  
Ketamine may be used in pediatric patients (fit within a Pediatric Medication/Skill Resuscitation System product,  $\leq 15$  years of age, or  $\leq 49$  kg) with DIRECT ONLINE MEDICAL ORDER by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR only.  
Agencies using Ketamine in the pediatric population must also be using in their adult population.
- **KETAMINE:**  
Ketamine may be used with or without a paralytic agent in conjunction with either an OPA, NPA, BIAD or endotracheal tube. BIAD is preferred over endotracheal tube until hypoxia and/ or hypotension are corrected.  
Ketamine may be used during the resuscitation of hypoxia or hypotension in conjunction with airway management. Once hypoxia and hypotension are corrected, use of a sedative and paralytic can proceed if indicated.  
Ketamine may be used in the dangerously combative patient requiring airway management IM, IV, IO should be established as soon as possible.  
Ketamine may be used for sedation once a BIAD or endotracheal tube are established and confirmed.  
Agencies using Ketamine must follow Standards Policy: Medical Policy Section Ketamine Program Requirements. Medical Policy 2.
- **Intubation:**  
Attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.  
Use of a stylet is recommended in all pediatric intubations.  
Endotracheal tube: Depth = 3 x the diameter of the ETT. Estimated Size =  $16 + \text{age (years)} / 4$ . Term newborn = 3.5 mm.  
If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)
- **NC EMS Airway Evaluation Form:**  
Fully complete and have receiving healthcare provider sign confirming BIAD or endotracheal tube placement.  
Complete online in region specific ReadyOp and upload completed form.  
Complete when Ketamine, Etomidate, Succinylcholine and/ or Rocuronium or used to facilitate use of a BIAD and/ or endotracheal intubation. Paramedics/ AEMT should consider using a BIAD if endotracheal intubation is unsuccessful.
- Secure the endotracheal tube well and consider c-collar in pediatric patients (even in absence of trauma) to better maintain ETT placement.  
Manual stabilization of endotracheal tube should be used during all patient moves / transfers.
- **Airway Cricothyrotomy Percutaneous Needle Procedure:**  
Indicated as a lifesaving / last resort procedure in pediatric patients  $< 10$  years of age.  
Very little evidence to support its use and safety.  
A variety of alternative pediatric airway devices now available make the use of this procedure rare.  
Agencies who utilize this procedure must develop a written procedure, establish a training program, maintain equipment and submit procedure and training plan to the State Medical Director/ Regional EMS Office.  
 $\geq 10$  years: Surgical cricothyrotomy or commercial kits based on agency preference recommended.
- **DOPE:** Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.



# Pediatric Asthma Respiratory Distress

## History


- Time of onset
- Possibility of foreign body
- Past Medical History
- Medications
- Fever / Illness
- Sick Contacts
- History of trauma
- History / possibility of choking
- Ingestion / OD
- Congenital heart disease

## Signs and Symptoms

- Wheezing / Stridor / Crackles / Rales
- Nasal Flaring / Retractions / Grunting
- Increased Heart Rate
- AMS
- Anxiety
- Attentiveness / Distractability
- Cyanosis
- Poor feeding
- JVD / Frothy Sputum
- Hypotension

## Differential

- Asthma / Reactive Airway Disease
- Aspiration
- Foreign body
- Upper or lower airway infection
- Congenital heart disease
- OD / Toxic ingestion / CHF
- Anaphylaxis
- Trauma



WHEEZING / Asthma

Following Assessment

STRIDOR / Croup

<b>B</b>	<b>Albuterol Nebulizer 1.25 - 2.5 mg</b> Repeat as needed x 3
	Pediatric Airway Protocol(s) 3, 5 - 7 <i>as indicated</i>
	Albuterol Nebulizer 1.25 - 2.5 mg +/- Ipratropium 0.5 mg (DuoNeb) Repeat as needed x 3 <i>if indicated</i>
	Consider Epinephrine 1:1000 0.01 mg / kg IM  Maximum 0.3 mg
	Methylprednisolone 2 mg / kg IV / IO / IM Maximum 125 mg
	Magnesium Sulfate 40 mg/kg IV / IO Infuse over 10 – 20 minutes  Maximum 2 gm

<b>B</b>	<b>Albuterol Nebulizer 1.25 - 2.5 mg</b> Repeat as needed x 3
	<b>Epinephrine Nebulizer 1 mg (1:1000) in 2 mL NS</b> May repeat x 1
	Pediatric Airway Protocol(s) 3, 5 - 7 <i>as indicated</i>
	Albuterol Nebulizer 1.25 - 2.5 mg +/- Ipratropium 0.5 mg (DuoNeb) Repeat as needed x 3 <i>if indicated</i>
	Consider Epinephrine 1:1000 0.01 mg / kg IM  Maximum 0.3 mg
	Methylprednisolone 2 mg / kg IV / IO / IM Maximum 125 mg

Notify Destination or  
Contact Medical Control



# Pediatric Asthma Respiratory Distress

## Pearls

- Recommended Exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro
- Items in Red Text are key performance measures used to evaluate protocol compliance and care.
- This protocol includes all patients with respiratory distress, Asthma, Reactive Airway Disease, croup, or bronchospasm.
- Patients may also have wheezing and respiratory distress with viral upper respiratory tract infections and pneumonia.
- Pulse oximetry should be monitored continuously and consider End-tidal CO<sub>2</sub> monitoring if available.
- Combination nebulizers containing albuterol and ipratropium (DuoNeb):

Patients may require more than 3 nebulizer treatments, treatments should continue until improvement.

Following 3 combination nebulizers (DuoNeb), it is preferable to continue albuterol solely with subsequent treatments as there is no proven benefit to continual use of ipratropium.

- Epinephrine:
  - If allergic reaction or anaphylaxis is suspected, give immediately and repeat until improvement.
  - If allergic reaction is not suspected, administer with no improvement and/ or impending respiratory failure.
- Consider Magnesium Sulfate with impending respiratory failure and/ or no improvement.
- Consider IV access when Pulse oximetry remains  $\leq 92\%$  after first beta-agonist nebulizer treatment.
- Do not force a child into a position, allow them to assume position of comfort, typically the tripod position.
- Bronchiolitis is a viral infection typically affecting infants which results in wheezing which may not respond to beta-agonists. Consider Epinephrine nebulizer if patient  $< 18$  months and not responding to initial beta-agonist treatment.
- Croup typically affects children  $< 2$  years of age. It is viral, possible fever, gradual onset, no drooling is noted.
- Epiglottitis typically affects children  $> 2$  years of age. It is bacterial, with fever, rapid onset, possible stridor, patient wants to sit up to keep airway open, drooling is common. Airway manipulation may worsen the condition.
- In patients using levalbuterol (Xopenex) you may use Albuterol for the first treatment then use the patient's supply for repeat nebulizers or agency's supply.
- A silent chest in respiratory distress is a pre-respiratory arrest sign.

- EMR/ EMT:

The use of Epinephrine IM is limited to the treatment of anaphylaxis and may be given only by autoinjector, unless manual draw-up is approved by the Agency Medical Director and the NC office of EMS.

Administration of diphenhydramine is limited to the oral route only.

- EMT administration of beta-agonist is limited to only patients currently prescribed the medication, unless approved by the Agency Medical Director and the NC office of EMS.
- Agency Medical Director may require contact of medical control prior to EMT/ EMR administering any medication(s).




# Post-intubation/ BIAD Management

## Capnography Monitoring

- End-tidal (EtCO<sub>2</sub>) monitoring is mandatory following placement of an endotracheal tube.
- EtCO<sub>2</sub> monitoring is mandatory following placement of a BIAD once available on scene.

Protocols AR 1, 2, 3, 5, and 6 should be utilized together (even if agency is not using Drug Assisted Airway Protocol) as they contain useful information for airway management.





# Post-intubation/ BIAD Management

## Pearls

- **Recommended Exam:** Mental Status, HEENT, Heart, Lungs, Neuro
- Patients requiring advanced airways and ventilation commonly experience pain and anxiety.
- Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.
- Ventilated patients cannot communicate pain/ anxiety and providers are poor at recognizing pain/ anxiety.
- Vital signs such as tachycardia and/ or hypertension can provide clues to inadequate sedation, however they are not always reliable indicators of a patient's lack of adequate sedation.
- **Sedation strategy:**
  - Pain is the primary reason patients experience agitation and must be addressed first.
  - Opioids and/ or Ketamine are the first line agents, alone or in combination.
  - Benzodiazepines may be utilized if patient is not responding to adequate opioid and/ or Ketamine doses.
  - Paralysis is considered a last resort, only when patients are not responding to opioids, Ketamine, or benzodiazepines.
  - Patients that have received paralytics may be experiencing pain with no obvious signs or symptoms.
  - Consider sedation early after giving paralytics, especially in patients receiving Rocuronium.
- **Ventilation rate:**
  - Guidelines: 30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 10 – 12 per minute.
  - Maintain EtCO<sub>2</sub> between 35 - 45 and avoid hyperventilation.
- Ventilator/ Ventilation strategies will need to be tailored to individual patient presentations. Medical director can indicate different strategies above.
- **Propofol:**
  - Use restricted to agencies approved by the OEMS State Medical Director.
  - Agencies must submit a use policy and education plan to the OEMS.
  - Infusion must be supplied and initiated by a medical facility and may be used only during interfacility transfer.
  - Paramedic may titrate infusion to maintain appropriate sedation but cannot initiate or bolus the medication.
- In general, ventilation with BVM should cause chest rise. With mechanical ventilation a reasonable tidal volume should be about 6 - 8 mL/kg and peak pressures should be < 30 cmH<sub>2</sub>O. Plateau Pressures should be < 30 cmH<sub>2</sub>O.
- Head of bed should be maintained at least 10 – 20 degrees of elevation when possible, to decrease aspiration risk.
- With abrupt clinical deterioration, if mechanically ventilated, disconnect from ventilator to assess lung compliance.
- **DOPE:** Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.



# Ventilator Emergencies

## History


- Birth defect (tracheal atresia, tracheomalacia, craniofacial abnormalities)
- Surgical complications (damage to phrenic nerve)
- Trauma (post-traumatic brain or spinal cord injury)
- Medical condition (bronchopulmonary dysplasia, muscular dystrophy)

## Signs and Symptoms

- Transport requiring maintenance of a mechanical ventilator
- Power or equipment failure at residence

## Differential

- Disruption of oxygen source
- Dislodged or obstructed tracheostomy tube
- Detached or disrupted ventilator circuit
- Cardiac arrest
- Increased oxygen requirement / demand
- Ventilator failure



## Pearls

- Always talk to family/ caregivers as they have specific knowledge and skills.
- If using the patient's ventilator bring caregiver knowledgeable in ventilator operation during transport.
- Take patient's ventilator to hospital even if not functioning properly.
- Always use patient's equipment if available and functioning properly.
- Continuous pulse oximetry and End Tidal CO<sub>2</sub> monitoring must be utilized during assessment and transport.
- Unable to correct ventilator problem: Remove patient from ventilator and manually ventilate using BVM.
- Typical alarms: Low Pressure/ Apnea: Loose or disconnected circuit, leak in circuit or around tracheostomy site.  
Low Power: Internal battery depleted.  
High Pressure: Plugged/ obstructed airway or circuit.
- DOPE: Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.

**AR 9**



# Tracheostomy Tube Emergencies

## History


- Birth defect (tracheal atresia, tracheomalacia, craniofacial abnormalities)
- Surgical complications (accidental damage to phrenic nerve)
- Trauma (post-traumatic brain or spinal cord injury)
- Medical condition (bronchial or pulmonary dysplasia, muscular dystrophy)

## Signs and Symptoms

- Nasal flaring
- Chest wall retractions (with or without abnormal breath sounds)
- Attempts to cough
- Copious secretions noted coming out of the tube
- Faint breath sounds on both sides of chest despite significant respiratory effort
- AMS
- Cyanosis

## Differential

- Allergic reaction
- Asthma
- Aspiration
- Septicemia
- Foreign body
- Infection
- Congenital heart disease
- Medication or toxin
- Trauma



## Pearls

- Always talk to family/ caregivers as they have specific knowledge and skills.
- Important to ask if patient has undergone laryngectomy. This does not allow mouth/ nasal ventilation by covering stoma.
- Use patients equipment if available and functioning properly.
- Estimate suction catheter size by doubling the inner tracheostomy tube diameter and rounding down.
- Suction depth: Ask family/ caregiver. No more than 3 to 6 cm typically. Instill 2 – 3 mL of NS before suctioning.
- Do not suction more than 10 seconds each attempt and pre-oxygenate before and between attempts.
- DO NOT force suction catheter. If unable to pass, then tracheostomy tube should be changed.
- Always deflate tracheal tube cuff before removal. Continual pulse oximetry and EtCO<sub>2</sub> monitoring if available.
- **DOPE:** Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.



# Mechanical Ventilation; Adult (Optional)

## History


- Multiple etiologies leading to need for advanced airway control
- Requires ventilation support
- Height and underlying lung conditions

## Signs and Symptoms


- Loss of consciousness or AMS with inability to protect airway
- Difficult oxygenation and/or ventilation
- 

## Differential

- ROSC
- Trauma
- Stroke
- Seizure
- Shock (see Shock Protocol)
- Toxicological



<b>MODE:</b> Volume – Assist Control
<b>FiO<sub>2</sub>:</b> 100%
<b>PEEP:</b> 5 cmH <sub>2</sub> O
<b>TIDAL VOLUME (V<sub>t</sub>):</b> 8 mL/kg <i>Follow PBW and V<sub>t</sub> on page 3</i>
<b>BPM: RESPIRATORY RATE:</b> 18 BPM
<b>FLOW RATE:</b> 60 mL/min (preset)
<b>Check Plateau Pressure</b> <b>Press Manual Breath</b> <i>P</i> Pressure button Goal Pressure < 30 cm/H <sub>2</sub> O
<b>Decrease Tidal Volume</b> 1 mL/kg increments Until ≤ 29 cm/H <sub>2</sub> O <b>(DO NOT DECREASE &lt; 4 mL/kg)</b>



### Alarming Ventilator and unsure how to troubleshoot

- Immediately disconnect patient and use BVM.
- Once oxygenation and ventilation stabilized, restart ventilator set-up procedure.

### Home Ventilator Inter-facility Transfer with Ventilator

- Set initial parameters to home or facility settings
- Titrate to oxygenation, work of breathing, SpO<sub>2</sub>, and EtCO<sub>2</sub>.
- Use home ventilator if functioning properly.

<b>MODE:</b> Volume – Assist Control
<b>FiO<sub>2</sub>:</b> 100%
<b>PEEP:</b> 5 cmH <sub>2</sub> O
<b>TIDAL VOLUME:</b> 8 mL/kg <i>Follow PBW and V<sub>t</sub> on page 3</i>
<b>BPM: RESPIRATORY RATE:</b> 12 BPM
<b>FLOW RATE:</b> 60 mL/min (preset)
<b>I:E Ratio</b> Increase to 1:4 or 1:5
<b>Check Plateau Pressure</b> <b>Press Manual Breath</b> <i>P</i> Pressure button Goal Pressure < 30 cm/H <sub>2</sub> O
<b>Decrease Tidal Volume</b> 1 mL/kg increments Until ≤ 29 cm/H <sub>2</sub> O <b>(DO NOT DECREASE &lt; 4 mL/kg)</b>

<b>After 10 minutes</b>
Decrease FiO <sub>2</sub> down to 50% Then adjust PEEP and FiO <sub>2</sub> Goal SpO <sub>2</sub> 92 – 98%
Step 1: PEEP =10 FiO <sub>2</sub> =50%
Step 2: PEEP =10 FiO <sub>2</sub> =60%
Step 3: PEEP =10 FiO <sub>2</sub> =70%
Step 4: PEEP =12 FiO <sub>2</sub> =70%
Step 5: PEEP =14 FiO <sub>2</sub> =70%

<b>Check Peak Inspiratory Pressure (PIP)</b> Goal V <sub>t</sub> is 8 mL/kg
<b>ADJUST PIP Alarm Settings</b>

Notify Destination or Contact Medical Control

**AR 11**



# Mechanical Ventilation; Adult (Optional)

## Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro
- **Mechanical ventilation may be used in any patient  $\geq 1$  year old.**
- **MODE:**
  - In all adult patients use Volume – Assist Control.
  - This mode requires adequate sedation as it can be uncomfortable in a patient who is awakening.
- **TIDAL VOLUME:**
  - Tidal volume is very important in preventing lung injury and calculated by height and predicted body weight, or ideal body weight, and NOT actual body weight.
  - Follow Tidal Volume by Height Table on page 3.
  - Follow Tidal Volume by Height Table on page 3 when adjusting Peak Inspiratory Pressure alarms to allow full exhalation.
  - High Tidal Volumes are well known to cause alveolar damage and lung injury.
- **FLOW RATE:**
  - A normal breath (non-mechanical ventilation) has highest flow and volume at the beginning and both decrease as inspiration comes to an end.
  - Setting Flow Rate at 60 L/minute allows patient to take full breath without air hunger toward end of inspiration. This is more comfortable for the patient.
  - If patient looks like they are trying to take in more volume initially, the Flow Rate can be increased by increments of 5 as needed to improve patient comfort.
- **$\text{FiO}_2$  and PEEP Adjustments:**
  - Seems intuitive that when  $\text{SpO}_2$  is less than desired the  $\text{FiO}_2$  should be increased.
  - When  $\text{FiO}_2$  is  $\geq 50\%$  and  $\text{SpO}_2$  remains low, this indicates a shunt, and PEEP must be used in conjunction with  $\text{FiO}_2$  to correct the shunt and increase oxygenation.
  - Follow PEEP adjustment recommendations on page 1.
- **$\text{EtCO}_2$ :**
  - $\text{EtCO}_2$  and arterial  $\text{CO}_2$  do not always correlate well in patients with lung disease or during serious illness or injury.
  - Use caution in adjusting respiratory rate to reach a goal of 35 – 45 mmHg. Most intubated patients do not need tight control in this range.
  - Patients with suspected head injury do need  $\text{EtCO}_2$  with a target of 35 – 45 mmHg.
  - Allowing patients with COPD and asthma exacerbations to have higher  $\text{EtCO}_2$  outside the 35 – 45 mmHg range is acceptable. Lower ventilation rates allow more time for exhalation and prevents auto-PEEP and/ or air trapping.
- **DOPE:** Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.



# Mechanical Ventilation; Adult (Optional)

## TIDAL VOLUME INITIAL SETTINGS By HEIGHT

### FEMALE

Height / Predicted body weight / Vt

HEIGHT	PBW	4 ml	5 ml	6 ml	7 ml	8 ml
4' 0" (48)	17.9	72	90	107	125	143
4' 1" (49)	20.2	81	101	121	141	162
4' 2" (50)	22.5	90	113	135	158	180
4' 3" (51)	24.8	99	124	149	174	198
4' 4" (52)	27.1	108	136	163	190	217
4' 5" (53)	29.4	118	147	176	206	235
4' 6" (54)	31.7	127	159	190	222	254
4' 7" (55)	34	136	170	204	238	272
4' 8" (56)	36.3	145	182	218	254	290
4' 9" (57)	38.6	154	193	232	270	309
4' 10" (58)	40.9	164	205	245	286	327
4' 11" (59)	43.2	173	216	259	302	346
5' 0" (60)	45.5	182	228	273	319	364
5' 1" (61)	47.8	191	239	287	335	382
5' 2" (62)	50.1	200	251	301	351	401
5' 3" (63)	52.4	210	262	314	367	419
5' 4" (64)	54.7	219	274	328	383	438
5' 5" (65)	57	228	285	342	399	456
5' 6" (66)	59.3	237	297	356	415	474
5' 7" (67)	61.6	246	308	370	431	493
5' 8" (68)	63.9	256	320	383	447	511
5' 9" (69)	66.2	265	331	397	463	530
5' 10" (70)	68.5	274	343	411	480	548
5' 11" (71)	70.8	283	354	425	496	566
6' 0" (72)	73.1	292	366	439	512	585
6' 1" (73)	75.4	302	377	452	528	603
6' 2" (74)	77.7	311	389	466	544	622
6' 3" (75)	80	320	400	480	560	640
6' 4" (76)	82.3	329	412	494	576	658
6' 5" (77)	84.6	338	423	508	592	677
6' 6" (78)	86.9	348	435	521	608	695
6' 7" (79)	89.2	357	446	535	624	714
6' 8" (80)	91.5	366	458	549	641	732
6' 9" (81)	93.8	375	469	563	657	750
6' 10" (82)	96.1	384	481	577	673	769
6' 11" (83)	98.4	394	492	590	689	787
7' 0" (84)	100.7	403	504	604	705	806

### MALE

Height / Predicted body weight / Vt

HEIGHT	PBW	4 ml	5 ml	6 ml	7 ml	8 ml
4' 0" (48)	22.4	90	112	134	157	179
4' 1" (49)	24.7	99	124	148	173	198
4' 2" (50)	27	108	135	162	189	216
4' 3" (51)	29.3	117	147	176	205	234
4' 4" (52)	31.6	126	158	190	221	253
4' 5" (53)	33.9	136	170	203	237	271
4' 6" (54)	36.2	145	181	217	253	290
4' 7" (55)	38.5	154	193	231	270	308
4' 8" (56)	40.8	163	204	245	286	326
4' 9" (57)	43.1	172	216	259	302	345
4' 10" (58)	45.4	182	227	272	318	363
4' 11" (59)	47.7	191	239	286	334	382
5' 0" (60)	50	200	250	300	350	400
5' 1" (61)	52.3	209	262	314	366	418
5' 2" (62)	54.6	218	273	328	382	437
5' 3" (63)	56.9	228	285	341	398	455
5' 4" (64)	59.2	237	296	355	414	474
5' 5" (65)	61.5	246	308	369	431	492
5' 6" (66)	63.8	255	319	383	447	510
5' 7" (67)	66.1	264	331	397	463	529
5' 8" (68)	68.4	274	342	410	479	547
5' 9" (69)	70.7	283	354	424	495	566
5' 10" (70)	73	292	365	438	511	584
5' 11" (71)	75.3	301	377	452	527	602
6' 0" (72)	77.6	310	388	466	543	621
6' 1" (73)	79.9	320	400	479	559	639
6' 2" (74)	82.2	329	411	493	575	658
6' 3" (75)	84.5	338	423	507	592	676
6' 4" (76)	86.8	347	434	521	608	694
6' 5" (77)	89.1	356	446	535	624	713
6' 6" (78)	91.4	366	457	548	640	731
6' 7" (79)	93.7	375	469	562	656	750
6' 8" (80)	96	384	480	576	672	768
6' 9" (81)	98.3	393	492	590	688	786
6' 10" (82)	100.6	402	503	604	704	805
6' 11" (83)	102.9	412	515	617	720	823
7' 0" (84)	105.2	421	526	631	736	842

### TROUBLESHOOTING Hypoxia or Deterioration DOPES

### RESPONSE to Hypoxia or Deterioration DOTT

D	Dislodged ETT or cuff leak	D	Disconnect ventilator, squeeze chest if auto-PEEP, Decompress if pneumothorax
O	Obstruction of ETT or circuit	O	Oxygen 100% FiO <sub>2</sub> , BVM and check compliance
P	Pneumothorax, Pneumonia, Pulmonary embolism or edema, Plug (mucous)	T	Tube position and function, check EtCO <sub>2</sub>
E	Equipment problem	T	Tweak ventilator settings or equipment
S	Stacked breaths, air trapping, or auto-PEEP		

Pressure Alarm Troubleshooting			Problem Location	Consider
High PIP	+	High Plateau > 30	Alveoli	Compliance problem: Pneumothorax, Pneumonia, Pulmonary Edema or Embolism, CHF
High PIP	+	Normal Plateau < 30	Airway problem	Airway, ventilator, or circuit problem: DOPE, Right Main stem intubation, Air trapping or auto-PEEP, Mucous plug, Patient out of synchrony with ventilator



# Pediatric Asystole / PEA

## History

- Events leading to arrest
- Estimated downtime
- SAMPLE
- Existence of terminal illness
- Airway obstruction
- Hypothermia
- Suspected abuse

## Signs and Symptoms

- Pulseless
- Apneic
- No electrical activity on ECG
- No heart tones on auscultation

## Differential

- Respiratory failure
- Foreign body
- Infection (croup, epiglottitis)
- Congenital heart disease
- See Reversible Causes below

### Pediatric Pulseless Arrest Protocol

#### Criteria for Death / No Resuscitation Review DNR / MOST Form

YES

Decomposition  
Rigor mortis  
Dependent lividity  
Blunt force trauma  
Injury incompatible with life  
Extended downtime with asystole

Do not begin resuscitation

Follow  
Deceased Subjects  
Policy

NO

**Begin Continuous CPR Compressions**  
Push Hard ( $\geq 1/3$  AP Diameter of Chest)  
(1.5 inches Infant / 2 inches in Children)  
Push Fast (100 - 120 / min)  
Change Compressors every 2 minutes  
(sooner if fatigued)  
(Limit changes / pulse checks  $\leq 10$  seconds)

#### Ventilation rate:

1 breath every 2 seconds when age  $< 1$   
1 breath every 3 seconds when age  $\geq 1$   
15:2 Compression:Ventilation if no Advanced Airway

AED Procedure  
*if available*

Cardiac Monitor

IV or IO Access Protocol UP 6

**P** Epinephrine 1:10,000  
0.01 mg/kg IV / IO Maximum Single Dose 1mg  
Or  
Epinephrine 1:1000 0.1 mg / kg ETT Maximum 2.5 mg  
Repeat every 3 – 5 minutes

**A** Normal Saline Bolus 20 mL/kg IV / IO  
May repeat as needed  
Maximum 60 mL/kg

Search for Reversible Causes

Blood Glucose Analysis Procedure  
*if applicable*

### Reversible Causes

Hypovolemia  
Hypoxia  
Hydrogen ion (acidosis)  
Hypothermia  
Hypo / Hyperkalemia

Tension pneumothorax  
Tamponade; cardiac  
Toxins  
Thrombosis; pulmonary (PE)  
Thrombosis; coronary (MI)

## AT ANY TIME

Return of  
Spontaneous  
Circulation



Go to  
Post Resuscitation  
Protocol

**P**

**A**



**PC 1**



# Pediatric Asystole / PEA

## Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks.**
- **Refer to optional protocol AC 11 or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Compress  $\geq 1/3$  anterior-posterior diameter of chest, in infants 1.5 inches and in children 2 inches.**
- **Majority of pediatric arrests stem from a respiratory insult or hypoxic event. Compressions should be coupled with ventilations.**
- **When advanced airway not in place perform 15 compressions with 2 ventilations.**
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children  $< 10$  kg.**
- **DO NOT HYPERVENTILATE:**  
If advanced airway in place ventilate:
  - Age  $< 1$  year: 1 breath every 2 seconds with continuous, uninterrupted compressions.
  - Age  $\geq 1$  year: 1 breath every 3 seconds with continuous, uninterrupted compressions.
- **Airway is a more important intervention in pediatric arrests. This should be accomplished quickly with BVM or BIAD.**
- **Patient survival is often dependent on proper ventilation and oxygenation / airway Interventions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **High-Quality CPR:**  
Make sure chest compressions are being delivered at 100 – 120 / min.  
Make sure chest compressions are adequate depth for age and body habitus.  
Make sure you allow full chest recoil with each compression to provide maximum perfusion.  
Minimize all interruptions in chest compressions to  $< 10$  seconds.  
Use AED or apply ECG monitor / defibrillator as soon as available.
- **Defibrillation:** Follow manufacturer's recommendations concerning defibrillation / cardioversion energy when specified.
- **End Tidal CO<sub>2</sub> (EtCO<sub>2</sub>)**  
If EtCO<sub>2</sub> is  $< 10$  mmHg, improve chest compressions. Goal is  $\geq 20$  mmHg.  
If EtCO<sub>2</sub> spikes, typically  $> 40$  mmHg, consider Return of Spontaneous Circulation (ROSC)
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Special Considerations**  
**Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm.  
Defibrillation is safe at all energy levels.
- **Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
- **Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol UP 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
- **Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.**



# Pediatric Bradycardia With a Pulse

## History


- Past medical history
- Foreign body exposure
- Respiratory distress or arrest
- Apnea
- Possible toxic or poison exposure
- Congenital disease
- Medication (maternal or infant)

## Signs and Symptoms

- Decreased heart rate
- Delayed capillary refill or cyanosis
- Mottled, cool skin
- Hypotension or arrest
- Altered level of consciousness

## Differential

- Respiratory failure, Foreign body, Secretions, Infection (croup, epiglottitis)
- Hypovolemia (dehydration)
- Congenital heart disease
- Trauma
- Tension pneumothorax
- Hypothermia
- Toxin or medication
- Hypoglycemia
- Acidosis





# Pediatric Bradycardia With Poor Perfusion

## Pearls

- Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro
- Bradycardia is often associated with hypoxia so insure patent airway, breathing, and circulation as needed.
- Begin CPR immediately with persistent bradycardia and poor perfusion despite adequate oxygenation and ventilation.
- Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.
- Rhythm should be interpreted in the context of symptoms and pharmacological treatment given only when symptomatic, otherwise monitor and reassess.
- Consider hyperkalemia with wide complex, bizarre appearance of QRS complex, and bradycardia.
- 12-Lead ECG:
  - 12 Lead ECG not necessary to diagnose and treat
  - Obtain when patient is stable and/or following rhythm conversion.
- Epinephrine is first drug choice for persistent, symptomatic bradycardia.
- Atropine:
  - Second choice, unless there is evidence of increased vagal tone or a primary AV conduction block, then give atropine first.
  - Ineffective and potentially harmful in cardiac transplantation. May cause paradoxical bradycardia.
- Symptomatic bradycardia causing shock or peri-arrest condition:
  - If no IV or IO access immediately available, start Transcutaneous Pacing, establish IV / IO access, and then administer epinephrine.
  - Epinephrine should be administered followed Atropine if no response.
- Symptomatic condition
  - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
  - Symptomatic bradycardia usually occurs at rates < 50 beats per minute.
  - Search for underlying causes such as hypoxia or impending respiratory failure.
- Serious Signs / Symptoms:
  - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.
- Transcutaneous Pacing Procedure (TCP)
  - Indicated with unstable bradycardia unresponsive to medical therapy.
  - If time allows transport to specialty center because transcutaneous pacing is a temporizing measure.
  - Transvenous / permanent pacemaker will probably be needed.
  - Immediate TCP with high-degree AV block (2d or 3d degree) with no IV / IO access.
- Most maternal medications pass through breast milk to the infant so maintain high-index of suspicion for OD-toxins.
- Hypoglycemia, severe dehydration and narcotic effects may produce bradycardia. Many other agents a child ingests can cause bradycardia, often is a single dose.



# Pediatric Pulmonary Edema / CHF

## History


- Congenital Heart Disease
- Chronic Lung Disease
- Congestive heart failure
- Past medical history

## Signs/Symptoms

- Infant: Respiratory distress, poor feeding, lethargy, weight gain, +/- cyanosis
- Child/Adolescent: Respiratory distress, bilateral rales, apprehension, orthopnea, jugular vein distention (rare), pink, frothy sputum, peripheral edema, diaphoresis, chest pain
- Hypotension, shock

## Differential

- Congestive heart failure
- Asthma
- Anaphylaxis
- Aspiration
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pericardial tamponade
- Toxic Exposure



## Pearls

- **Recommended exam: Mental status, Respiratory, Cardiac, Skin, Neuro**
- **Contact Medical Control early in the care of the pediatric cardiac patient.**
- **Most children with CHF have a congenital heart defect, obtain a precise past medical history.**
- **Congenital heart disease varies by age:**
  - < 1 month: Tetralogy of Fallot, Transposition of the great arteries, Coarctation of the aorta.
  - 2 – 6 months: Ventricular septal defects (VSD), Atrioseptal defects (ASD).
  - Any age: Myocarditis, Pericarditis, SVT, heart blocks.
- **Treatment of Congestive Heart Failure / Pulmonary edema may vary depending on the underlying cause and may include the following with consultation by Medical Control:**
  - Morphine Sulfate: 0.1 mg/kg IV / IO. Max single dose 5mg/dose
  - Fentanyl: 1 mcg/kg IV / IO. Max single dose 50 mcg.
  - Nitroglycerin: Dose determined after consultation of Medical Control.
  - Lasix 1 mg/kg IV / IO.
  - Agency specific vasopressor.
- Do not assume all wheezing is pulmonary, especially in a cardiac child: avoid albuterol unless strong history of recurrent wheezing secondary to pulmonary etiology (discuss with Medical Control)



# Pediatric Cardiac Arrest

## History

- Time of arrest
- Medical history
- Medications
- Possibility of foreign body
- Hypothermia

## Signs and Symptoms

- Unresponsive
- Cardiac arrest

## Differential

- Respiratory failure: Foreign body, Secretions, Infection (croup, epiglottitis)
- Hypovolemia (dehydration)
- Congenital heart disease
- Trauma
- Tension pneumothorax, cardiac tamponade, pulmonary embolism
- Hypothermia
- Toxin or medication
- Electrolyte abnormalities (Glucose, K)
- Acidosis

## Protocol Age Guidance:

Newborn – 3 days:  
AO2 Newly Born

3- days to 15 years:  
PC4 Pediatric Cardiac Arrest

≥ 16 years:  
AC3 Cardiac Arrest;  
Adult

## Criteria for Death / No Resuscitation Review DNR / MOST Form

NO

Do not begin resuscitation  
Follow Deceased Subjects Policy

**Begin Continuous CPR Compressions**  
Push Hard ( $\geq 1/3$  AP Diameter of Chest)  
(1.5 inches Infant / 2 inches in Children)  
Push Fast (100 - 120 / min)  
Change Compressors every 2 minutes  
(sooner if fatigued)  
(Limit changes / pulse checks  $\leq 10$  seconds)

### Ventilation rate:

1 breath every 2 seconds when age  $< 1$

1 breath every 3 seconds when age  $\geq 1$

15:2 Compression:Ventilation if no Advanced Airway

AED Procedure  
*if available*

ALS Available

P Cardiac Monitor

Shockable Rhythm

Defibrillation Automated

Continue CPR  
2 Minutes

Repeat and reassess

Pediatric Airway  
Protocol(s) 5, 6

Pediatric Asystole /  
PEA  
Protocol PC 1

Pediatric Airway  
Protocol(s) 5, 6

Pediatric VF / VT  
Protocol PC 6  
Pediatric  
Tachycardia  
Protocol PC 5

Pediatric Airway  
Protocol(s) 5, 6

Naloxone 0.4 – 2 mg IN / IM  
Peds: 0.1 mg/kg IN  
Maximum 4 mg

Naloxone 0.4 – 2 mg  
Peds: 0.1 mg/kg  
IV / IO / IM / IN / ETT  
Maximum 4 mg

NO

Arrest secondary  
to Opioid OD?

Notify Destination or  
Contact Medical Control



# Pediatric Cardiac Arrest

## Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Compress  $\geq 1/3$  anterior-posterior diameter of chest, in infants 1.5 inches and in children 2 inches.**
- **Majority of pediatric arrests stem from a respiratory insult or hypoxic event. Compressions should be coupled with ventilations.**
- **When advanced airway not in place perform 15 compressions with 2 ventilations.**
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children  $< 10$  kg.**
- **DO NOT HYPERVENTILATE:**  
If advanced airway in place ventilate:
  - Age  $< 1$  year: 1 breath every 2 seconds with continuous, uninterrupted compressions.
  - Age  $\geq 1$  year: 1 breath every 3 seconds with continuous, uninterrupted compressions.
- **Patient survival is often dependent on proper ventilation and oxygenation / airway Interventions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **High-Quality CPR:**  
Make sure chest compressions are being delivered at 100 – 120 / min.  
Make sure chest compressions are adequate depth for age and body habitus.  
Make sure you allow full chest recoil with each compression to provide maximum perfusion.  
Minimize all interruptions in chest compressions to  $< 10$  seconds.  
Use AED or apply ECG monitor / defibrillator as soon as available.
- **Defibrillation:**  
Follow manufacturer's recommendations concerning defibrillation / cardioversion energy when specified.  
Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.  
Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO<sub>2</sub> (EtCO<sub>2</sub>)**  
If EtCO<sub>2</sub> is  $< 10$  mmHg, improve chest compressions. Goal is  $\geq 20$  mmHg.  
If EtCO<sub>2</sub> spikes, typically  $> 40$  mmHg, consider Return of Spontaneous Circulation (ROSC)
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Special Considerations**  
**Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm.  
Defibrillation is safe at all energy levels.
- **Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
- **Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol UP 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
- **Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** - Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.



# Pediatric Tachycardia

## Narrow Complex ( $\leq 0.09$ sec)

### History


- Past medical history
- Medications or Toxic Ingestion (Aminophylline, Diet pills, Thyroid supplements, Decongestants, Digoxin)
- Drugs (nicotine, cocaine)
- Congenital Heart Disease
- Respiratory Distress
- Syncope or Near Syncope

### Signs and Symptoms

- Heart Rate: Child > 180/bpm  
Infant > 220/bpm
- Pale or Cyanosis
- Diaphoresis
- Tachypnea
- Vomiting
- Hypotension
- Altered Level of Consciousness
- Pulmonary Congestion
- Syncope

### Differential

- Heart disease (Congenital)
- Hypo / Hyperthermia
- Hypovolemia or Anemia
- Electrolyte imbalance
- Anxiety / Pain / Emotional stress
- Fever / Infection / Sepsis
- Hypoxia, Hypoglycemia
- Medication / Toxin / Drugs (see HX)
- Pulmonary embolus
- Trauma, Tension Pneumothorax





# Pediatric Tachycardia

## Narrow Complex ( $\leq 0.09$ sec)

### Pearls

- **Recommended Exam:** Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro
- **Monomorphic QRS:**
  - All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
  - QRS complexes in a single lead will change from complex to complex.
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance.** Pediatric paddles should be used in children  $< 10$  kg.
- **Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.**
- **12-Lead ECG:**
  - 12-Lead ECG not necessary to diagnose and treat.
  - Obtain when patient is stable and/or following rhythm conversion.
  - When administering adenosine, obtaining a continuous 12-Lead can be helpful to physicians.
- **Unstable condition:**
  - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
  - If at any point patient becomes unstable move to unstable arm in algorithm
  - If IV or IO access is in place, may administer adenosine and repeat, prior to synchronized cardioversion.
- **Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.**
- **Serious Signs and Symptoms:**
  - Respiratory distress / failure.
  - Signs of shock / poor perfusion with or without hypotension.
  - AMS
  - Sudden collapse with rapid, weak pulse
- **Narrow Complex Tachycardia ( $\leq 0.09$  seconds):**
  - Sinus tachycardia: P waves present. Variable R-R waves. Infants usually  $< 220$  beats / minute. Children usually  $< 180$  beats / minute.
  - SVT:  $> 90\%$  of children with SVT will have a narrow QRS ( $\leq 0.09$  seconds.) P waves absent or abnormal. R-R waves not variable. Usually abrupt onset. Infants usually  $> 220$  beats / minute. Children usually  $> 180$  beats / minute.
  - Atrial Flutter / Fibrillation
- **Vagal Maneuvers:**
  - Breath holding. Blowing a glove into a balloon. Have child blow out “birthday candles” or through an obstructed straw. Infants: May put a bag of ice water over the upper half of the face careful not to occlude the airway.
- Separating the child from the caregiver may worsen the child's clinical condition.
- Monitor for respiratory depression and hypotension associated if Diazepam, Lorazepam, or Midazolam is used.
- Continuous pulse oximetry is required for all SVT Patients if available.



# Pediatric Tachycardia

## Wide Complex (> 0.09 sec)

### History

- Past medical history
- Medications or Toxic Ingestion (Aminophylline, Diet pills, Thyroid supplements, Decongestants, Digoxin)
- Drugs (nicotine, cocaine)
- Congenital Heart Disease
- Respiratory Distress
- Syncope or Near Syncope

### Signs and Symptoms

- Heart Rate: Child > 180/bpm  
Infant > 220/bpm
- Pale or Cyanosis
- Diaphoresis
- Tachypnea
- Vomiting
- Hypotension
- Altered Level of Consciousness
- Pulmonary Congestion
- Syncope

### Differential

- Heart disease (Congenital)
- Hypothermia/ Hyperthermia
- Hypovolemia or Anemia
- Electrolyte imbalance
- Anxiety/ Pain/ Emotional stress
- Fever/ Infection/ Sepsis
- Hypoxia, Hypoglycemia
- Medication/ Toxin/ Drugs (see HX)
- Pulmonary embolus
- Trauma, Tension Pneumothorax

Assess tachycardia in context of clinical condition  
Identify and treat underlying cause of tachycardia

Unstable / Serious Signs and Symptoms  
AMS, shock, hypotension  
HR Typically > 180 Child  
HR Typically > 220 Infant

YES → P

Cardiac Monitor
Cardioversion Procedure
<b>0.5 – 1 J / kg</b> Repeat and increase to <b>2 J / Kg</b> May increase to <b>4 J / Kg</b> Or adult maximum
Consider sedation <u>Do NOT delay cardioversion</u> <b>Midazolam 0.1 – 0.2 mg / kg IV / IO / IN</b> May repeat if needed <b>Maximum Single Dose 2 mg</b> <b>Maximum Total Dose 5 mg</b>

NO

<b>B</b>	12 Lead ECG Procedure
<b>P</b>	Cardiac Monitor
	IV or IO Access Protocol UP 6

Single lead ECG able to diagnose and treat arrhythmia

12 Lead ECG not necessary to diagnose and treat, but preferred when patient is stable.

Regular Rhythm?

YES

Probable Ventricular Tachycardia

Contact Medical Control

<b>P</b>	<b>If QRS Regular and Monomorphic</b> Adenosine 0.1 mg / kg IV / IO Maximum 6 mg  May repeat Adenosine 0.2 mg / kg IV / IO Maximum 12 mg
	Expert consultation recommended for further medication management and/ or cardioversion procedure

**AT ANY TIME**

Pulseless



Go to  
Pediatric Pulseless Arrest Protocol

	Notify Destination or Contact Medical Control	
--	---	--



# Pediatric Tachycardia

## Wide Complex (> 0.09 sec)

### Pearls

- **Recommended Exam:** Mental Status, Skin, Neck, Lung, Heart, Abdomen, Neuro
- **Monomorphic QRS:**
  - All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
  - QRS complexes in a single lead will change from complex to complex.
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance.** Pediatric paddles should be used in children < 10 kg.
- **Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.**
- **12-Lead ECG:**
  - 12-Lead ECG is not necessary to diagnose and treat arrhythmia. A single lead ECG is often all that is needed.
  - Obtain 12-Lead when patient is stable and/ or following a rhythm conversion.
  - When administering adenosine, obtaining a continuous 12-Lead can be helpful later to physicians.
- **Unstable condition:**
  - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
  - If at any point patient becomes unstable move to unstable arm in algorithm
- **Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.**
- **Serious Signs and Symptoms:**
  - Respiratory distress/ failure.
  - Signs of shock/ poor perfusion with or without hypotension.
  - AMS
  - Sudden collapse with rapid, weak pulse
- **Serious Signs and Symptoms:**
  - Respiratory distress/ failure.
  - Signs of shock/ poor perfusion with or without hypotension.
  - AMS
  - Sudden collapse with rapid, weak pulse
- **Wide Complex Tachycardia (≥ 0.09 seconds):**
  - SVT with aberrancy.
  - VT: Uncommon in children. Rates may vary from near normal to > 200/ minute.
  - Most children with VT have underlying heart disease / cardiac surgery/ long QT syndrome/ cardiomyopathy.
  - Amiodarone 5 mg / kg over 20 – 60 minutes or Procainamide 15 mg / kg over 30 – 60 minutes IV / IO** are recommended agents. They should not be administered together. Consultation with Medical Control is advised when these agents are considered.
- **Torsade's de Pointes/ Polymorphic (multiple shaped) Tachycardia:**
  - Rate is typically 150 to 250 beats/ minute.
  - Associated with long QT syndrome, hypomagnesaemia, hypokalemia, many cardiac drugs.
  - May quickly deteriorate to VT.
  - Separating the child from the caregiver may worsen the child's clinical condition.
- Monitor for respiratory depression and hypotension associated if Diazepam, Lorazepam, or Midazolam is used.
- Continuous pulse oximetry is required for all SVT patients if available.



# Pediatric Ventricular Fibrillation Pulseless Ventricular Tachycardia

## History

- Events leading to arrest
- Estimated downtime
- Past medical history
- Medications
- Existence of terminal illness
- Airway obstruction
- Hypothermia

## Signs and Symptoms

- Unresponsive
- Cardiac Arrest

## Differential

- Respiratory failure / Airway obstruction
- Hyper / hypokalemia, Hypovolemia
- Hypothermia, Hypoglycemia, Acidosis
- Tension pneumothorax, Tamponade
- Toxin or medication
- Thrombosis: Coronary / Pulmonary Embolism
- Congenital heart disease

### Pediatric Pulseless Arrest Protocol PC 4

**Begin Continuous CPR Compressions**  
**Push Hard ( $\geq 1/3$  AP Diameter of Chest)**  
**(1.5 inches Infant / 2 inches in Children)**  
**Push Fast (100 - 120 / min)**  
**Change Compressors every 2 minutes**  
**(sooner if fatigued)**  
**(Limit changes / pulse checks  $\leq 10$  seconds)**

#### Ventilation rate:

1 breath every 2 seconds when age  $< 1$   
 1 breathe every 3 seconds when age  $\geq 1$

**15:2 Compression:Ventilation if no Advanced Airway**

#### Automated Defibrillation Procedure

#### Defibrillation Manual Procedure

A

- First shock: 2 J / Kg
- Second shock: 4 J / Kg
- Subsequent shocks  $\geq 4$  J / kg  
Maximum 10 J / kg or adult dose

#### IV / IO Protocol UP 6

#### Epinephrine 1:10,000

0.01 mg/kg IV / IO Maximum 1mg

Or

#### Epinephrine 1:1000 0.1 mg / kg ETT

Maximum 2.5 mg

Repeat every 3 – 5 minutes

#### If Rhythm Refractory to defibrillation

- Continue CPR and give Agency specific Anti-arrhythmic(s) in a drug-shock-drug-shock pattern.
- Continue CPR up to point where you are ready to defibrillate with device charged.

Repeat pattern during resuscitation.

P

Amiodarone 5mg/kg (max 300mg)

A

#### Magnesium Sulfate 40 mg/kg IV / IO

Infuse over 2 – 3 minutes

Maximum 2 gm

Notify Destination or  
Contact Medical Control

## AT ANY TIME

### Return of Spontaneous Circulation



Go to Post Resuscitation Protocol

## Reversible Causes

Hypovolemia  
 Hypoxia  
 Hydrogen ion (acidosis)  
 Hypothermia  
 Hypo / Hyperkalemia  
 Hypoglycemia

Tension pneumothorax  
 Tamponade; cardiac  
 Toxins  
 Thrombosis; pulmonary (PE)  
 Thrombosis; coronary (MI)



# Pediatric Ventricular Fibrillation Pulseless Ventricular Tachycardia (Optional)

## Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Compress  $\geq 1/3$  anterior-posterior diameter of chest, in infants 1.5 inches and in children 2 inches.**
- **Majority of pediatric arrests stem from a respiratory insult or hypoxic event. Compressions should be coupled with ventilations.**
- **When advanced airway not in place perform 15 compressions with 2 ventilations.**
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children  $< 10$  kg.**
- **DO NOT HYPERVENTILATE:**  
**If advanced airway in place ventilate:**  
**Age  $< 1$  year: 1 breath every 2 seconds with continuous, uninterrupted compressions.**  
**Age  $\geq 1$  year: 1 breath every 3 seconds with continuous, uninterrupted compressions.**
- **Patient survival is often dependent on proper ventilation and oxygenation / airway Interventions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **High-Quality CPR:**  
Make sure chest compressions are being delivered at 100 – 120 / min.  
Make sure chest compressions are adequate depth for age and body habitus.  
Make sure you allow full chest recoil with each compression to provide maximum perfusion.  
Minimize all interruptions in chest compressions to  $< 10$  seconds.  
Use AED or apply ECG monitor / defibrillator as soon as available.
- **Defibrillation:**  
Follow manufacturer's recommendations concerning defibrillation / cardioversion energy when specified.  
Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.  
Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.  
**Manual Defibrillation at the AEMT level is permissible only during pulseless cardiac arrest with VF or VT.**
- **End Tidal CO<sub>2</sub> (EtCO<sub>2</sub>)**  
If EtCO<sub>2</sub> is  $< 10$  mmHg, improve chest compressions. Goal is  $\geq 20$  mmHg.  
If EtCO<sub>2</sub> spikes, typically  $> 40$  mmHg, consider Return of Spontaneous Circulation (ROSC)
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Special Considerations**  
**Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm.  
Defibrillation is safe at all energy levels.
- **Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
- **Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol UP 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
- **Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.



# Pediatric Post Resuscitation

## History

- Respiratory arrest
- Cardiac arrest

## Signs/Symptoms

- Return of pulse

## Differential

- Continue to address specific differentials associated with the original dysrhythmia

### Transport Destination Decision

Post-resuscitation patient is medically complex.

### Consider facility capabilities:

- Pediatric ICU service
- Pediatric Cardiology service
- Pediatric Neurology service
- Targeted Temperature Management

Pediatric Airway Protocol(s) AR 5 - 7 <i>as needed</i>	
	Monitor Vital Signs / Reassess
	Blood Glucose Analysis Procedure
	<b>Optimize Ventilation and Oxygenation</b>
	<ul style="list-style-type: none"> <li>• Maintain SpO<sub>2</sub> ≥ 92 – 98%</li> <li>• Advanced airway if indicated</li> <li>• Age Appropriate Respiratory Rate</li> <li>• Remove Impedence Threshold Device</li> </ul> <p><b>DO NOT HYPERVENTILATE</b></p>
	ETCO <sub>2</sub> ideally 35 – 45 mm Hg
<b>B</b>	12 Lead ECG Procedure
	IV or IO Protocol UP 6
<b>P</b>	Cardiac Monitor
	Pediatric Diabetic Protocol PM 2 <i>if indicated</i>
	Pediatric Hypotension / Shock Protocol PM 3 <i>if indicated</i>
	Pediatric Bradycardia Protocol PC 2 <i>if indicated</i>
	Pediatric Tachycardia Protocol PC 5, 6 <i>as indicated</i>

### Hypotension Age Based

**0 – 31 Days**  
< 60 mmHg

**1 Month to 1 Year**  
< 70 mmHg

**> than 1 Year**  
< 70 + ( 2 x age) mmHg

Arrhythmias are common and usually self limiting after ROSC



If Arrhythmia Persists follow Rhythm Appropriate Protocol

Antiarrhythmic Medication Given During Arrest

NO

YES

Continue Antiarrhythmic Utilized  
Refer to Appropriate Pediatric Arrhythmia Protocol

Agency Specific Antiarrhythmic

Post-intubation / BIAD Management Protocol AR 8

Notify Destination or Contact Medical Control



# Pediatric Post Resuscitation

## Pearls

- **Recommended Exam:** Mental Status, Neck, Skin, Lungs, Heart, Abdomen, Extremities, Neuro
- **Goals of care are to preserve neurologic function, prevent secondary organ damage, treat the underlying cause of illness, and optimize prehospital care.** Frequent reassessment is necessary.
- **Hyperventilation is a significant cause of hypotension and recurrence of cardiac arrest in the post resuscitation phase and must be avoided.** Titrate  $\text{FIO}_2$  to maintain  $\text{SpO}_2$  of 92 - 98%.
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance.** Pediatric paddles should be used in children < 10 kg.
- **Pain/sedation:**
  - Patients requiring advanced airways and ventilation commonly experience pain and anxiety.
  - Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.
  - Ventilated patients cannot communicate pain / anxiety and providers are poor at recognizing pain / anxiety.
  - Vital signs such as tachycardia and / or hypertension can provide clues to inadequate sedation, however they both are not always reliable indicators of patient's lack of adequate sedation.
  - Pain must be addressed first, before anxiety. Opioids are typically the first line agents before benzodiazepines. Ketamine is also a reasonable first choice agent.
- **Ventilator / Ventilation strategies:**
  - Tailored to individual patient presentations. Medical Control can indicate different strategies above.
  - In general ventilation with BVM should cause chest rise. With mechanical ventilation a reasonable tidal volume should be about 6 mL/kg and peak pressures should be < 30 cmH2O.
  - Continuous pulse oximetry and capnography should be maintained during transport for monitoring.
  - Head of bed should be maintained at least 10 – 20 degrees of elevation when possible to decrease aspiration risk.
- **EtCO<sub>2</sub> Monitoring:**
  - Initial End tidal CO<sub>2</sub> may be elevated immediately post-resuscitation, but will usually normalize.
  - Goal is 35 – 45 mmHg but DO NOT hyperventilate to achieve.
  - EtCO<sub>2</sub> should be continually monitored with advanced airway in place.
- Administer resuscitation fluids and vasopressor agents to maintain SBP at targets listed on page 1. This table represents minimal SBP targets.
- Targeted Temperature Management is recommended in pediatrics, but prehospital use is not associated with improved outcomes. Transport to facility capable of intensive pediatric care.
- Consider transport to facility capable of managing the post-arrest patient including hypothermia therapy, cardiology / cardiac catheterization, intensive care service, and neurology services.
- The condition of post-resuscitation patients fluctuates rapidly and continuously, and they require close monitoring. Appropriate post-resuscitation management may best be planned in consultation with Medical Control.



# Pediatric Allergic Reaction

## History


- Onset and location
- Insect sting or bite
- Food allergy/ exposure
- Medication allergy/ exposure
- New clothing, soap, detergent
- Past medical history/ reactions
- Medication history

## Signs and Symptoms

- Itching or hives
- Coughing/ wheezing or respiratory distress
- Chest or throat constriction
- Difficulty swallowing
- Hypotension or shock
- Edema

## Differential

- Urticaria (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration/ Airway obstruction
- Vasovagal event
- Asthma/ COPD /CHF





# Pediatric Allergic Reaction (Optional)

## Pearls

- **Recommended Exam:** Mental Status, Skin, Heart, Lungs, Abdomen
- **Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.**
- **Epinephrine administration:**
  - Drug of choice and the FIRST drug that should be administered in acute anaphylaxis (Moderate/ Severe Symptoms.)
  - IM Epinephrine should be administered in priority before or during attempts at IV or IO access.
- **Diphenhydramine and steroid administration:**
  - Diphenhydramine/ steroids have no proven benefit in Moderate/ Severe anaphylaxis.
  - Diphenhydramine/ steroids should NOT delay initial or repeat Epinephrine administration.
  - In Moderate and Severe anaphylaxis, Diphenhydramine may decrease mental status.
  - Diphenhydramine should NOT be given to a patient with decreased mental status and/ or a hypotensive patient as this may cause nausea, vomiting, and/ or worsening mental status.
- **Anaphylaxis unresponsive to repeat doses of IM epinephrine may require IV epinephrine administration by IV push or epinephrine infusion. Contact Medical Control for appropriate dosing.**
- **Symptom Severity Classification:**
  - Mild symptoms:**
    - Flushing, hives, itching, erythema with normal blood pressure and perfusion.
  - Moderate symptoms:**
    - Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with normal blood pressure and perfusion.
  - Severe symptoms:**
    - Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with hypotension and poor perfusion.
- **Allergic reactions may occur with only respiratory and gastrointestinal symptoms and have no rash/ skin involvement.**
- **Angioedema** is seen in moderate to severe reactions and is swelling involving the face, lips or airway structures. This can also be seen in patients taking blood pressure medications like Prinivil / Zestril (lisinopril)-typically end in -ii.
- **Hereditary Angioedema** involves swelling of the face, lips, airway structures, extremities, and may cause moderate to severe abdominal pain. Some patients are prescribed specific medications to aid in reversal of swelling. **Paramedic may assist or administer this medication per patient/ package instructions.**
- **Fluids and Medication titrated to maintain a SBP  $>70 + (\text{age in years} \times 2)$  mmHg.**
- **Patients with moderate and severe reactions should receive a 12-Lead ECG and should be continually monitored, but this should NOT delay administration of epinephrine.**
- **EMR/ EMT:**
  - The use of Epinephrine IM is limited to the treatment of anaphylaxis and may be given only by autoinjector, unless manual draw-up is approved by the Agency Medical Director and the NC office of EMS.
  - Administration of diphenhydramine is limited to the oral route only.**
- **EMT administration of beta-agonist is limited to only patients currently prescribed the medication,** unless approved by the Agency Medical Director and the NC office of EMS.
- Agency Medical Director may require contact of medical control prior to EMT/ EMR administering any medication(s). Medical Director may require contact of medical control prior to EMT/ EMR administering any medication.
- The shorter the onset from exposure to symptoms the more severe the reaction.



# Pediatric Diabetic

## History

- Past medical history
- Medications
- Recent blood glucose check
- Last meal

## Signs and Symptoms

- Altered mental status
- Combative/ irritable
- Diaphoresis
- Seizures
- Abdominal pain
- Nausea/ vomiting
- Weakness
- Dehydration
- Deep/ rapid breathing

## Differential

- Alcohol/ drug use
- Toxic ingestion
- Trauma; head injury
- Seizure
- CVA
- Altered baseline mental status.

	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure <i>if indicated</i>
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Hypotension/ Shock Protocol AM 5 <i>if indicated</i>
	Seizure Protocol UP 13 <i>if indicated</i>

**Blood glucose  $\leq 69$  mg/dl**

**Symptomatic with NO IV / IO Access:** Awake, alert and able to tolerate oral agent:

Give **oral glucose solution**.

If unable to tolerate oral: **Glucagon 0.1 mg/kg IM (Maximum 1 mg)**  
Repeat every 15 minutes as needed to keep Blood glucose  $> 60$  mg / dl.

**Blood Sugar  $\leq 69$  mg / dl**

**Blood Sugar 70 – 249 mg / dl**

**Blood Sugar  $\geq 250$  mg / dl**

**Oral Glucose ½ to 1 Tube**  
If age appropriate

Consider Oral Solution (Juices / Food)

**A**   
**< 1 Year of Age:**  
D10: 5 mL/kg IV / IO  
Repeat as needed

**1 to 2 Years:**  
D25: 2 mL/kg IV / IO  
Repeat as needed

**$\geq 2$  Years:**  
D50: 1 mL/kg IV / IO  
**Maximum 25 g per dose**  
Repeat as needed

**Blood Glucose Analysis Procedure**  
***if condition changes***

Exit to  
Appropriate  
Protocol(s)

**Blood Sugar  $\geq 250$  mg / dl**

**A** **Normal Saline Bolus 10 - 20 mL/kg IV / IO**  
**Maximum 20 mL/kg**

**Monitor and Reassess**  
**Every 5 minutes**  
**Until Blood Glucose  $\geq 80$  mg / dl**

**Notify Destination or**  
**Contact Medical Control**



# Pediatric Diabetic

## Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Respirations and effort, Abdomen, Neuro.**
- **Patients with prolonged hypoglycemia or those who are malnourished may not respond to glucagon.**
- **Do not administer oral glucose to patients that are not able to swallow or protect their airway.**
- **Quality control checks should be maintained per manufacturers recommendation for all glucometers.**
- **D10/ D25 Preparation:**
  - D10: Remove 10 mL of D50 from a D50 vial. Add 40 mL of NS with the 10 mL of D50 with a total volume of 50 mL.
  - D10: Alternative, Discard 40 mL from the D50 vial and draw up 40 mL of NS with a total volume of 50 mL.
  - D25: Remove 25 mL of D50 and draw up 25 mL of NS with a total volume of 50 mL.
- **Patient's refusing transport to medical facility after treatment of hypoglycemia:**
  - Adult caregiver must be present with pediatric patient.
  - Blood sugar must be  $\geq 80$ , patient has ability to eat and availability of food with responders on scene.
  - Blood sugar trending up is necessary with 2 or more readings above 80, obtained over 15-minute period.
  - Patient must have known history of diabetes and not taking any oral diabetic agents.
  - Patient returns to normal mental status and has a normal neurological exam with no new neurological deficits.
  - Must demonstrate capacity to make informed health care decisions. See Universal Patient Care Protocol UP-1.
  - Otherwise contact medical control.
- **Hypoglycemia with Oral Agents:**
  - Patients taking oral diabetic medications should be strongly encouraged to allow transportation to a medical facility.
  - They are at risk of recurrent hypoglycemia that can be delayed for hours and require close monitoring even after normal blood glucose is established.
  - Not all oral agents have prolonged action so Contact Medical Control or NC Poison Control Center for advice.
  - Patients who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.
- **Hypoglycemia with Insulin Agents:**
  - Many forms of insulin now exist. Longer acting insulin places the patient at risk of recurrent hypoglycemia even after a normal blood glucose is established.
  - Not all insulins have prolonged action so Contact Medical Control for advice. Patients who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.
  - In extreme circumstances with no IV and no response to glucagon, Dextrose 50 % can be administered rectally. Contact medical control for advice.



# Pediatric Hypotension/ Shock (Optional)

## History

- Blood loss
- Fluid loss
- Vomiting
- Diarrhea
- Fever
- Infection

## Signs and Symptoms

- Restlessness, confusion, weakness
- Dizziness
- Tachycardia
- Hypotension (Late sign)
- Pale, cool, clammy skin
- Delayed capillary refill
- Dark-tarry stools

## Differential

- Shock
  - Hypovolemic
  - Cardiogenic
  - Septic
  - Neurogenic
  - Anaphylactic
- Trauma
- Infection
- Dehydration
- Congenital heart disease
- Medication or Toxin

	Blood Glucose Analysis Procedure
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Pediatric Airway Protocol(s) <i>if indicated</i>
	Diabetic Protocol PM 2 <i>if indicated</i>

## Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60  
Ages  $\geq$  1 month: SBP < 70  
Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90  
Ages  $\geq$  65: SBP < 110

All ages Shock Index:  
HR > SBP

History and Exam Suggest Type of Shock

Cardiogenic

Hypovolemic

Distributive

Obstructive

Chest Pain: Cardiac and STEMI  
Protocol AC 4  
Appropriate Pediatric Arrhythmia Protocol(s)  
*if indicated*

**A** Normal Saline Bolus  
5 – 10 mL / kg IV / IO  
Titrate to age appropriate  
SBP  $\geq$  70 + (2 x Age)  
Maximum 10 mL / kg

Pediatric Allergy Protocol PM 1  
*if indicated*

Suspected Sepsis Protocol UP 15  
*if indicated*

Multiple Trauma Protocol TB 6  
*if indicated*

**P** Chest Decompression-Needle Procedure  
*if indicated*

**A** Normal Saline Bolus  
20 mL / kg IV / IO  
Titrate to age appropriate  
SBP  $\geq$  70 + (2 x Age)  
Maximum 60 mL / kg

**P** Vasopressor(s)

Notify Destination or Contact Medical Control



# Pediatric Hypotension/ Shock (Optional)

## Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- Consider all possible causes of shock and treat per appropriate protocol. Majority of decompensation in pediatrics is airway or respiratory related.
- Decreasing heart rate and hypotension occur late in children and are signs of impending cardiac arrest.
- Shock may be present with a normal blood pressure initially or even elevated.
- Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the first and only sign.
- Consider all possible causes of shock and treat per appropriate protocol.

### Hypovolemic Shock:

Hemorrhage, trauma, GI bleeding, or pregnancy-related bleeding.

### Tranexamic Acid (TXA):

Agencies utilizing TXA must submit letters from their receiving trauma centers for approval by the OEMS Medical Director.

Receiving trauma centers must agree to continue TXA therapy with repeat dosing.

TXA is NOT indicated and should NOT be administered where trauma occurred > 3 hours prior to EMS arrival.

- Cardiogenic Shock:

Heart failure: MI, Cardiomyopathy, Myocardial contusion, Ruptured ventricle/ septum/ valve/ toxins.

- Distributive Shock:

Septic/ Anaphylactic/ Neurogenic/ Toxic

Hallmark is warm, dry, pink skin with normal capillary refill time and typically alert.

- Obstructive Shock:

Pericardial tamponade. Pulmonary embolus. Tension pneumothorax.

Signs may include hypotension with distended neck veins, tachycardia, unilateral decreased breath sounds or muffled heart sounds.

- Acute Adrenal Insufficiency or Congenital Adrenal Hyperplasia:

Body cannot produce enough steroids (glucocorticoids/ mineralocorticoids.)

May have primary or secondary adrenal disease, congenital adrenal hyperplasia, or more commonly have stopped a steroid like prednisone. Injury or illness may precipitate.

Usually hypotensive with nausea, vomiting, dehydration and/ or abdominal pain.


If suspected, AEMT or Paramedic should give Methylprednisolone 125 mg IM / IV / IO or Dexamethasone 10 mg IM / IV / IO. Use steroid agent specific to your drug list.

May administer prescribed steroid carried by patient IM / IV / IO. Patient may have Hydrocortisone (Cortef or Solu-Cortef). Dose: < 1y.o. give 25 mg, 1-12 y.o. give 50 mg, and > 12 y.o. give 100 mg or dose specified by patient's physician.



# Suspected Viral Hemorrhagic Fever

## Ebola



# Suspected Viral Hemorrhagic Fever Ebola

PARTICULAR ATTENTION MUST BE PAID TO PROTECTING MUCOUS MEMBRANES OF THE EYES, NOSE, and MOUTH FROM SPLASHES OF INFECTIOUS MATERIAL OR SELF INOCULATION FROM SOILED PPE / GLOVES.

THERE SHOULD BE NO EXPOSED SKIN

**DONNING PPE: BEFORE** you enter the patient area.

## Recommended PPE

**PAPR:** A PAPR with a full face shield, helmet, or headpiece. Any reusable helmet or headpiece must be covered with a single-use (disposable) hood that extends to the shoulders and fully covers the neck and is compatible with the selected PAPR.

**N95 Respirator:** Single-use (disposable) N95 respirator in combination with single-use (disposable) surgical hood extending to shoulders and single-use (disposable) full face shield. If N95 respirators are used instead of PAPRs, careful observation is required to ensure healthcare workers are not inadvertently touching their faces under the face shield during patient care.

**Single-use (disposable) fluid-resistant or impermeable gown** that extends to at least mid-calf or coverall without integrated hood. Coveralls with or without integrated socks are acceptable.

**Single-use (disposable) nitrile examination gloves with extended cuffs.** Two pairs of gloves should be worn. At a minimum, outer gloves should have extended cuffs.

**Single-use (disposable), fluid-resistant or impermeable boot covers** that extend to at least mid-calf or single-use (disposable) shoe covers. Boot and shoe covers should allow for ease of movement and not present a slip hazard to the worker.

**Single-use (disposable) fluid-resistant or impermeable shoe covers** are acceptable only if they will be used in combination with a coverall with integrated socks.

**Single-use (disposable), fluid-resistant or impermeable apron that covers** the torso to the level of the mid-calf should be used if Ebola patients have vomiting or diarrhea. An apron provides additional protection against exposure of the front of the body to body fluids or excrement. If a PAPR will be worn, consider selecting an apron that ties behind the neck to facilitate easier removal during the doffing procedure

## DOFFING PPE: OUTSIDE OF PPE IS CONTAMINATED! DO NOT TOUCH

1) PPE must be carefully removed without contaminating one's eyes, mucous membranes, or clothing with potentially infectious materials.

Use great care while doffing your PPE so as not to contaminate yourself (e.g. Do not remove your N-95 facemask or eye protection BEFORE you remove your gown). There should be a dedicated monitor to observe donning and doffing of PPE. It is very easy for personnel to contaminate themselves when doffing. A dedicated monitor should observe doffing to insure it is done correctly. Follow CDC guidance on doffing.

2) PPE must be double bagged and placed into a regulated medical waste container and disposed of in an appropriate location.

3) Appropriate PPE must be worn while decontaminating / disinfecting EMS equipment or unit.

3) Re-useable PPE should be cleaned and disinfected according to the manufacturer's reprocessing instructions.

Hand Hygiene should be performed by washing with soap and water with hand friction for a minimum of 20 seconds.

Alcohol-based hand rubs may be used if soap and water are not available.

EVEN IF AN ALCOHOL-BASED HAND RUB IS USED, WASH HANDS WITH SOAP AND WATER AS SOON AS FEASIBLE.

## THE USE OF GLOVES IS NOT A SUBSTITUTE FOR HAND WASHING WITH SOAP & WATER


For any provider exposure or contamination contact occupational health.

If the patient is being transported via stretcher then a disposable sheet can be placed over them.

## Pearls

- **Transmission to another individual is the greatest after a patient develops fever. Once there is fever, the viral load in the bodily fluids appears to be very high and thus a heightened level of PPE is required.**
- **Patient contact precautions are the most important consideration.**
- **Incubation period 2-21 days**
- **Ebola must be taken seriously; however using your training, protocols, procedures and proper Personal Protective Equipment (PPE), patients can be cared for safely.**
- When an infection does occur in humans, the virus can be spread in several ways to others. The virus is spread through direct contact (through broken skin or mucous membranes) with a sick person's blood or body fluids (urine, saliva, feces, vomit, and semen) objects (such as needles) that have been contaminated with infected body fluids.
- Limit the use of needles and other sharps as much as possible. All needles and sharps should be handled with extreme care and disposed in puncture-proof, sealed containers. Safety devices must be employed immediately after use.
- **Ebola Information:** For a complete review of Ebola go to:  
<http://www.cdc.gov/vhf/ebola/index.html>  
<https://www.cdc.gov/vhf/ebola/clinicians/emergency-services/ems-systems.html>

# Suspected Viral Hemorrhagic Fever Ebola



## Pearls

- **Ebola Information:** For a complete review of Ebola EMS Vehicle Disinfection go to:  
<https://www.cdc.gov/vhf/ebola/clinicians/emergency-services/ems-systems.html>

# Suspected Viral Hemorrhagic Fever Ebola

## Special Circumstances Section

Decedent Known or suspected carrier of HVF / Ebola Requires Transportation

**Only personnel trained in handling infected human remains, and wearing full PPE, should touch, or move any Ebola-infected remains.**

**Handling human remains should be kept to a minimum.**

Donning / Doffing PPE

**PPE should be in place BEFORE contact with the body**

- 1) Prior to contact with body, postmortem care personnel must wear PPE consisting of: surgical scrub suit, surgical cap, impervious Tyvek-Coveralls, eye protection (e.g., face shield, goggles), facemask, shoe covers, and double surgical gloves.
- 2) Additional PPE (leg coverings,) might be required in certain situations (e.g., copious amounts of blood, vomit, feces, or other body fluids that can contaminate the environment).

**PPE should be removed immediately after and discarded as regulated medical waste.**

- 1) Use caution when removing PPE as to avoid contaminating the wearer.
- 2) Hand hygiene (washing your hands thoroughly with soap and water or an alcohol based hand rub) should be performed immediately following the removal of PPE. If hands are visibly soiled, use soap and water.

Preparation of Body Prior to Transport

- 1) At the site of death, the body should be wrapped in a plastic shroud. Wrapping of the body should be done in a way that prevents contamination of the outside of the shroud.
- 2) Change your gown or gloves if they become heavily contaminated with blood or body fluids.
- 3) Leave any intravenous lines or endotracheal tubes that may be present in place.
- 4) Avoid washing or cleaning the body.
- 5) After wrapping, the body should be immediately placed in a leak-proof plastic bag not less than 150 µm thick and zippered closed. The bagged body should then be placed in another leak-proof plastic bag not less than 150 µm thick and zippered closed before being transported to the morgue.

Surface Decontamination

- 1) Prior to transport to the morgue, perform surface decontamination of the corpse-containing body bags by removing visible soil on outer bag surfaces with EPA-registered disinfectants which can kill a wide range of viruses.
- 2) Follow the product's label instructions. Once the visible soil has been removed, reapply the disinfectant to the entire bag surface and allow to air dry.
- 3) Following the removal of the body, the patient room should be cleaned and disinfected.
- 4) Reusable equipment should be cleaned and disinfected according to standard procedures.

Transportation of VHV / Ebola Remains

PPE is required for individuals driving or riding in a vehicle carrying human remains. DO NOT handle the remains of a suspected / confirmed case of Ebola. The remains must be safely contained in a body bag where the outer surface of the body bag has been disinfected prior to the transport.

### Pearls

- **Ebola Information:** For a complete review of Handling Remains of Ebola Infected Patients go to:


<http://www.cdc.gov/vhf/ebola/hcp/guidance-safe-handling-human-remains-ebola-patients-us-hospitals-mortuaries.html>



# High Consequence Pathogens

## (Respiratory Diseases, SARS, MERS-CoV, COVID-19)

### Special Circumstances Section



# High Consequence Pathogens

## (Respiratory Diseases, SARS, MERS-CoV, COVID-19)

### Pearls

- **First Responders:** Because community spread is now present, every patient contact should be considered to have potential for infection with COVID-19. Limit number of FR when caring for patients to limit exposures and PPE use.
- **Place facemask on any patient complaining of respiratory problems with or without a fever.**
- **Dispatch Screening:**
  - If caller interrogation results in positive screen first responders are assigned based on local agency direction.
  - This screening process will result in many False Positive screens in order to be very sensitive.
- **First Responder and EMS Screening:**
  - Limit distance initially to  $\geq$  6 feet and conduct a quick screening using the EMD specific question. If this results in a positive screen, immediately place a facemask on the source patient and all providers don appropriate PPE and limit provider number to that which necessary for patient care.
- **Close Contact and Duration Definition:**
  - Healthcare provider exposure is defined as being within 6 feet for  $\geq$  15 minutes in a patient with suspected illness.
  - Unprotected (no or incorrect PPE) with direct contact with body fluids, including respiratory generated body fluids.
- **Transport:**
  - Occupants in cab of vehicle all should wear facemasks. Riders should be discouraged in order to limit PPE use.
  - Limit number of providers in vehicle required to provide patient care in order to limit exposures.
  - Ensure use of correct PPE for crew and passengers when aerosol-producing procedures utilized.
- **Recommend facemask and gloves with every patient contact. It is reasonable to wear eye protection on every patient contact.**
- **Reasonable to wear simple/surgical mask during entire duty-shift when not able to maintain social distance of  $\geq$  6 feet among fellow providers when not engaged in patient care.**
- **Negative Pressure in care compartment:**
  - Door or window available to separate driver's and care compartment space:**
    - Close door/window between driver's and care compartment and operate rear exhaust fan on full.
  - No door or window available to separate driver's and care compartment space:**
    - Open outside air vent in driver's compartment and set rear exhaust fan to full.
    - Set vehicle ventilation system to non-recirculating to bring in maximum outside air.
    - Use recirculating HEPA ventilation system if equipped.
- **Airborne precautions:**
  - Standard PPE with fit-tested N95 mask (or PAPR respirator) and utilization of a gown or coveralls, change of gloves after every patient contact, and strict hand washing precautions. This level is utilized with Aspergillus, SARS/MERS/COVID-19, Tuberculosis, Measles (rubeola) Chickenpox (varicella-zoster), Smallpox, Influenza, disseminated herpes zoster, or Adenovirus/Rhinovirus.
- **Contact precautions:**
  - Standard PPE with utilization of a gown or coveralls, change of gloves after every patient contact, and strict hand washing precautions. This level is utilized with GI complaints, blood or body fluids, C diff, scabies, wound and skin infections, MRSA. Clostridium difficile (C diff) is not inactivated by alcohol-based cleaners and washing with soap and water is indicated.
- **Droplet precautions:**
  - Standard PPE plus a standard surgical mask for providers who accompany patients in the treatment compartment and a surgical mask or NRB O2 mask for the patient.
  - This level is utilized when Influenza, Meningitis, Mumps, Streptococcal pharyngitis, Pertussis, Adenovirus, Rhinovirus, and undiagnosed rashes.
- **All-hazards precautions:**
  - Standard PPE plus airborne precautions plus contact precautions.
  - This level is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. SARS, MERS-CoV, COVID-19).
- **COVID-19 (Novel Coronavirus): For most current criteria to guide evaluations of patients under investigation:**  
<https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>

# High Consequence Pathogens

## (Respiratory Diseases, SARS, MERS-CoV, COVID-19)

### Decontamination Recommendations

#### EMS Personnel Requires Decontamination

##### Driver:

- Should wear full PPE as described when caring for patient.
- Remove all PPE, except respiratory (N95, PAPR, or equivalent) and perform hand hygiene prior to entering cab to prevent contamination of driver's compartment. **Cab occupants only need to wear facemasks if respirator not already used.**

##### Wash hands:

- Thoroughly after transferring patient care and/or cleaning ambulance

##### Maintain records:

- All prehospital providers exposed to patient at the scene and during ambulance transport (self-monitoring for symptoms for 14 days is recommended, even if wearing appropriate PPE).  
**This does not mean the providers can no longer work.**
- List all prehospital provider names (students, observers, supervisors, first response etc.) in the Patient Care Report.

#### EMS Equipment / Transport Unit Requires Decontamination

##### Safely clean vehicles used for transport:

- Follow standard operating procedures for the containment and disposal of regulated medical waste.
- Follow standard operating procedures for containing and reprocessing used linen.

##### Wear appropriate PPE when:

- Removing soiled linen from the vehicle. Avoid shaking the linen.
- Clean and disinfect the vehicle in accordance with agency standard operating procedures.
- Personnel performing the cleaning should wear a disposable gown and gloves (a respirator should not be needed) during the clean-up process; the PPE should be discarded after use.
- All surfaces that may have come in contact with the patient or materials contaminated during patient care (e.g., stretcher, rails, control panels, floors, walls, work surfaces) should be thoroughly cleaned and disinfected using an **EPA-registered disinfectant** appropriate for SARS, MERS-CoV, or coronavirus in healthcare settings in accordance with manufacturer's recommendations. **Keep doors open to patient care compartment while cleaning to allow air exchanges.**

### EMS Provider Exposure Risk and Monitoring Recommendations

Close Contact				Close Contact			
Less than 6 feet for ≥ 15 minutes Source patient NOT WEARING A MASK				Less than 6 feet for ≥ 15 minutes Source patient WEARING A MASK			
PPE Utilized	Exposure Risk	Monitoring	Work Restrictions	PPE Utilized	Exposure Risk	Monitoring	Work Restrictions
NONE	HIGH		<b>If symptomatic:</b> Fever and Respiratory symptoms (cough, difficulty breathing or other respiratory symptoms) <b>THEN</b> <b>Exclude from work:</b>	NONE	MEDIUM		<b>If symptomatic:</b> Fever and Respiratory symptoms (cough, difficulty breathing or other respiratory symptoms) <b>THEN</b> <b>Exclude from work:</b>
No facemask N95 or PAPR	HIGH			No facemask N95 or PAPR	MEDIUM		
No Eye Protection	MEDIUM	Self-monitor Supervision	<b>THEN</b> <b>Exclude from work:</b> <ul style="list-style-type: none"> <li>At least 72 hours after fever resolution with no use of fever reducing medications.</li> <li>AND</li> <li>At least 10 days since symptom onset.</li> </ul>	No Eye Protection	LOW	Self-monitor Supervision	<b>THEN</b> <b>Exclude from work:</b> <ul style="list-style-type: none"> <li>At least 72 hours after fever resolution with no use of fever reducing medications.</li> <li>AND</li> <li>At least 10 days since symptom onset.</li> </ul>
No Gown/ Coveralls or Gloves	LOW			No Gown/ Coveralls or Gloves	LOW		
All recommended PPE Except facemask instead of N95 or PAPR	LOW			All recommended PPE Except facemask instead of N95 or PAPR	LOW		

Placing a simple/surgical mask on the patient within 15 minutes of contact decreases exposure risk.

#### Return to Work Practice and Work Restrictions (if excluded from work OR exposure to suspected or known COVID-19 patient):

- Prior to duty shift, measure temperature and assess for illness symptoms either by provider, infection control officer, or occupational or public health.
- Self-monitoring with oversight by agency's infection control officer, occupation or public health department per agency policy.
- Wear mask at all times and restrict care of immunocompromised patients (Cancer, Transplant, Steroid use) until all symptoms have resolved or 14 days after onset of illness, whichever is longest.
- Social distance: Employee should maintain 6 feet of separation as work duties permit in the workspace.
- Remove from work if employee becomes symptomatic.
- <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-for-ems.html>
- <https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/hcp-return-work.html>
- <https://www.epa.gov/pesticide-registration/list-n-disinfectants-use-against-sars-cov-2-covid-19>

# High Consequence Pathogens

## (Respiratory Diseases, SARS, MERS-CoV, COVID-19)

### First Responder Guidance

#### **COVID-19 Declared Pandemic with both State and Federal Emergencies Declared**

- Many systems are heavily dependent on First Responder agencies to supplement critical prehospital medical care services.
- Community spread is now evident both in NC and in the US.
- Every patient, regardless of medical or injury complaint, is at risk of COVID-19 and all should undergo routine screening questions.
- While EMD is a first step, all providers must screen every patient contact and don appropriate PPE based on clinical situation and COVID-19 screening.
- The citizens we serve continue to have a variety of illness and injury unrelated to COVID-19.

- Limiting PPE use:

First Responders should consider staging with all incidents and sending 1 provider (or more dependent on situation) into the scene to assess for fever and respiratory complaints.

Request staged resources as needed only to provide necessary medical care.

Where patients do not require immediate intervention, first responders may stay in contact with patient, but remain beyond 6 feet until EMS providers arrive to begin assessment and further care.

Consider calling patient on mobile phones to maintain contact and provide reassurance and explain current situation.

### PPE Crisis or Alternative Strategies

#### **N95 Respirators**

- Use only for aerosol generating procedures (Nebulizer, NIPPV, Suctioning, BVM, BIAD, Intubation).
- Use facemasks in all other scenarios.
- Use respirators (N95 or equivalent) beyond the manufacturing expiration date when not soiled, ripped, torn, or otherwise damaged. Securing straps should also be in good repair and operational:
  - Visually inspect straps, nose bridge/foam, and mask in general.
  - Perform seal check: <https://www.youtube.com/watch?v=pGXuUyAoEd8>
- Models tested by CDC and are believed to function properly beyond expiration date:

3M: 1860, 1860s, 1870, 8210, 9010, 8000	Medline/Alpha Protech NON27501
Gerson 1730	Moldex: 1512, 2201
- Minimize providers caring for patient to the extent possible to conserve.
- Use Self-Contained Breathing Apparatus (SCBA) if needed.
- Re-use respiratory (N95 or equivalent) masks and place in paper bag between use. Do not touch inside of mask. Wash hands thoroughly before removing mask.
- When to discard a respirator (N95 or equivalent):
  - After using during an aerosol producing procedure.
  - Contamination with blood, body fluids or secretions, following close contact with known COVID-19 patient.


#### **Gowns:**

- Use only for aerosol generating procedures (Nebulizer, NIPPV, Suctioning, BVM, BIAD, Intubation).
- Use only for close patient contact, lifting, moving, or transferring where provider contacts patients body.
- May use removable and washable coveralls.

<https://www.cdc.gov/coronavirus/2019-ncov/hcp/respirators-strategy/index.html>



# Hospice or Palliative Care Patient (Optional)





# Hospice or Palliative Care Patient (Optional)

Acute Pain / Air Hunger:

Severity	Medication		
	Morphine (IV/IM/SQ)	Dilauidid (IV/IM/SQ)	Fentanyl (IV/IM/SQ)
Mild	2 mg	0.5 mg	25 mcg
Moderate	4 mg	1 mg	50 mcg
Severe	8 mg	2 mg	100 mcg
Titration	2 mg q 15 minutes IV	0.5 mg q 15 minutes IV	25 mcg q 15 minutes IV

**Due to pain associated with IM injection, IM administration should only be used if alternative medications or routes of administration are not available. PICC lines may be accessed for use by EMS with sterile techniques. May access port-a-cath if appropriate equipment is available and provider is trained.**

If using IM or SQ injections, delay repeat dosing by 30 minutes to prevent dose stacking.

**Consider using moderate / severe dose in opiate tolerant patients:**

Opiate tolerant patients have typical daily dose of narcotic is equivalent to  $\geq 60$  mg of oral Morphine per day (60 OME (Oral Morphine Equivalents)).

**Examples of opiate dosages equivalent to 60 mg of oral Morphine:**

40 mg/day of Oxycodone	60 mg/day Hydrocodone
25 mcg/hr Fentanyl Transdermal	15 mg/day of Methadone
200 mg/day of Tapentadol	16 mg/day of Oxymorphone
Suboxone	

Consider total use of multiple types of opiates. If in doubt about the patient's level of opiate tolerance, or amount of total daily opiate use, treat with a lower initial dose of opiate.

**Anxiety / Agitation:**

Severity	Medication			
	Ativan (IV/IM/SQ)	Versed (IV/IM/SQ)	Valium (IV/IM/SQ)	Haldol (IV/IM/SQ)
Mild / Moderate	0.5 mg	1 mg	2 mg	2 mg
Severe	1 mg	2 mg	5 mg	4 mg

May repeat dose in 15 minutes for IV administration, or 30 minutes for IM or SQ injections.

**Nausea / Vomiting:**

Zofran IV / IM	Phenergan IV / IM	Haldol IV / IM / SQ	Ativan IV / IM / SQ
4 mg	25 mg	2 mg	0.5 mg

**Pearls**

- MOST form Section A and DNR forms are equivalent – if valid, Do Not Resuscitate.**
- MOST form and DNR forms may be revoked by Health Care Power of Attorney or other appropriate surrogate decision-makers.**
- Palliative care is specialized care for patients with a chronic and/ or terminal illness which focuses on managing symptoms exacerbation and the stress of illness.
- Hospice care is specialized care (similar to palliative care) for patients within the last 6 months of life.
- Hospice patient may not have a DNR or MOST form completed and still be enrolled in Hospice care.
- Emergency Kits (eKit):**

May be given to patient by Hospice to use at home for acute symptom exacerbation.

Each eKit is individualized and will be different for each patient, but typically addresses pain, nausea/ vomiting, anxiety, and/ or secretions. (EMS is able to administer if within provider's scope of practice.)

- Interaction on-scene with Hospice personnel:**

Hospice nurses are valuable resources in helping patients/ families make care/ transport decisions.

EMS should discuss care/ transport decision with Hospice nurse.

After medication administration, if no transport occurs, care may be transferred to Hospice nurse.



# Mass Vaccination/Immunization Medication Distribution

## History

- Follow local public health department criteria for specific immunization or medication administered.
- Patient receiving medication or vaccination must be without evidence of active infection.
- AEMT and Paramedic providers may participate
- EMT may participate when DHHS/NCMB allows special provision during local or state emergency.


## Situation

- Local implementation of this protocol must be done as a component of the EMS system's local public health department community immunization or medication distribution program.
- May initiate protocol when a community has limited public health department resources or when local or state health emergency is declared.

## Special Circumstances Section

<b>B</b>	<b>Review immunization/vaccination or medication guide provided by the local public health department:</b> <ul style="list-style-type: none"><li>Patient selection criteria per local public health department (may vary)</li><li>Vaccine/immunization or medication indications</li><li>Vaccine/immunization or medication contraindications</li><li>Vaccine/immunization or medication distribution procedure</li></ul> <ul style="list-style-type: none"><li>EMT may provide vaccinations when DHHS/NCMB allows special provision during local or state emergency.</li></ul>
----------	---

<b>B</b>	<b>Confirm patient eligibility for the vaccination or medication including:</b> <ul style="list-style-type: none"><li>Age</li><li>Medical history</li><li>Contraindications</li><li>Allergies</li></ul>
----------	---



<b>Administer vaccination or medication:</b> <ul style="list-style-type: none"><li>Dose dependent on local public health department</li><li>Route dependent on local public health department (PO, IN, IM, IV, SQ)</li></ul>
--

### Administer Over-the-Counter medication and/or vaccination (if applicable):

- Undergo specific "just-in-time" training
- Dose dependent on local public health department
- Route dependent on local public health department (PO, IN, IM). SQ when specified by NCOEMS.
- Complete required local public health department documentation
- Provide post immunization or medication written instructions and monitoring

<b>Allergic Reaction or Complications</b> 
<ul style="list-style-type: none"><li>Exit to age appropriate Protocol(s)</li><li>Notify appropriate local public health department provider/official</li></ul>

<b>Do not administer:</b> <ul style="list-style-type: none"><li>Refer to local public health department providers/officials for further care and instructions.</li></ul>
--

## Pearls

- Purpose:**  
**Provide protocol driven process for EMS providers to assist with public health immunization or medication distribution initiatives.**
- Documentation of the immunization or medication:**  
Complete using local public health department approved record system.  
Creation of an EMS patient care report is not required and is not required to submit to NCOEMS.  
Must create a log of all patient contacts associated with the immunization or medication distribution program maintained by the EMS system.  
If local public health department is maintaining a log of all patients, EMS may use the public health log and keep copies in the EMS system.
- Injection site:**  
Most common injection site for subcutaneous is tissue of an upper arm; follow procedure USP-4 otherwise.  
Injection volume is limited to 1 - 2 mL per site unless specific guidance is given per local public health department.  
Most common sites for intramuscular injections are upper arm, buttocks, and thighs, follow procedure USP-4.  
Injection volume is limited to 1 mL in the upper arm, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.  
Injection volume is limited to 2 mL (1 mL in pediatrics) in buttocks and thighs, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.

## SC 4



# SARS CoV2

## Monoclonal Antibody Administration

### History

- FDA has issued an Emergency Use Authorization permitting the administration of REGEN-COV (casirivimab and imdevimab) for the treatment of mild to moderate COVID-19.
- Monoclonal antibodies are used to neutralize and prevent progression of the SARS CoV2 virus.

### Situation

- Local implementation of this protocol must be done as a component of the EMS system's local public health department community immunization or medication distribution program.
- May initiate protocol when a community has limited public health department resources or when local or state health emergency is declared.

#### **A** Review monoclonal antibody eligibility criteria:


- [https://www.ems.gov/pdf/EMS\\_Template\\_Protocol\\_for\\_COVID-19\\_Monoclonal\\_Antibody\\_Administration\\_August\\_2021.pdf](https://www.ems.gov/pdf/EMS_Template_Protocol_for_COVID-19_Monoclonal_Antibody_Administration_August_2021.pdf)
- See page 2.
- Determine medication route for either intravenous or subcutaneous administration

#### Allergic Reaction or Complications

- Exit to age appropriate Protocol(s)
- Notify appropriate local public health department provider/official

#### **A** Confirm patient eligibility for monoclonal antibody including:

- Age
- Medical history
- Contraindications
- Allergies



**A** Intravenous Administration:  
Mix  
Casirivimab 600 mg and Imdevimab 600 mg  
In 100 mL NS  
Infuse IV piggyback in a NS line at KVO over 21 minutes (310 mL/hr)  
Infuse through NS primed micron filter 0.20 or 0.22 size

**A** Subcutaneous Administration:  
Draw up  
Casirivimab 600 mg (2.5 mL) in 2 separate syringes and  
Imdevimab 600 mg (2.5 mL) in 2 separate syringes  
Administer 4 SQ injections in separate muscle areas  
Avoid the waistline and 5 cm perumbilically  
  
**Injections: Subcutaneous and Intramuscular Procedure USP - 4**

### Pearls

- Purpose:**  
Provide protocol driven process for EMS providers to assist with public health medication distribution initiatives.
- Documentation of the medication:**  
Creation of an EMS patient care report is required and is required to submit to NCOEMS.  
Must create a log of all patient contacts associated with the medication distribution program maintained by the EMS system unless local public health department is maintaining log. EMS may then retain copies of health department patient logs.
- Injection site:**  
Most common injection site for subcutaneous is tissue of an upper arm; follow procedure USP-4 otherwise.  
Injection volume is limited to 1 - 2 mL per site unless specific guidance is given per local public health department.  
Most common sites for intramuscular injections are upper arm, buttocks, and thighs, follow procedure USP-4.  
Injection volume is limited to 1 mL in the upper arm, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.  
Injection volume is limited to 2 mL (1 mL in pediatrics) in buttocks and thighs, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.

**SC 5**



# SARS CoV2

## Monoclonal Antibody Administration

### Special Circumstances Section

#### Eligibility criteria:

- Age > 12 and weight  $\geq$  40 kg.
- Not requiring hospitalization
- Not requiring oxygen therapy
- High risk for disease progression
  - Age  $\geq$  65
  - Obesity
  - Pregnancy
  - Chronic kidney disease
  - Dementia
  - Diabetes
  - Immunocompromised or immunosuppressive treatments
  - Cardiovascular disease (MI, CVA, CHF, hypertension, hyperlipidemia, diabetes)
  - Chronic lung disease (COPD, asthma, interstitial lung disease)
  - Cancer
  - Sickle cell disease
  - Liver disease
  - Neurodevelopmental disorders, metabolic syndromes, or congenital abnormalities
  - Medical technology dependent, tracheostomy, gastrostomy, or NIPPV/ventilator
- Post-Exposure Prophylaxis (PEP)
  - Not fully vaccinated and immunocompromised or taking immunosuppressive
    - Only 1 of 2 doses and/or less than 2 weeks from 2d dose in 2 dose series or weeks from 1<sup>st</sup> does in vaccine only requiring 1 dose.*
  - OR: Individuals at high risk of exposure to a SARS-CoV-2 infected individual
    - (Nursing home or prison resident)*
- High risk of death
  - Age  $\geq$  80
  - Male sex
  - Black and South Asian descent

medications  
*less than 2*




# Scene Rehabilitation: General (Optional)

Injury / Illness / Complaint should be treated using appropriate treatment protocol beyond need for oral or IV hydration.






**Initial Process**


1. Personnel logged into General Rehabilitation Section
2. VS Assessed / Recorded (If HR > 110 then obtain Temp)  
Carbon Monoxide monitoring if indicated
3. Personnel assessed for signs / symptoms
4. Remove PPE, Body Armor, Haz-Mat Suits, Turnout Gear, Other equipment as indicated



Exit to Scene Rehabilitation Responder Protocol



Reassess responder after 20 Minutes in General Rehabilitation Section  
Reassess VS



Responder Cannot Wear Protective Gear  
Extend Rehabilitation Time Until VS Improve

Extend Rehabilitation Time Until VS Improve

Discharge Responder from General Rehabilitation Section  
Reports for Reassignment

## VITAL SIGN CAVEATS

### **Blood Pressure:**

Prone to inaccuracy on scenes. Must be interpreted in context.

Firefighters have elevated blood pressure due to physical exertion and is not typically pathologic.

Firefighters with Systolic BP  $\geq$  160 or Diastolic BP  $\geq$  100 may need extended rehabilitation. However this does not necessarily prevent them from returning to duty.

### **Temperature:**

Firefighters may have increased temperature during rehabilitation.

# Scene Rehabilitation: General (Optional)

## Pearls

- **This protocol is optional and given only as an example. Agencies may and are encouraged to develop their own.**
- **Rehabilitation officer has full authority in deciding when responders may return to duty and may adjust rest / rehabilitation time frames depending on existing conditions.**
- **Rehabilitation goals:**
  - Relief from climatic conditions.
  - Rest, recovery, and hydration prior to incident, during, and following incident.
  - Active and / or passive cooling or warming as needed for incident type and climate conditions.
- **May be utilized with adult responders on fire, law enforcement, rescue, EMS and training scenes.**
- **Responders taking anti-histamines, blood pressure medication, diuretics or stimulants are at increased risk for cold and heat stress.**
- **General indications for rehabilitation:**
  - 20-minute rehabilitation following use of a second 30-minute SCBA, 45-minute SCBA or single 60-minute SCBA cylinder.
  - 20-minute rehabilitation following 40 minutes of intense work without SCBA.
- **General work-rest cycles:**
  - 10-minute self-rehabilitation following use of one 30-minute SCBA cylinder or performing 20 minutes of intense work without SCBA.
- **Serious signs / symptoms:**
  - Chest pain, dizziness, dyspnea, weakness, nausea, or headache.
  - Symptoms of heat stress (cramps) or cold stress.
  - Changes in gait, speech, or behavior.
  - Altered Mental Status.
  - Abnormal Vital Signs per agency SOP or Policy / Procedure.
- **Rehabilitation Section:**
  - Integral function within the Incident Management System.
  - Establish section such that it provides shelter / shade, privacy and freedom from smoke or other hazards
  - Large enough to accommodate expected number of personnel.
  - Separate area to remove PPE.
  - Accessible to EMS transport units and water supply.
  - Away from media agencies and spectators / bystanders.



# Scene Rehabilitation: Responder (Optional)

## Special Operations Section

### Remove:

PPE  
Body Armor  
Chemical Suits  
SCBA  
Turnout Gear  
Other equipment as indicated


### Continue:

Heat and Cold Stress treatment techniques from General Rehab Section

Injury / Illness / Complaint should be treated using appropriate treatment protocol beyond need for oral or IV hydration.



NFPA Age Predicted 85 % Maximum Heart Rate		
20 - 25		170
26 - 30		165
31 - 35		160
36 - 40		155
41 - 45		152
46 - 50		148
51 - 55		140
56 - 60		136
61 - 65		132



### Pearls

- This protocol is optional and given only as an example. Agencies may and are encouraged to develop their own.
- Rehabilitation officer has full authority in deciding when responders may return to duty.
- Utilized when responder is not appropriate for General Rehabilitation Protocol.
- May be utilized with adult responders on fire, law enforcement, rescue, EMS and training scenes.
- Responders taking anti-histamines, blood pressure medication, diuretics or stimulants are at increased risk for cold and heat stress.
- Rehabilitation Section is an integral function within the Incident Management System.
- Establish section such that it provides shelter, privacy and freedom from smoke or other hazards.



# Bites and Envenomations

## History

- Type of bite/ sting
- Description/ photo for identification
- Time, location, size of bite/ sting
- Previous reaction to bite/ sting
- Domestic vs. Wild
- Tetanus and Rabies risk
- Immunocompromised patient

## Signs and Symptoms

- Rash, skin break, wound
- Pain, soft tissue swelling, redness
- Blood oozing from the bite wound
- Evidence of infection
- Shortness of breath, wheezing
- Allergic reaction, hives, itching
- Hypotension or shock


## Differential

- Animal bite
- Human bite
- Snake bite (poisonous)
- Spider bite (poisonous)
- Insect sting / bite (bee, wasp, ant, tick)
- Infection risk
- Rabies risk
- Tetanus risk

Call for help/ additional resources  
Stage until scene safe

	General Wound Care Procedure
	Immobilize Injury
	Remove any constricting clothing/ bands/ jewelry
■■■	IV or IO Access Protocol UP 6 <b>if indicated</b>
■■■	Age Appropriate Trauma Protocol(s) TB 4, 5, 6 <b>if indicated</b>
■■■	Age Appropriate Allergic Reaction/ Anaphylaxis Protocol AM 1/ PM 1 <b>if indicated</b>
■■■	Age Appropriate Hypotension/ Shock Protocol AM 5 / PM 3 <b>if indicated</b>
■■■	Pain Control Protocol UP 11 <b>if indicated</b>
■■■	Extremity Trauma Protocol TB 4 <b>if indicated</b>

Contact  
**Carolinias Poison Control**  
**1-800-222-1222**  
Or  
**Agency Specific Number**





# Bites and Envenomations

## Pearls

- **Recommended Exam:** Mental Status, Skin, Extremities (Location of injury), and a complete Neck, Lung, Heart, Abdomen, Back, and Neuro exam if systemic effects are noted
- Immunocompromised patients are at an increased risk for infection: diabetes, chemotherapy, transplant patients.
- Consider contacting the North Carolina Poison Control Center for guidance (1-800-222-1222).
- Do not put responders in danger attempting to capture an animal or insect for identification purposes.
- Evidence of infection: swelling, redness, drainage, fever, red streaks proximal to wound.
- **Human bites:**
  - Human bites have higher infection rates than animal bites due to normal mouth bacteria.
  - Hand and foot bites have highest rates of infection.
- **Dog/ Cat/ Carnivore bites:**
  - Carnivore bites are much more likely to become infected and all have risk of Rabies exposure.
  - Cat bites may progress to infection rapidly due to a specific bacteria (Pasteurella multecida).
- **Snake bites:**
  - Poisonous snakes in this area are generally of the pit viper family: rattlesnake and copperhead.
  - Coral snake bites are rare: Very little pain but very toxic. "Red on yellow - kill a fellow, red on black - venom lack."
  - Amount of envenomation is variable, generally worse with larger snakes and early in spring.
  - Snake bites are treated based on signs and symptoms and progression.
  - It is not important to attempt to identify the type of snake and attempts may endanger providers.**
  - Do not bring a snake to the facility for identification as accidental bites to providers may occur.**
- **Spider bites:**
  - Black Widow spider bites tend to be minimally painful, but over a few hours, muscular pain and severe abdominal pain may develop (spider is black with red hourglass on belly).
  - Brown Recluse spider bites are minimally painful to painless. Little reaction is noted initially but tissue necrosis at the site of the bite develops over the next few days (brown spider with fiddle shape on back).
- **Animal bite(s) in subjects declining transport to a medical facility for evaluation:**
  - NCGS 130A-196 requires that all animal bites be reported to the local health department even if the bite is by the owner's animal, and even if accidental.
  - Reporting requirements can be satisfied by reporting to local animal control official.



# Carbon Monoxide/ Cyanide

## History

- Smoke inhalation
- Ingestion of cyanide
- Eating large quantity of fruit pits
- Industrial exposure
- Trauma
- Reason: Suicide, criminal, accidental
- Past Medical History
- Time/ Duration of exposure

## Signs and Symptoms

- AMS
- Malaise, weakness, flu like illness
- Dyspnea
- GI Symptoms: N/V; cramping
- Dizziness
- Seizures
- Syncope
- Reddened skin
- Chest pain

## Differential

- Diabetic related
- Infection
- MI
- Anaphylaxis
- Renal failure/ dialysis problem
- Head injury/ trauma
- Co-ingestant or exposures

Immediately Remove from Exposure

Appropriate Airway Protocol(s) 1 - 7  
**as indicated**

High Flow Oxygen

Blood Glucose Analysis Procedure

**B** 12 Lead ECG Procedure

IV or IO Access Protocol UP 6

**P** Cardiac Monitor/ CO Monitor

Altered Mental Status  
Protocol UP 4  
**if indicated**

Age Appropriate  
Diabetic  
Protocol AM 2/ PM 2  
**if indicated**

Age Appropriate  
Multiple Trauma Protocol TB 6  
Head Injury TB 5  
**if indicated**

Age Appropriate  
Hypotension/ Shock  
Protocol AM 5/ PM 3  
**if indicated**

High Suspicion  
of Cyanide

YES P

Hydroxocobalamin 70 mg / kg IV / IO  
Maximum 5 g

*If available*

NO

Continue Care  
Continue High Flow Oxygen

Monitor and Reassess

Notify Destination or  
Contact Medical Control



## Pearls

- **Recommended exam: Neuro, Skin, Heart, Lungs, Abdomen, Extremities**
- **Scene safety is priority.**
- Consider CO and Cyanide with any product of combustion.
- Normal environmental CO level does not exclude CO poisoning.
- Symptoms present with lower CO levels in pregnancy, children, and the elderly.
- Continue high flow oxygen regardless of pulse ox readings.

TE 2



# Drowning

## History


- Submersion in water regardless of depth
- Possible history of trauma
- Slammed into shore wave break
- Duration of submersion/ immersion
- Temperature of water or possibility of hypothermia

## Signs and Symptoms

- Unresponsive
- Mental status changes
- Decreased or absent vital signs
- Foaming/ Vomiting
- Coughing, Wheezing, Rales, Rhonchi, Stridor
- Apnea

## Differential

- Trauma
- Pre-existing medical problem
  - Hypoglycemia
  - Cardiac Dysrhythmia
- Pressure injury (SCUBA diving)
  - Barotrauma
  - Decompression sickness
- Post-immersion syndrome



## Pearls

- Recommended Exam: Respiratory, Mental status, Trauma Survey, Skin, Neuro
- Drowning is the process of experiencing respiratory impairment (any respiratory symptom) from submersion/ immersion in a liquid.
- Begin with BVM ventilations, if patient does not tolerate, then apply appropriate mode of supplemental oxygen.
- Ensure scene safety. Drowning is a leading cause of death among would-be rescuers.
- When feasible, only appropriately trained and certified rescuers should remove patients from areas of danger.
- Regardless of water temperature – resuscitate all patients with known submersion time of  $\leq 25$  minutes.
- Regardless of water temperature – If submersion time  $\geq 1$  hour consider moving to recovery phase instead of rescue.
- Foam is usually present in airway and may be copious, DO NOT waste time attempting to suction. Ventilate with BVM through foam (suction water and vomit only when present.)
- Cardiac arrest in drowning is caused by hypoxia, airway and ventilation are equally important to high-quality CPR.
- Encourage transport of all symptomatic patients (cough, foam, dyspnea, abnormal lung sounds, hypoxia) due to potential worsening over the next 6 hours.
- Predicting prognosis in prehospital setting is difficult and does not correlate with mental status. Unless obvious death, transport.
- Hypothermia is often associated with drowning and submersion injuries even with warm ambient conditions.
- Drowning patient typically has  $<1 - 3$  mL/kg of water in lungs (does not require suction) Primary treatment is reversal of hypoxia.
- Spinal motion restriction is usually unnecessary. When indicated it should not interrupt ventilation, oxygenation and/ or CPR.

TE 3



# Hyperthermia

## History

- Age, very young and old
- Exposure to increased temperatures and / or humidity
- Past medical history / Medications
- Time and duration of exposure
- Poor PO intake, extreme exertion
- Fatigue and / or muscle cramping

## Signs and Symptoms

- Altered mental status / coma
- Hot, dry or sweaty skin
- Hypotension or shock
- Seizures
- Nausea

## Differential

- Fever (Infection)
- Dehydration
- Medications
- Hyperthyroidism (Thyroid Storm)
- Delirium tremens (DT's)
- Heat cramps, exhaustion, stroke
- CNS lesions or tumors

Temperature Measurement  
Procedure  
***if available***

Temperature Measurement  
should NOT delay treatment of  
hyperthermia

Remove from heat source to cool environment

Cooling measures

Remove tight clothing

Blood Glucose Analysis Procedure

Age Appropriate  
Diabetic Protocol AM 2/ PM 2  
***as indicated***

## Heat Stroke

### Classic Heat Stroke

- Not common type
- Hot and Dry
- Altered Mental Status

### Exertional Heat Stroke

- **Most common type**
- Wet with prior sweating
- Altered Mental Status

## HEAT CRAMPS

Normal to elevated body temperature  
Warm, moist skin  
Weakness, Muscle cramping

## HEAT EXHAUSTION

Elevated body temperature  
Cool, moist skin  
Weakness, Anxious, Tachypnea

## HEAT STROKE

Fever, usually > 104°F (40°C)  
Hot, dry skin  
Hypotension, AMS / Coma

PO Fluids as tolerated

Monitor and Reassess

Age Appropriate  
Airway Protocol(s) AR 1 - 7  
***as indicated***

Altered Mental Status  
Protocol UP 4  
***as indicated***

Active cooling measures  
Target Temp < 102.5°F (39°C)

B 12 Lead ECG Procedure

IV or IO Access Protocol UP 6

P Cardiac Monitor

A Normal Saline Bolus  
500 mL IV / IO  
Repeat to effect SBP > 90  
Maximum 2 L  
PED: Bolus 20 mL/kg IV / IO  
Repeat to effect Age appropriate  
SBP  $\geq$  70 + 2 x Age  
Maximum 60 mL/kg

Age Appropriate  
Hypotension/ Shock  
Protocol AM 5/ PM 3  
***as indicated***

Monitor and Reassess

Notify Destination or  
Contact Medical Control



# Hyperthermia

## Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lungs, Neuro
- **Extremes of age are more prone to heat emergencies (i.e. very young and very old).**
- **Temperature measurement:**
  - Obtain and document patient temperature if able.
  - Many thermometers and routes of measurement are available.
  - Order of preference for route of measurement: Rectal > oral > temporal > axillary.
- Heat illness is predisposed by use of: tricyclic antidepressants, phenothiazines, anticholinergic medications, and alcohol.
- Cocaine, Amphetamines, and Salicylates may elevate body temperatures.
- Intense shivering may occur as patient is cooled.
- **Heat Cramps:**
  - Consists of benign muscle cramping secondary to dehydration and is not associated with an elevated temperature.
- **Heat Exhaustion:**
  - Consists of dehydration, salt depletion, dizziness, fever, mental status changes, headache, cramping, nausea and vomiting.
  - Vital signs usually consist of tachycardia, hypotension, and an elevated temperature.
- **Heat Stroke:**
  - Consists of dehydration, tachycardia, hypotension, temperature  $\geq$  104°F (40°C), and an altered mental status.
  - Sweating generally disappears as body temperature rises above 104°F (40°C).
  - The young and elderly are more prone to be dry with no sweating.
  - Exertional Heat Stroke:**
    - In exertional heat stroke (athletes, hard labor), the patient may have sweated profusely and be wet on exam.
    - Rapid cooling takes precedence over transport as early cooling decreases morbidity and mortality.**
    - If available, immerse in an ice water bath for 20 minutes. Monitor rectal temperature and remove patient when temperature reaches 102.5°F (39°C). Your goal is to decrease rectal temperature below 104°F (40°C) with target of 102.5°F (39°C) within 15 minutes. Stirring the water aids in cooling.
    - Nearly 66% of all exertional heat strokes occur in high school athletes during the month of August.
    - In NC, it is mandatory that all high school field houses have a dunk tank and available ice and water.
    - Other methods include cold wet towels below and above the body or spraying cold water over body continuously.
- **Neuroleptic Malignant Syndrome (NMS):**
  - Neuroleptic Malignant Syndrome is a hyperthermic emergency which is not related to heat exposure.
  - It occurs after taking neuroleptic antipsychotic medications.
  - This is a rare but often lethal syndrome characterized by muscular rigidity, AMS, tachycardia and hyperthermia.
  - Drugs Associated with Neuroleptic Malignant Syndrome:**
    - Prochlorperazine (Compazine), promethazine (Phenergan), clozapine (Clozaril), risperidone (Risperdal) metoclopramide (Reglan), amoxapine (Ascendin), and lithium.
  - Management of NMS:**
    - Supportive care with attention to hypotension and volume depletion.
    - Use benzodiazepines such as diazepam or midazolam for seizures and/ or muscular rigidity.



# Hypothermia/ Frostbite

## History

- Age, very young and old
- Exposure to decreased temperatures but may occur in normal temperatures
- Past medical history / Medications
- Drug use: Alcohol, barbituates
- Infections/ Sepsis
- Length of exposure/ Wetness/ Wind chill

## Signs and Symptoms

- Altered mental status/ coma
- Cold, clammy
- Shivering
- Extremity pain or sensory abnormality
- Bradycardia
- Hypotension or shock

## Differential

- Sepsis
- Environmental exposure
- Hypothyroidism
- Hypoglycemia
- CNS dysfunction
- Stroke
- Head injury
- Spinal cord injury

Temperature Measurement  
Procedure  
*if available*

Temperature Measurement  
should NOT delay treatment of  
hypothermia

Remove wet clothing  
Dry/ Warm Patient

Passive warming measures

Blood Glucose Analysis Procedure

Age Appropriate  
Diabetic Protocol AM 2/ PM 2  
*as indicated*

Hypothermia / Frost Bite

Systemic Hypothermia

Localized Cold Injury

General Wound Care

DO NOT Rub Skin to warm  
DO NOT Massage  
Skin to warm

DO NOT allow refreezing

Awake with / without AMS

Age Appropriate  
Airway Protocol(s) AR 1 - 7  
*as indicated*

Altered Mental Status  
Protocol UP 4  
*as indicated*

Active warming measures

B 12 Lead ECG Procedure

IV or IO Access Protocol UP 6

**A** Normal Saline Bolus  
500 mL IV / IO  
Repeat to titrate SBP > 90 mmHg

Maximum 2 L

Pediatric: 20 mL / kg IV / IO  
Repeat to titrate Age Appropriate  
SBP  $\geq$  70 + 2 x Age  
Maximum 60 mL / kg

**P** Cardiac Monitor

Age Appropriate

Hypotension/ Shock

Protocol AM 5/ PM 3

Multiple Trauma Protocol TB 6  
*as indicated*

Monitor and Reassess

Unresponsive

Pulse

YES  
Exit to  
Age Appropriate  
Cardiac Protocol(s)  
*See Pearls*

NO

Notify Destination or  
Contact Medical Control



# Hypothermia/ Frostbite

## Pearls

- Recommended Exam: Mental Status, Heart, Lungs, Abdomen, Extremities, Neuro
- NO PATIENT IS DEAD UNTIL WARM AND DEAD (Body temperature  $\geq 93.2^{\circ}$  F,  $32^{\circ}$  C.)
- Temperature measurement:
  - Obtain and document patient temperature if able.
  - Many thermometers and routes of measurement are available.
  - Order of preference for route of measurement: Rectal > oral > temporal > axillary.
  - Many thermometers do not register temperature below  $93.2^{\circ}$  F.
- Hypothermia categories:
  - Mild  $90 - 95^{\circ}$  F ( $32 - 35^{\circ}$  C)
  - Moderate  $82 - 90^{\circ}$  F ( $28 - 32^{\circ}$  C)
  - Severe  $< 82^{\circ}$  F ( $< 28^{\circ}$  C)
- Mechanisms of hypothermia:
  - Radiation: Heat loss to surrounding objects via infrared energy (60% of most heat loss.)
  - Convection: Direct transfer of heat to the surrounding air.
  - Conduction: Direct transfer of heat to direct contact with cooler objects (important in submersion.)
  - Evaporation: Vaporization of water from sweat or other body water losses.
- Contributing factors of hypothermia: Extremes of age, malnutrition, alcohol or other drug use.
- If the temperature is unable to be measured, treat the patient based on the suspected temperature.
- CPR:
  - Severe hypothermia may cause cardiac instability and rough handling of the patient theoretically can cause ventricular fibrillation. This has not been demonstrated or confirmed by current evidence. Intubation and CPR techniques should not be withheld due to this concern.
  - Intubation can cause ventricular fibrillation, so it should be done gently by the most experienced provider(s).
  - Below  $86^{\circ}$  F ( $30^{\circ}$  C) antiarrhythmics may not work and if given, should be given at increased time intervals. Contact medical control for direction. Epinephrine can be administered.
  - Below  $86^{\circ}$  F ( $30^{\circ}$  C) pacing should not be utilized.
  - Consider withholding CPR if patient has organized rhythm or has other signs of life. Contact Medical Control.
  - If the patient is below  $86^{\circ}$  F ( $30^{\circ}$  C) then defibrillate 1 time if defibrillation is required. Deferring further attempts until more warming occurs is controversial. Contact medical control for direction.
  - Hypothermia may produce severe bradycardia so take at least 60 seconds to palpate a pulse.
- Active Warming:
  - Remove from cold environment and into warm environment protected from wind and wet conditions.
  - Remove wet clothing and provide warm blankets/ warming blankets.
  - Hot packs can be activated and placed in the armpit and groin area if available. Care should be taken not to place the packs directly against the patient's skin.



# Marine Envenomation/ Injury

## History

- Type of bite/ sting
- Identification of organism
- Previous reaction to marine organism
- Immunocompromised
- Household pet

## Signs and Symptoms

- Intense localized pain
- Increased oral secretions
- Nausea/ vomiting
- Abdominal cramping
- Allergic reaction / anaphylaxis


## Differential

- Jellyfish sting
- Sea Urchin sting
- Sting ray barb
- Coral sting
- Swimmers itch
- Cone Shell sting
- Fish bite
- Lion Fish sting

Call for help/ additional resources  
Stage until scene safe

General Wound Care Procedure	
	IV or IO Access Protocol UP 6 <i>if indicated</i>
P	Cardiac Monitor <i>if indicated</i>
	Drowning Protocol TE 3 <i>if indicated</i>
	Age Appropriate Allergy/ Anaphylaxis Protocol AM 1/ PM 1 <i>if indicated</i>
	Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 <i>if indicated</i>
	Pain Control Protocol UP 11 <i>if indicated</i>

Contact  
Carolinas Poison Control  
1-800-222-1222  
Or  
Agency Specific Number





# Marine Envenomation/ Injury

## Pearls

- **Ensure your safety:** Avoid the organism or fragments of the organism as they may impart further sting or injury.
- **Priority is removal of the patient from the water to prevent drowning.**
- **Coral:**
  - Coral is covered by various living organisms which are easily dislodged from the structure.
  - Victim may swim into coral causing small cuts and abrasions and the coral may enter into cuts, causing little if any symptoms initially, but later causing inflammation, pain and/ or infection.
  - The next 24 – 48 hours may reveal an inflammatory reaction with swelling, redness, itching, tenderness, and ulceration.
  - Treatment is flushing with large amounts of fresh water or soapy water then repeating.
- **Jelly Fish/ Anemone/ Man-O-War:**
  - Wash the area with fresh seawater to remove tentacles and nematocysts.
  - Do not apply fresh water or ice as this will cause nematocysts firing as well.
  - Recent evidence does not demonstrate a clear choice of any solution that neutralizes nematocysts.
  - Vinegar (immersion for 30 seconds), 50:50 mixture of Baking Soda and Seawater, and even meat tenderizer may have similar effects.
  - Immersion in warm water for 20 minutes, 110 – 114°F (43 - 46°C), is effective in pain control.
  - Shaving cream may be useful in removing the tentacles and nematocysts with a sharp edge (card).
  - Stimulation of the nematocysts by pressure or rubbing cause the nematocyst to fire even if detached from the jellyfish.
  - Lift away tentacles as scrapping or rubbing will cause nematocysts firing.
  - Typically symptoms are immediate stinging sensation on contact, intensity increases over 10 minutes.
  - Redness and itching usually occur.
  - Papules, vesicles and pustules may be noted and ulcers may form on the skin.
  - Increased oral secretions and gastrointestinal cramping, nausea, pain, or vomiting may occur.
  - Muscle spasm, respiratory, and cardiovascular collapse may follow.
- **Lionfish:**
  - In North Carolina this would typically occur in a residence/ business as lionfish are often kept as pets in saltwater aquariums.
  - Remove any obvious protruding spines and irrigate area with copious amounts of saline.
  - The venom is heat labile so immersion in hot water, 110 – 114°F for 30 to 90 minutes is the treatment of choice but do not delay transport if indicated.
- **Stingrays:**
  - Typical injury is swimmer stepping on ray and muscular tail drives 1 – 4 barbs into victim.
  - Venom released when barb is broken.
  - Typical symptoms are immediate pain which increases over 1 – 2 hours.
  - Bleeding may be profuse due to deep puncture wound.
  - Nausea, vomiting, diarrhea, muscle cramping, and increased urination and salivation may occur.
  - Seizures, hypotension, and respiratory or cardiovascular collapse may occur.
  - Irrigate wound with saline. Extract the spine or barb unless in the abdomen or thorax, Contact Medical Control for advice.
  - Immersion in hot water, if available, for 30 to 90 minutes but do not delay transport.
- Patients can suffer cardiovascular collapse from both the venom and/ or anaphylaxis even in seemingly minor envenomation.
- Sea creature stings and bites impart moderate to severe pain.
- Arrest the envenomation by inactivation of the venom as appropriate.
- Ensure good wound care, immobilization and pain control.



# Overdose/ Toxic Ingestion

## History


- Ingestion or suspected ingestion of a potentially toxic substance
- Substance ingested, route, quantity
- Time of ingestion
- Reason (suicidal, accidental, criminal)
- Available medications in home
- Past medical history, medications

## Signs and Symptoms

- Mental status changes
- Hypotension / hypertension
- Decreased respiratory rate
- Tachycardia, dysrhythmias
- Seizures
- S.L.U.D.G.E.
- D.U.M.B.B.E.L.S


## Differential

- Tricyclic antidepressants (TCAs)
- Acetaminophen (Tylenol)
- Aspirin
- Depressants
- Stimulants
- Anticholinergic
- Cardiac medications
- Solvents, Alcohols, Cleaning agents
- Insecticides (organophosphates)





# Overdose/ Toxic Ingestion





# Overdose/ Toxic Ingestion

## Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro
- **Opioids and opiates** may require higher doses of Naloxone to improve respiration, in certain circumstances up to 10 mg.
- **Time of Ingestion:**
  1. Most important aspect is the **TIME OF INGESTION**, the substance(s), amount ingested, and any co-ingestants.
  2. Every effort should be made to elicit this information before leaving the scene.

- **Charcoal Administration:**

The American Academy of Clinical Toxicology DOES NOT recommend the routine use of charcoal in poisonings.

1. Consider Charcoal within the **FIRST HOUR** after ingestion. If a potentially life threatening substance is ingested or extended release agent(s) are involved and  $\geq$  one hour from ingestion, Contact Medical Control or NC Poison Control Center for direction.
2. If NG would be necessary to administer Charcoal, then DO NOT administer unless known to be adsorbed, airway secured by intubation, and ingestion is less than **ONE HOUR** confirmed and potentially lethal.
3. Charcoal in general, should only be given to a patient who is alert and awake such that they can self-administer the medication.

- Do not rely on patient history of ingestion, especially in suicide attempts. Make sure patient is still not carrying hiding other medications or has any weapons.

- **Pediatric:**

Age specific blood pressure 0 – 28 days  $> 60$  mmHg, 1 month - 1 year  $> 70$  mmHg, 1 - 10 years  $> 70 + (2 \times \text{age})$  mmHg and  $> 10$  years  $> 90$  mmHg.

**Pediatric IV Fluid maintenance rate:**

4 mL for the first 10 kg of weight +  
2 mL for the second 10 kg of weight +  
1 mL for every additional kg in weight after 20 kg

**Example: 34 kg pediatric**

First 10 kg:	4 mL/kg/hr = 40 mL/hr
Second 10 kg:	2 mL/kg/hr = 20 mL/hr
Final 14 Kg:	1 mL/kg/hr = 14 mL/hr

Total: 74 mL/hr rate

- Bring bottles, contents, emesis to ED.
- **S.L.U.D.G.E:** Salivation, Lacrimation, Urination, Defecation, GI distress, Emesis.
- **D.U.M.B.B.E.L.S:** Diarrhea, Urination, Miosis, Bradycardia, Bronchorrhea, Emesis, Lacrimation, Salivation.
- **Tricyclic:** 4 major areas of toxicity: seizures, dysrhythmias, hypotension, decreased mental status or coma; rapid progression from alert mental status to death.
- **Acetaminophen:** initially normal or nausea/ vomiting. If not detected and treated, causes irreversible liver failure.
- **Aspirin:** Early signs consist of abdominal pain and vomiting. Tachypnea and altered mental status may occur later. Renal dysfunction, liver failure, and or cerebral edema among other things can take place later.
- **Depressants:** decreased HR, decreased BP, decreased temperature, decreased respirations, non-specific pupils.
- **Stimulants:** increased HR, increased BP, increased temperature, dilated pupils, seizures.
- **Anticholinergic:** increased HR, increased temperature, dilated pupils, mental status changes.
- **Cardiac Medications:** dysrhythmias and mental status changes.
- **Solvents:** nausea, coughing, vomiting, and mental status changes.
- **Insecticides:** increased or decreased HR, increased secretions, nausea, vomiting, diarrhea, pinpoint pupils.
- **Nerve Agent Antidote kits** contain 2 mg of Atropine and 600 mg of pralidoxime in an autoinjector for self administration or patient care. These kits may be available as part of the domestic preparedness for Weapons of Mass Destruction.
- **EMR and EMT may administer naloxone by IN / IM route only and may administer from EMS supply. Agency medical director may require Contact of Medical Control prior to administration and may restrict locally.**
- **When appropriate contact the North Carolina Poison Control Center for guidance, reference Policy 18.**
- Consider restraints if necessary for patient's and/or personnel's protection per the Restraint Procedure.



# WMD-Nerve Agent Protocol

## History

- Exposure to chemical, biologic, radiologic, or nuclear hazard
- Potential exposure to unknown substance/hazard

## Signs and Symptoms

- Salivation
- Lacration
- Urination; increased, loss of control
- Defecation / Diarrhea
- GI Upset; Abdominal pain / cramping
- Emesis
- Muscle Twitching
- Seizure Activity
- Respiratory Arrest

## Differential

- Nerve agent exposure (e.g., VX, Sarin, Soman, etc.)
- Organophosphate exposure (pesticide)
- Vesicant exposure (e.g., Mustard Gas, etc.)
- Respiratory Irritant Exposure (e.g., Hydrogen Sulfide, Ammonia, Chlorine, etc.)

Call for help/ additional resources  
Stage until scene safe

Obtain history of exposure  
Observe for specific toxidromes  
Initiate triage and/or decontamination as indicated.

Contact  
**Carolinians Poison Control**  
1-800-222-1222  
Or  
Agency Specific Number

Symptom Severity

Asymptomatic

**Minor Symptoms:**  
Respiratory Distress + SLUDGEM

**Major Symptoms:**  
Altered Mental Status, Seizures, Respiratory Distress, Respiratory Arrest

Monitor and Reassess  
Every 15 minutes for symptoms  
Initiate Treatment per Appropriate Arm

IV or IO Access Protocol UP 6  
Nerve Agent Kit IM  
2 Doses Rapidly  
*if available*

IV or IO Access Protocol UP 6  
Nerve Agent Kit IM  
3 Doses Rapidly  
*if available*

Atropine 2 mg IV / IO / IM  
Pediatric: See Pearls  
IV / IO / IM  
Repeat every 3 to 5 minutes until symptoms resolve

Pralidoxime (2PAM) 600 mg IV / IO / IM  
Pediatric: 15 – 25 mg / kg  
IV / IO / IM  
Over 30 minutes

Seizure Protocol UP 13

**CDC/ ASPR CHEMPACK Program**  
NC -57 EMS containers  
-43 locations  
  
Almost all citizens within 50 miles of CHEMPACK  
See Page 2 and Pearls

Multiple Patients

Consider  
Activation and deployment of CHEMPACK  
**CHEMPACK ACTIVATION:**  
(insert local number)  
  
**Healthcare Coalition Activation**  
(insert local number)

YES

NO

Notify Destination or  
Contact Medical Control



# WMD-Nerve Agent Protocol

## Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lungs, Gastrointestinal, Neuro
- **Follow local HAZMAT protocols for decontamination and use of personal protective equipment.**
- **Adult/ Pediatric Atropine Dosing Guides:**

Confirmed attack: Begin with 1 Nerve Agent Kit for patients less than 7 years of age, 2 Nerve Agent Kits from 8 to 14 years of age, and 3 Nerve Agent Kits for patients 15 years of age and over.

If Triage/ MCI issues exhaust supply of Nerve Agent Kits, use pediatric atropines (if available).

Usual pediatric doses:  $0.5 \text{ mg} \leq 40 \text{ pounds (18 kg)}$ , 1 mg dose if patient weighs between 40 to 90 pounds (18 to 40 kg), and 2 mg dose  $\geq 90 \text{ pounds} (\geq 40 \text{ kg})$ .

- **Each Nerve Agent Kit contains 600 mg of Pralidoxime (2-PAM) and 2 mg of Atropine.**

- **Seizure Activity: Any benzodiazepine by any route is acceptable.**

- For patients with major symptoms, there is no limit for atropine dosing.

- Carefully evaluate patients to ensure they do not have exposure to other agent(s) (e.g., narcotics, vesicants, etc.)

- The main symptom that the atropine addresses is excessive secretions, so atropine should be given until secretions improve/ dry.

- EMS personnel, public safety officers and EMR/ EMT may carry, self-administer, or administer atropine/ pralidoxime to others by protocol. Agency medical director may require Contact of Medical Control prior to administration.


- **CHEMPACK Program:**

For multiple patients, call for **CHEMPACK** deployment per local emergency management and healthcare **coalition plans**.

**1 EMS CHEMPACK supports 454 patients.**

Medication in CHEMPACK may be used regardless of expiration date.

EMS Type CHEMPACK Container 454 Person Treatment Capacity			
Product	Cases	Units per case	Total Units
Mark 1 Auto-injector	5	240	1,200
-OR-			
ATNAA Auto-injector	6	200	1,200
-OR-			
Atropen 2mg Auto-injector	9	136	1,224
Pralidoxime 300mg Auto-injector	5	240	1,200
-AND-			
Diazepam 10mg Auto-injector	2	300	600
Seizalam (Midazolam) 5mg/ml vial 10ml	1	100	100
Atropen 0.5mg Auto-injector	1	225	225
Atropen 1mg Auto-injector	1	225	225
Atropine Sulfate 0.4mg/ml vial 20ml	1	100	100
Pralidoxime 1gm inj. 20ml	1	276	276
Sterile Water 20ml vials	1	150	150





# Blast Injury/ Incident

## History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history/ Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status

## Signs and Symptoms

- Hearing loss (TM rupture)
- Ocular burns/vision changes
- Multiple trauma/ penetrating trauma
- Hypotension/ shock
- Airway compromise/distress could be indicated by hoarseness/ wheezing
- Pneumothorax/ hemothorax
- Traumatic amputation (tourniquet)

## Differential

- Thermal / Chemical / Electrical Burn Injury  
Superficial  
(1<sup>st</sup> Degree) red – painful  
(Don't include in TBSA)
- Partial Thickness  
(2<sup>nd</sup> Degree) blistering
- Full Thickness  
(3<sup>rd</sup> Degree) painless/charred or leathery skin
- Radiation injury

**Nature of Device:** Agent/ Amount. Industrial Explosion. Terrorist Incident. Improvised Explosive Device.

**Method of Delivery:** Incendiary/ Explosive

**Nature of Environment:** Open / Closed.

**Distance from Device:** Intervening protective barrier. Other environmental hazards,

**Evaluate for:** Blunt Trauma/ Crush Injury/ Compartment Syndrome/ Traumatic Brain Injury/ Concussion/ Tympanic Membrane Rupture/ Abdominal hemorrhage or Evisceration, Blast Lung Injury and Penetrating Trauma.

## Scene Safety/ Quantify number and Triage Patients/ Load and Go with Assessment/ Treatment Enroute

Call for help/ additional resources  
Stage until scene safe

Accidental/ Intentional Explosions  
(See Pearls)

	Triage Protocol UP 2 <i>as indicated</i>
	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 <i>as indicated</i>
	Multiple Trauma Protocol TB 6 <i>if indicated</i>
	IV and IO Access Protocol UP 6 <i>if indicated</i>
P	Cardiac Monitor <i>if indicated</i>
	Thermal Burn Protocol TB 9 Chemical and Electrical Burn Protocol TB 2 <i>if indicated</i>
	Crush Injury Protocol TB 3 <i>if indicated</i>
	Radiation Incident Protocol TB 7 <i>if indicated</i>
	Decontamination Procedure USP 2 <i>if indicated</i>
	Pain Control Protocol UP 11 <i>if indicated</i>



Rapid Transport to appropriate destination using  
Trauma and Burn:  
EMS Triage and Destination Plan

Notify Destination or  
Contact Medical Control



# Blast Injury/ Incident

## Pearls

- **Types of Blast Injury:**

Primary Blast Injury: From the blast pressure (air) wave.

Secondary Blast Injury: Impaled objects. Debris which becomes missiles/ shrapnel.

Tertiary Blast Injury: Patient falling or being thrown/ pinned by debris.

Most Common Cause of Death: Secondary Blast Injuries.

- **Triage of Blast Injury patients:**

Blast Injury patients with burn injuries should be triaged using the Thermal Burn/ Chemical and Electrical Burn Protocol Guidelines for Critical/ Serious/ Minor Trauma and Burns and the Trauma and Burn: EMS Triage and Destination Plan.

Patients may be hard of hearing due to tympanic membrane rupture.

- **Care of Blast Injury Patients:**

Patients may suffer multi-system injuries including blunt and penetrating trauma, shrapnel, barotrauma, burns, and toxic chemical exposure.

Consider airway burns, which should prompt early and aggressive airway management as indicated.

Cover open chest wounds with semi-occlusive dressing or commercial chest seal product.

Use Lactated Ringers (if available) for all Critical or Serious Burns.

Minimize IV fluids resuscitation in patients with no signs of shock or poor perfusion.

- **Blast Lung Injury:**

Blast Lung Injury is characterized by respiratory difficulty and hypoxia. Can occur (rarely) in patients without external thoracic trauma. More likely to occur in an enclosed space or in close proximity to explosion.

Symptoms: Dyspnea, hemoptysis, cough, chest pain, wheezing, and hemodynamic instability.

Signs: Apnea, tachypnea, hypopnea, hypoxia, cyanosis, and diminished breath sounds.

Air embolism should be considered and patient transported in left-lateral decubitus position.

Blast Lung Injury patients may require early intubation but positive pressure ventilation may worsen the injury, avoid hyperventilation, which can cause further injury.

Air transport may worsen lung injury, monitor oxygenation and ventilation closely. Tension pneumothorax may occur requiring chest decompression. Be judicious with fluids as volume overload may worsen lung injury.

- **Accidental Explosions or Intentional Explosions:**

**All explosions or blasts should be considered intentional until determined otherwise.**

**Greatest concern is potential threat for a secondary device.**

Attempt to determine the source of the blast to include any potential threat for aerosolization of hazardous materials.

Evaluate scene safety including the source of the blast, which may continue to spill explosive liquids or gases.

Consider structural collapse, environmental hazard, and fire.

Conditions that led to the initial explosion may reoccur and lead to a second explosion.

Patients who physically able, typically will attempt to move as far away from the explosive source.

Evaluate surroundings for suspicious items; unattended back packs or packages, or unattended vehicles.

**If patient(s) is unconscious or there is fatalities and you are evaluating patient(s) for signs of life:**

**Before moving, note if there are wires coming from the patient(s), or if it appears the patient(s) is lying on a package/ pack, or bulky item. If so, do not move the patient(s), quickly back away and immediately notify a law enforcement officer.**

If there are no indications the patient is connected to a triggering mechanism for a secondary device, expeditiously remove the patient(s) from the scene and begin transport to the hospital.

Protect the airway and cervical spine, however beyond the primary survey, care and a more detailed assessment should be deferred until rapid transport begins.

If there are signs the patient was carrying the source of the blast, notify law enforcement immediately, and most likely a law enforcement officer will accompany your patient to the hospital.



# Chemical and Electrical Burn

## History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history/ Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status

## Signs and Symptoms

- Burns, pain, swelling
- Ocular burns/ vision changes
- Loss of consciousness
- Hypotension/ shock
- Compartment syndrome
- Airway compromise/ distress could be indicated by hoarseness/ wheezing
- Electrical burn may be misleading with small contact/ external burn and major internal injury – burn/ trauma center transport is recommended

## Differential

- Thermal / Chemical / Electrical Burn Injury
- Superficial (1<sup>st</sup> Degree) red – painful (Don't include in TBSA)
- Partial Thickness (2<sup>nd</sup> Degree) blistering
- Full Thickness (3<sup>rd</sup> Degree) painless/charred or leathery skin
- Radiation injury
- Blast injury

**Assure Chemical Source is NOT Hazardous to Responders.**

**Assure Electrical Source is NO longer in contact with patient before touching patient.**

## Assess Burn/ Concomitant Injury Severity

**< 5% TBSA 2<sup>nd</sup>/3<sup>rd</sup> Degree Burn**  
No inhalation injury, Not Intubated, Normotensive GCS 14 or Greater Minor Burn

**5-15% TBSA 2<sup>nd</sup>/3<sup>rd</sup> Degree Burn**  
Suspected inhalation injury or requiring intubation for airway stabilization Hypotension or GCS 13 or Less (When reasonably accessible, transport to a Burn Center) Serious Burn

**>15% TBSA 2<sup>nd</sup>/3<sup>rd</sup> Degree Burn**  
Burns with Multiple Trauma Burns with definitive airway compromise (When reasonably accessible, transport to a Burn Center) Critical Burn

<p><b>Age Appropriate Airway Protocol(s) AR 1, 2, 3, 4, 5, 6, 7 if indicated</b></p> <p></p> <p>IV or IO Access Protocol UP 6 Consider 2 IV sites if <math>\geq 15\%</math> TBSA</p> <p></p> <p>Thermal Burn Protocol TB 9</p> <p></p> <p>Pain Control Protocol UP 11 <b>if indicated</b></p>	
	<p>Identify Contact Points</p> <p><b>Eye Involvement</b> Irrigate Involved Eye(s) with Normal Saline + for 30 minutes Continue irrigation during transport</p> <p><b>Chemical Exposure/ Burn</b> Flush Contact Area with Normal Saline for 15 minutes Continue irrigation during transport</p> <p>Decontamination Procedure USP 2 <b>if indicated</b></p>
	<p></p> <p>Age Appropriate Cardiac Protocol(s) <b>if indicated</b></p>
	<p><b>Rapid Transport</b> to appropriate destination using <u>Trauma and Burn: EMS Triage and Destination Plan</u></p>
	<p><b>Notify Destination or Contact Medical Control</b></p>



# Chemical and Electrical Burn

## Pearls

- **Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, and Neuro**
- Green, Yellow, and Red in burn severity do not apply to the Start/ JumpStart Triage System.
- Refer to Rule of Nines.
- **Transport and Destination:**
  - In general, chemical and electrical burns should be transported to a burn center.
  - Burn center should be initial destination choice unless EMS system access is limited by time and/ or distance.
  - When EMS transport to burn center is limited, transport to and stabilization at local center is appropriate.
- **Chemical Burns:**
  - Refer to Decontamination Procedure.
  - With dry powders/ substances, gently brush or wipe off prior to irrigation. Do not aerosolize by brushing too vigorously.
  - Normal Saline or Sterile Water is preferred, however if not available, do not delay irrigation and use tap water. Other water sources may be used based on availability.
  - Flush the area as soon as possible with the cleanest, most readily available water or saline solution and use copious amounts of fluids.
  - Flush contact area for a minimum of 15 minutes and continue until arrival at receiving facility.
  - Hydrofluoric acid burns:**
    - Monitor ECG for peaked T waves, which can be sign of hypocalcemia.
  - Eye involvement:**
    - Irrigation is recommended for a minimum of 30 minutes and continue until arrival at receiving facility.
- **Electrical Burns:**
  - Remember the extent of the obvious external burn from an electrical source does not always reflect more extensive internal damage. Small external injury may have large internal injury.**
  - Do not refer to wounds as an entry and exit wound.**
  - DO NOT contact patient until you are certain the source of the electrical shock is disconnected.**
  - Attempt to locate contact points (generally there will be two or more.) A point where the patient contacted the source and a point(s) where the patient is grounded.
  - Sites will generally be full thickness (3<sup>rd</sup>).
  - Cardiac Monitor: Anticipate ventricular or atrial irregularity including VT, VF, atrial fibrillation, and/ or heart blocks.
  - Attempt to identify the nature of the electrical source (AC or DC), the amount of voltage, and the amperage the patient may have been exposed to during the electrical shock.
  - Lightning strike:**
    - Lightning strike victims are amenable to airway, breathing, cardiac compressions, as well as early defibrillation.
    - Use concept of reverse triage with multiple casualties. Resuscitate lightning strikes as the priority.**
    - Lightning strike victims found alive do not often deteriorate quickly.



# Crush Syndrome Trauma

## History

- Entrapped and crushed under heavy load > 30 minutes
- Extremity / body crushed
- Building collapse, trench collapse, industrial accident, pinned under heavy equipment

## Signs and Symptoms

- Hypotension
- Hypothermia
- Abnormal ECG findings
- Pain
- Anxiety

## Differential

- Entrapment without crush syndrome
- Vascular injury with perfusion deficit
- Compartment syndrome
- Altered mental status

	Age Appropriate Airway Protocol(s) AR 1 - 7 <i>as indicated</i>
<b>B</b>	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
<b>P</b>	Cardiac Monitor
	Multiple Trauma Protocol TB 6 <i>if indicated</i>
	Thermal Burn Protocol TB 9 Chemical and Electrical Burn Protocol TB 2 <i>if indicated</i>
	Pain Control Protocol UP 11 <i>as indicated</i>


## Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60  
Ages  $\geq$  1 month: SBP < 70  
Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90  
Ages  $\geq$  65: SBP < 100

All ages:  
Shock Index HR > SBP

<b>P</b>	Consider Midazolam 0.5 – 2 mg IV / IO Midazolam 1 – 2 mg IN Maximum 5 mg Pediatric: 0.1 – 0.2 mg / kg IV / IO / IN Pediatric Maximum 2 mg Over 2 – 3 minutes as needed
<b>A</b>	Normal Saline Infusion 1 Liter per hour IV / IO Pediatric: 3 x maintenance fluid rate
<b>A</b>	Decrease Normal Saline Infusion 500 mL per hour IV / IO Pediatric: Maintenance fluid rate
<b>P</b>	Calcium Gluconate 2 g IV / IO Or (Calcium Chloride 1 g IV / IO) Pediatric: 20 mg / kg IV / IO Over 2- 3 minutes  Albuterol Nebulizer 2.5 – 5 mg May repeat x 3  Sodium Bicarbonate 50 mEq IV / IO Pediatric: 1 mEq / kg IV / IO



**Rapid Transport** to appropriate destination using  
**Trauma and Burn:**  
**EMS Triage and Destination Plan**

Notify Destination or Contact Medical Control

Exit to  
Age Appropriate  
Cardiac Arrest  
Protocol AC 3 / PC 4  
Arrhythmia Protocol(s)  
*if indicated*





# Crush Syndrome Trauma

## Pearls

- **Recommended exam: Mental Status, Musculoskeletal, Neuro**
- **Scene safety is of paramount importance as typical scenes may pose hazards to rescuers. Call for appropriate resources.**
- **Crush Injury is a localized crush injury with systemic signs and symptoms causing muscle breakdown and release of potentially toxic muscle cell components and electrolytes into the circulation.**
- **Crush syndrome typically manifests after 1 – 4 hours of crush injury.**
- **Fluid resuscitation strategy:**
  - If possible, administer IV / IO fluids prior to release of crushed body part, especially with crush > 1 hour.
  - If access to patient and initiation of IV / IO fluids occurs after 2 hours, give 2 liters of IV fluids in adults and 20 mL/kg of IV fluids in pediatrics, and then begin > 2 hour dosing regimen.
  - **If not able to perform IV / IO fluid resuscitation immediately, place tourniquet on crushed limb until IV / IO fluids can be initiated (even if tourniquet is not being used for hemorrhage control).**
- **Pediatric IV Fluid maintenance rate:**
  - 4 mL for the first 10 kg of weight +
  - 2 mL for the second 10 kg of weight +
  - 1 mL for every additional kg in weight after 20 kg
- **Example: 28 kg pediatric**

First 10 kg:	4 mL/kg/hr = 40 mL/hr
Second 10 kg:	2 mL/kg/hr = 20 mL/hr
Final 8 Kg:	1 mL/kg/hr = 8 mL/hr
Total: 68 mL/hr rate	
- Consider all possible causes of shock and treat per appropriate protocol.
- Majority of decompensation in pediatrics is airway or respiratory related.
- Decreasing heart rate and hypotension occur late in children and are signs of impending cardiac arrest.
- Shock may be present with a normal blood pressure initially or even elevated.
- Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the only sign.
- **Patients may become hypothermic even in warm environments. Maintain warmth.**
- Hyperkalemia from crush syndrome can produce ECG changes described in protocol, but may also be a bizarre, wide complex rhythm. Wide complex rhythms should also be treated using the VF/ Pulseless VT Protocol if indicated (AC 9 VF Pulseless VT Protocol and/ or PC 7 Pediatric VF Pulseless VT Protocol).

## TB 3



# Extremity Trauma

## History


- Type of injury
- Mechanism: crush/ penetrating/ amputation
- Time of injury
- Open vs. closed wound / fracture
- Wound contamination
- Medical history
- Medications

## Signs and Symptoms

- Pain and/ or swelling
- Deformity
- Altered sensation/ motor function
- Diminished pulse/ capillary refill
- Decreased extremity temperature

## Differential

- Abrasion
- Contusion
- Laceration
- Sprain
- Dislocation
- Fracture
- Amputation



## Trauma and Burn Protocol Section

## Pearls

- **Recommended Exam: Mental Status, Extremity, Neuro, Perfusion**
- Peripheral neurovascular status is important to assess and document, as well as time of assessment.
- In amputations, time is critical. Transport and notify medical control immediately, so that the appropriate destination can be determined.
- Hip dislocations as well as knee and elbow fracture/ dislocations have a high incidence of vascular compromise.
- Urgently transport any injury with neurological or vascular compromise.
- Blood loss may be concealed or not apparent with extremity injuries.
- Lacerations optimally should be evaluated for repair within 6 hours from the time of injury.
- **Multiple casualty incident: Tourniquet Procedure may be considered first instead of direct pressure.**

**TB 4**



# Head Trauma

## History

- Time of injury
- Mechanism (blunt vs. penetrating)
- Loss of consciousness
- Bleeding
- Past medical history
- Medications
- Evidence for multi-trauma

## Signs and Symptoms

- Pain, swelling, bleeding
- Altered mental status
- Unconscious
- Respiratory distress/ failure
- Vomiting
- Major traumatic mechanism of injury
- Seizure

## Differential

- Skull fracture
- Brain injury (Concussion, Contusion, Hemorrhage)
- Epidural hematoma
- Subdural hematoma
- Subarachnoid hemorrhage
- Spinal injury
- Abuse

## Prevent hypoxia, hypotension, and hyperventilation

**A single episode of hypoxia, hypotension, and hyperventilation increases mortality**

<b>Age Appropriate</b>  Airway Protocol(s) AR 1, 2, 3, 5, 6 <i>if indicated</i>	
	<b>Obtain and Record GCS</b>
	All patients
	Titrate <b>target SpO2 100%</b>
	Monitor HR, BP and O2 every 3-5 minutes
	Blood Glucose Analysis Procedure
<b>B</b>	Maintain EtCO2 35 – 45 mmHg
<b>A</b>	IV or IO Access - UP 6 <i>if indicated</i>
<b>P</b>	Cardiac Monitor
	Altered Mental Status - UP 4 <i>if indicated</i>
	Multiple Trauma - TB 6 <i>if indicated</i>
	<b>Age Appropriate</b>  Hypotension/ Shock - AM 5/ PM 3 <i>if indicated</i>
	Seizure - UP 13 <i>if indicated</i>
	Spinal Motion Restriction Protocol TB 8 Procedure WTP 2 <i>if indicated</i>
	Pain Control - UP 11 <i>if indicated</i>
	Monitor and Reassess

**Rapid Transport** to appropriate destination using  
**Trauma and Burn:**  
**EMS Triage and Destination Plan**

**Hyperventilation:**  
Hyperventilation is NOT recommended in patients who require BVM, BIAD, or ETT.

Maintain ventilation rate to target EtCO2 of 35 – 45 mmHg  
See Pearls

## Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60  
Ages  $\geq$  1 month: SBP < 70  
Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90  
Ages  $\geq$  65: SBP < 110

All ages Shock Index:  
 $SI = HR \div SBP$

Use Shock Index, Pediatric Adjusted (SIPA) for children  $<12$  (see pearls)

**Notify Destination or Contact Medical Control**



# Head Trauma

Eye Opening Response	Verbal Response	Motor Response
4 = Spontaneous	5 = Oriented	6 = Obeys commands
3 = To verbal stimuli	4 = Confused	5 = Localizes pain
2 = To pain	3 = Inappropriate words	4 = Withdraws from pain
1 = None	2 = Incoherent	3 = Flexion to pain or decorticate
	1 = None	2 = Extension to pain or decerebrate
		1 = None

Age	HR	SBP	SIPA cutoff value
1-3 years	70-110	90-110	1.2
4-6 years	65-110	90-110	1.2
7-12 years	60-100	100-120	1.0
>12 years	55-90	100-135	0.9

SIPA, shock index, pediatric age-adjusted; HR, heart rate; SBP, systolic blood pressure.

## Pearls

- Recommended Exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Back, Neuro
- Hypoxia:
  - Single episode of hypoxia can worsen head injury and double mortality.
  - Titrate SpO<sub>2</sub> as close to 100% as possible.
- Hyperventilation in head injury requiring advanced airway:
  - Hyperventilation lowers CO<sub>2</sub> and causes vasoconstriction leading to increased intracranial pressure (ICP).
  - Hyperventilation is not recommended and can worsen the brain injury.
  - In patients requiring BVM, BIAD, or endotracheal tube, titrate ventilation rate to EtCO<sub>2</sub> between 35 - 45 mmHg.
  - Recommended ventilation rates with advanced airways:
    - Infant/ Toddler: 25 breaths / minute
    - Children: 20 Breaths / minute
    - Adolescents/ Adults: 10 – 12 Breaths / minute
- Hypotension:
  - Episodes of hypotension can worsen head injury and increase mortality:
  - In adults, minimal SBP is at least 90 - 100 mmHg.
  - In pediatrics, minimal SBP is at least > 70 + (2 x the age in years).
  - Usually indicates shock unrelated to the head injury and should be aggressively treated, otherwise limit fluid administration.
- GCS
  - Key performance measure used in the EMS Acute Trauma Care Toolkit.
  - Serial assessments of GCS with ongoing assessments should be performed.
- Do not place in Trendelenburg position as this may increase ICP and worsen blood pressure.
- Poorly fitted cervical collars may also increase ICP when applied too tightly.
- In areas with short transport times, Drug Assisted Airway protocol is not recommended for patients who are spontaneously breathing and who have oxygen saturations of  $\geq 90\%$  with supplemental oxygen including BIAD/ BVM.
- Increased intracranial pressure (ICP) may cause hypertension and bradycardia (Cushing's Response).
- Consider Restraints if necessary for patient's and/ or personnel's protection per the Restraints: Physical Procedure USP 5.
- Concussions:
  - Traumatic brain injuries involving any of a number of symptoms including confusion, loss of consciousness, vomiting, or headache.
  - Any prolonged confusion or mental status abnormality which does not return to normal within 15 minutes or any documented loss of consciousness should be evaluated by a physician ASAP.
  - EMS Providers should not make return-to-play decisions when evaluating an athlete with suspected concussion.**
  - This is outside the scope of practice.



# Multiple Trauma

## History

- Time and mechanism of injury
- Damage to structure or vehicle
- Location in structure or vehicle
- Others injured or dead
- Speed and details of MVC
- Restraints/ protective equipment
- Past medical history
- Medications

## Signs and Symptoms

- Pain, swelling
- Deformity, lesions, bleeding
- Altered mental status or unconscious
- Hypotension or shock
- Arrest

## Differential (Life threatening)

- Uncontrolled hemorrhage
- Airway obstruction/ deformity
- Chest:
  - Tension pneumothorax
  - Flail chest/ Open chest wound
  - Pericardial tamponade/ Hemothorax
- Head Trauma Protocol TB 5
- Intra-abdominal bleeding
- Pelvis/ Femur/ Extremity fracture
- Spine fracture/ Cord injury
- Hypothermia

	Age Appropriate Airway Protocol(s) AR 1 - 7 <i>as indicated</i>
	Chest Decompression Procedure WTP 1 <i>if indicated</i>
	Control External Hemorrhage Procedure(s) WTP 4, 5, 7 Consider Pelvic Binding Splint Fractures Procedure WTP 3
	IV or IO Access Protocol UP 6
	Spinal Motion Restriction Procedure WTP 2 Spinal Motion Restriction Protocol TB 8 <i>if indicated</i>
	<b>Obtain and Record GCS</b>

## TXA/ Blood Product Indicators:

V/S parameters  
for blunt or penetrating trauma:

### Adult:

- SBP  $\leq$  70 mmHg  
or
- SBP  $\leq$  90 mmHg + HR  $\geq$  110
- Age  $\geq$  65  
SBP  $<$  100 mmHg + HR  $>$  100

### Peds:

- SBP  $<$  {70 + 2(Age)}

VS / Perfusion Abnormal / Shock? YES

NO

	Head Injury Protocol TB 5 <i>if indicated</i>
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Pain Control Protocol UP 11 <i>if indicated</i>
	Extremity Trauma Protocol TB 4 <i>if indicated</i>
	Crush Syndrome Protocol TB 3 <i>if indicated</i>
	Repeat Assessment Adult Procedure
	Monitor and Reassess

	Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 <i>if indicated</i>
	TXA 1 - 2 g over 10 minutes IV / IO Peds: (Per local trauma guidelines) over 10 minutes IV / IO Maximum 1 gm <i>if indicated</i>

P

Blood Products

*if indicated*

Rapid Transport to appropriate destination using <b>Trauma and Burn:</b> <b>EMS Triage and Destination Plan</b> Limit Scene Time $\leq$ 15 minutes Provide Early Notification		
	Notify Destination or Contact Medical Control	

**TB 6**



# Multiple Trauma

## Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Back, Neuro
- **Items in Red Text** are key performance measures used in the EMS Acute Trauma Care Toolkit
- Scene time should not be delayed for procedures and all should be performed during rapid transport of unstable patients.
- Ask all patients if they are taking any anticoagulants and report during facility transition of care.
- **Airway:**
  - BVM and BIAD are acceptable for airway management to maintain SpO<sub>2</sub> of 92 – 98%.
  - Endotracheal intubation, if performed, should be completed during transport and should not delay scene time.
- **Breathing:**
  - Consider Chest Decompression with signs of shock and/ or injury to torso with evidence of tension pneumothorax.
- **Circulation:**
  - Control external hemorrhage and prevent hypothermia by keeping patient warm.
  - IV or IO access should be established during rapid transport of unstable patients.
- **Head Injury with multiple trauma (Refer to Head Trauma Protocol TB 5):**
  - Higher SBP targets are needed to maintain cerebral perfusion pressure.
  - Single episodes of Hypotension and/ or hypoxia are associated with worse outcomes in head injured patients.
  - Adult SBP target is  $\geq 100$  mmHg.
  - Pediatric SBP target is  $\geq 70 + 2(\text{Age})$  mmHg.
- **Trauma Triad of Death:**
  - Metabolic acidosis/ Coagulopathy/ Hypothermia
- **Address by appropriate resuscitation measures and keeping patient warm, regardless of ambient temperature, which helps to treat metabolic acidosis, coagulopathy, and hypothermia.**
- **Tranexamic Acid (TXA):**
  - Agencies utilizing TXA must submit letters from their receiving trauma centers for approval by the OEMS Medical Director.
  - Receiving trauma centers must agree to continue TXA therapy with repeat dosing.
  - TXA is NOT indicated and should NOT be administered where trauma occurred  $> 3$  hours prior to EMS arrival.
- **Trauma in Pregnancy:**
  - Providing optimal care for the mother = optimal care for the fetus.
  - After 20 weeks gestation (fundus at or above umbilicus) transport patient on left side with 10 – 20° of elevation.
- **Geriatric Trauma:**
  - Age  $\geq 65$ : SBP  $< 110$  mmHg or HR  $>$  SBP may indicate shock.
  - Evaluate with a high index of suspicion, occult injuries difficult to recognize and with unexpected patient decompensation.
  - Risk of death with trauma increases after age 55.
  - Low impact mechanisms, such as ground level falls might result in severe injury especially in age over 65.
- See Regional Trauma Guidelines when declaring Trauma Activation.
- Maintain high-index of suspicion for domestic violence or abuse, pediatric non-accidental trauma, or geriatric abuse.
- **Refer to your Regional Trauma Guidelines when declaring Trauma Activation.**
- Severe bleeding from an extremity, not rapidly controlled with direct pressure, needs application of a tourniquet.
- Maintain high-index of suspicion for domestic violence or abuse, pediatric non-accidental trauma, or geriatric abuse.



# Radiation Incident

## History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history/ Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status


## Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/ distress could be indicated by hoarseness/ wheezing
- Hypotension
- Thermal or Chemical Injury

## Differential

- Thermal / Chemical / Electrical Burn Injury
- Superficial  
(1<sup>st</sup> Degree) red – painful  
(Don't include in TBSA)
- Partial Thickness  
(2<sup>nd</sup> Degree) blistering
- Full Thickness  
(3<sup>rd</sup> Degree) painless/charred or leathery skin

**Scene Safety / Quantify number and Triage Patients/ Load and Go with Assessment/ Treatment Enroute**




**Collateral Injury:** Most all injuries immediately seen will be a result of collateral injury, such as heat from the blast, trauma from concussion, treat collateral injury based on typical care for the type of injury displayed.

**Qualify:** Determine exposure type; external irradiation, external contamination with radioactive material, internal contamination with radioactive material.

**Quantify:** Determine exposure (generally measured in Grays/Gy). Information may be available from those on site who have monitoring equipment, do not delay transport to acquire this information.



# Radiation Incident



Time Phases of Radiation Injury  
(Exposure Dose vs Clinical Outcome)

Exposure Dose (Gy)	Prodrome Severity	Manifest Illness - Symptom Severity			Prognosis
		Hematologic	Gastrointestinal	Neurologic	
0.5 to 1.0	+	+	0	0	Survival almost certain
1.0 to 2.0	+/++	+	0	0	Survival >90 percent
2.0 to 3.5	++	++	0	0	Probable survival
3.5 to 5.5	+++	+++	+	0	Death in 50% at 3.5 to 6 wks
5.5 to 7.5	+++	+++	++	0	Death probable in 2-3 wks
7.5 to 10	+++	+++	+++	0*	Death probable in 1-2.5 wks
10 to 20	+++	+++	+++	+++	Death certain in 5-12 days
> 20	+++	+++	+++	+++**	Death certain in 2-5 days

**Abbreviations:** Gy: dose in Grey;  
0: no effects; +: mild; ++: moderate; +++: severe or marked

\* Hypotension

\*\* Also cardiovascular collapse, fever, shock


Modified from : Waselenko, JK, MacVittie, TJ, Blakely, WF, et al. Medical management of the acute radiation syndrome: Recommendations of the strategic national stockpile radiation working group. Ann Int Med 2004; 140:1039.

## Pearls

- The three primary methods of protection from radiation sources:**
  - Limiting time of exposure
  - Distance from
  - Shielding from the source
- Dealing with a patient with a radiation exposure can be a frightening experience. Do not ignore the ABC's, a dead but decontaminated patient is not a good outcome. Refer to the Decontamination Procedure USP 2 for more information.
- Normal Saline or Sterile Water is preferred, however if not available, do not delay irrigation using tap water. Other water sources may be used based on availability. Flush the area as soon as possible with the cleanest, most readily available water or saline solution using copious amounts of fluids.
- Three methods of exposure:**
  - External irradiation
  - External contamination
  - Internal contamination
- Two classes of radiation:**
  - Ionizing radiation (greater energy) is the most dangerous and is generally in one of three states:  
Alpha Particles, Beta Particles and Gamma Rays.
  - Non-ionizing (lower energy) examples include microwaves, radios, lasers and visible light.
- Radiation burns with early presentation are unlikely, it is more likely this is a combination event with either thermal or chemical burn being presented as well as a radiation exposure. When the burn is from a radiation source, it indicates the patient has been exposed to a significant source, (> 250 rem).
- Patients experiencing radiation poisoning are not contagious. Cross contamination is only a threat with external and internal contamination.
- Typical ionizing radiation sources in the civilian setting include soil density probes used with roadway builders and medical uses such as x-ray sources as well as radiation therapy. Sources used in the production of nuclear energy and spent fuel are rarely exposure threats as is military sources used in weaponry. Nevertheless, these sources are generally highly radioactive and in the unlikely event they are the source, consequences could be significant and the patient's outcome could be grave.
- Dirty bomb ingredients generally include previously used radioactive material and are usually combined with a conventional explosive device to spread and distribute the contaminated material.
- Refer to Decontamination Procedure USP 2/ WMD and Nerve Agent Protocol TE 8 for contamination events.
- If there is a time lag between the time of exposure and the encounter with EMS, key clinical symptom evaluation includes: nausea/ vomiting, hypothermia/ hyperthermia, diarrhea, neurological/ cognitive deficits, headache, and hypotension.
- This event may require an activation of the National Radiation Injury Treatment Network (RITN). UNC Hospitals, Atrium Health Wake Forest Baptist and Duke are the RITN hospitals, with burns managed at UNC and Wake Forest.



# Selective Spinal Motion Restriction



## Pearls

- Recommended Exam: Mental Status, Skin, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro
- Patients meeting all the above criteria do not require spinal motion restriction. However, patients who fail one or more criteria above require spinal motion restriction, but does NOT require use of the long spine board for immobilization.
- Long spine boards are NOT considered standard of care in most cases of potential spinal injury. Spinal motion restriction with cervical collar, and securing patient to cot, while padding all void areas is appropriate.
- True spinal immobilization is not possible. Spine protection and spinal motion restriction do not equal long spine board.
- Spinal motion restriction is always utilized in at-risk patients. This includes cervical collar, securing to stretcher, minimizing movement/ transfers, and maintenance of in-line spine stabilization during any necessary movement/ transfers. This includes the elderly, or others with body or spine habitus preventing them from lying flat.
- Consider spinal motion restriction in patients with arthritis, cancer, dialysis, and underlying spine or bone disease.
- Range of motion (ROM) is tested by touching chin to chest (look down), extending neck (look up), and turning head from side to side (chin to each shoulder) only in patients without posterior cervical mid-line pain. ROM should NOT be assessed if patient has midline spinal tenderness. Patient's range of motion should not be assisted, they must be able to complete alone.
- EMR may participate in spinal motion restriction per Agency Medical Director.
- Immobilization on a long spine board is not necessary where:
  - Penetrating trauma to the head, neck or torso with no signs and/ or symptoms of spinal injury.
- Concerning mechanisms that may result in spinal column injury:
  - Fall from  $\geq 3$  feet and/ or  $\geq 5$  stairs or steps. Ground level falls in patients  $\geq 65$  years of age.
  - MVC  $\geq 30$  mph, rollover, and/or ejection
  - Motorcycle, bicycle, other mobile device, or pedestrian-vehicle crash
  - Diving or axial load to spine
  - Electric shock



# Thermal Burn

## History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history/ Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status


## Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/ distress could be indicated by hoarseness/ wheezing

## Differential

- Thermal / Chemical / Electrical Burn Injury
- Superficial (1<sup>st</sup> Degree) red – painful (Don't include in TBSA)
- Partial Thickness (2<sup>nd</sup> Degree) blistering
- Full Thickness (3<sup>rd</sup> Degree) painless/charred or leathery skin
- Radiation injury
- Blast injury

## Assess Burn/ Concomitant Injury Severity



## Airway Protocol(s) AR 1, 2, 3, 5, 6 *as indicated*

### Multiple Trauma Protocol TB 6 *if indicated*

Remove Rings, Bracelets, Constricting Items

Dry Clean Sheet or Dressings

### IV or IO Access Protocol UP 6 Consider 2 IV sites if $\geq 15\% \text{ TBSA}$

Lactated Ringers  
or  
Normal Saline

$0.25 \text{ mL/kg} \times (\text{TBSA}\%) \text{ /hr}$

A

Lactated Ringers  
or  
Normal Saline

Age:  $\leq 5 = 125 \text{ mL/hr}$   
Age: 6–14 = 250 mL/hr  
Age:  $\geq 15 = 500 \text{ mL/hr}$

A

NO

YES

TBSA estimate at  $\geq 20\%$

## Pain Control Protocol UP 11 *if indicated*

## Carbon Monoxide/ Cyanide Protocol TE 2 *if indicated*

Monitor and Reassess

**Rapid Transport** to appropriate destination using  
**Trauma and Burn:  
EMS Triage and Destination Plan**

**Formula example:**  
70 kg patient with 15% TBSA


$$0.25 \times 70 \text{ kg} = 17.5 \\ 17.5 \times 15 = 263 \text{ mL/hr}$$

**Notify Destination or  
Contact Medical Control**

**1. Lactated Ringers preferred over Normal Saline. Use if available, if not change over once available.**



# Thermal Burn



## Rule of Nines

- Rarely find a complete isolated body part that is injured as described in the Rule of Nines.
- More likely, it will be portions of one area, portions of another, and an approximation will be needed.
- For the purpose of determining the extent of serious injury, differentiate the area with minimal or 1<sup>st</sup> degree burn(superficial) from those of partial (2<sup>nd</sup>) or full (3<sup>rd</sup>) thickness burns.
- For the purpose of determining Total Body Surface Area (TBSA) of burn, include only Partial (2<sup>nd</sup>) and Full Thickness (3<sup>rd</sup>) burns.** Report the observation of other superficial (1<sup>st</sup> degree) burns but do not include those burns in your TBSA estimate.
- Some texts will refer to 4<sup>th</sup> 5<sup>th</sup> and 6<sup>th</sup> degree burns. There is significant debate regarding the actual value of identifying a burn injury beyond that of the superficial, partial and full thickness burn at least at the level of emergent and primary care. For our work, all are included in Full Thickness burns


Estimate spotty areas of burn by using the size of the patient's palm as 1 %

## Pearls

- Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, and Neuro**
- Green, Yellow, and Red In burn severity do not apply to the Start/ JumpStart Triage System.
- Airway considerations:**
  - For systems performing RSI, Rocuronium is preferred agent (succinylcholine can be used in the first 24-hours)
  - Singed nasal hairs, facial burns, and/ or carbonaceous sputum are NOT absolute indications for intubation in a burn patient.
  - Utilizing non-rebreather face mask as well as NIPPV procedure are acceptable as tolerated.
- Critical or Serious Burns:**
  - > 5-15% total body surface area (TBSA) 2<sup>nd</sup> or 3<sup>rd</sup> degree burns
  - 3<sup>rd</sup> (full thickness) degree burns > 5% TBSA for any age group
  - Circumferential burns of extremities
  - Electrical or lightning injuries
  - Suspicion of abuse or neglect
  - Inhalation injury
  - Chemical burns
  - Burns of face, hands, perineum, or feet
  - Require direct transport to a Burn Center. Local facility should be utilized only if distance to Burn Center is excessive or critical interventions such as airway management are not available in the field.
- Burn patients are trauma patients, evaluate for multisystem trauma.
- Assure whatever has caused the burn is no longer contacting the injury. (Stop the burning process!)
- Circumferential burns to extremities are dangerous due to potential vascular compromise secondary to soft tissue swelling.
- Burn patients are prone to hypothermia - never apply ice or cool the burn, must maintain normal body temperature.
- Evaluate the possibility of geriatric abuse with burn injuries in the elderly.
- Do not administer IM pain injections to a burn patient. IM dosing is variable in burn patients and may result in over or under dose.



# Traumatic Arrest





# Traumatic Arrest


## Pearls.

- Recommended Exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Back, Neuro
- Items in Red Text are key performance measures used in the EMS Acute Trauma Care Toolkit.
- Scene time should not be delayed for procedures and all should be performed during rapid transport.
- First arriving EMS personnel should make the assessment concerning agonal respirations, pulselessness, asystole or PEA < 40, pupillary reflexes, and spontaneous body movements.
- Withholding resuscitative efforts with blunt and penetrating trauma victims who meet criteria, is appropriate.
- Airway:
  - Airway is a priority in traumatic arrest.
  - BVM and BIAD are acceptable for airway management.
  - Endotracheal intubation, if performed, should be completed during transport and should not delay scene time.
- Breathing:
  - Consider Chest Decompression in both blunt and penetrating trauma.
  - Needle Chest Decompression permissible at the AEMT level only involving TRAUMATIC PULSELESS ARREST.
- Circulation:
  - Control external hemorrhage, including use of tourniquets, and prevent hypothermia by keeping patient warm.
  - IV or IO access should be established during rapid transport of unstable patients.
  - If transport time to Trauma Center is < 15 minutes, use of ECG monitor may delay resuscitation and transport.
  - Rhythm determination is more helpful in rural settings, or where transport to nearest facility is > 15 minutes. Omit from algorithm where appropriate.
  - Organized rhythms, for purpose of protocol, include Ventricular Tachycardia, Ventricular Fibrillation, and PEA.
  - Wide, bizarre rhythms (Idioventricular and severely bradycardic rhythms < 40 BPM), are not organized rhythms.
- Trauma Triad of Death:
  - Metabolic acidosis/ Coagulopathy/ Hypothermia
  - Performance of appropriate resuscitation measures and keeping patient warm, regardless of ambient temperature, helps to treat metabolic acidosis, coagulopathy, and hypothermia.
- Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.
- DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT) compressions to ventilations are 30:2. If advanced airway in place ventilate 10 – 12 breaths per minute.
- ALS procedures should optimally be performed during rapid transport.
- Time considerations:
  - From the time cardiac arrest is identified, if CPR is performed ≥ 15 minutes with no ROSC, consider termination of resuscitation on scene.
  - From the time cardiac arrest is identified, if transport time to closest Trauma Center is > 15 minutes consider termination of resuscitation on scene.
- Lightning strike, drowning, or in situations causing hypothermia, resuscitation should be initiated.
- Where multiple lightning strike victims are found, use Reverse Triage: Begin CPR in apneic/ pulseless victims.
- Agencies utilizing Targeted Temperature Management Protocol should not cool the trauma patient, but rather make every effort to maintain warmth.



# Universal Patient Care

## Universal Protocol Section





# Universal Patient Care

Identify potential hazards to rescuers, patient and public.

Identify number of patients and utilize START protocol if indicated.

Observe patient position and surroundings.

## General:

All patient care must be appropriate to your level of training and documented in the PCR.

The PCR / EMR narrative should be considered a story of the circumstances, events and care of the patient and should allow a reader to understand the complaint, the assessment, the treatment, why procedures were performed and why indicated procedures were not performed as well as ongoing assessments and response to treatment and interventions.

## Adult Patient:

An adult is considered hypotensive when Systolic Blood Pressure is less than 90 mmHg.

Diabetic patients and women may have atypical presentations of cardiac related problems such as MI.

General weakness can be the symptom of a very serious underlying process.

Beta blockers and other cardiac drugs may prevent a reflexive tachycardia in shock with low to normal pulse rates.

## Geriatric Patient:

Hip fractures and dislocations have high mortality.

Altered mental status is not always dementia. Always check Blood Sugar and assess signs of stroke, trauma, etc. with any alteration in a patient's baseline mental status.

Minor or moderate injury in the typical adult may be very serious in the elderly.

Special note on oxygen administration and utilization:

Oxygen is ubiquitous in prehospital patient care and probably over utilized. Oxygen is a pharmaceutical with indications, contraindications as well as untoward side effects. Recent research demonstrates a clear link with increased mortality when given in overdose (hyperoxia / hyperventilation) in cardiac arrest. Utilize oxygen when indicated and not because it is available. A reasonable target oxygen saturation in all treatment protocols is 94 % regardless of delivery device.

## Pearls

- **Recommended Exam: Minimal exam if not noted on the specific protocol is vital signs, mental status with GCS, and location of injury or complaint.**
- Any patient contact, which does not result in an EMS transport, must have a completed Patient Care Report.
- Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.
- Two complete vital sign acquisitions should occur at a minimum with any patient encounter.
- **Patient Refusal (Declining Treatment and/ or Transport):**

Patient refusal is a high risk situation. Encourage patient to accept transport to medical facility.

Encourage patient to allow an assessment, including vital signs. Documentation of the event is very important including a mental status assessment describing the patient's capacity to refuse care.

### Guide to Assessing capacity:

**C: Patient should be able to communicate a clear choice:** This should remain stable over time. Inability to communicate a choice or an inability to express the choice consistently demonstrates incapacity.

**R: Relevant information is understood:** Patient should be able to voice a factual understanding of the illness/ injury, the options, and the risks and benefits of recommended treatment or transport.

**A: Appreciation of the situation:** Ability to communicate an understanding of the facts of the situation. The patient should be able to recognize the significance of the outcome potentially from their decision.

**M: Manipulation of information in a rational manner:** Demonstrate a rational process to come to a decision.

Should be able to describe the logic they are using to come to the decision, though you may not agree with decision.

- **Pediatric Patient General Considerations:**

A pediatric patient is defined by fitting with a Pediatric Medication/ Skill Resuscitation System, Age ≤ 15, weight ≤ 49 kg.

Special needs children may require continued use of Pediatric based protocols regardless of age and weight.

Initial assessment should utilize the **Pediatric Assessment Triangle** which encompasses Appearance, Work of Breathing and Circulation to skin.

The order of assessment may require alteration dependent on the developmental state of the pediatric patient.


Generally the child or infant should not be separated from the caregiver unless absolutely necessary during assessment and treatment.

- Timing of transport should be based on patient's clinical condition and the agency transport policy.
- Consider consultation with Medical Control for patient(s) refusing treatment/ transport.
- Blood Pressure is defined as a Systolic/ Diastolic reading. A palpated Systolic reading may be necessary at times.
- SAMPLE: Signs/ Symptoms; Allergies; Medications; PMH; Last oral intake; Events leading to illness/ injury



# Triage

## Universal Protocol Section





# Triage

## Pearls

- **When approaching a multiple casualty incident where resources are limited:**  
Triage decisions must be made rapidly with less time to gather information  
Emphasis shifts from ensuring the best possible outcome for an individual patient to ensuring the best possible outcome for the greatest number of patients.
- **Scene Size Up:**
  1. Conduct a scene size up. Assure well being of responders. Determine or ensure scene safety before entering. If there are several patients with the same complaints consider HazMat, WMC or CO poisoning.
  2. Take Triage system kit.
  3. Determine number of patients. Communicate the number of patients and nature of the incident and establish incident command.
  4. Direct incoming resources. Identify ingress and egress path. Establish a staging area. Assign a medical officer, triage officer, transportation officer, and staging officer as personal become available.
- **Triage is a continual process and is a continuous process in each section as resources allow.**
- **Step 1: Global sorting:**  
Call out to those involved in the incident to walk to a designated area and assess group last.  
For those who cannot walk, have them wave/ indicate a purposeful movement and assess them second.  
Those involved who are not moving, or have an obvious life threat, assess first.
- **Step 2: Individual assessments:**  
Control major hemorrhage.  
Open airway and if child, give 2 rescue breaths.  
Perform Needle Chest Decompression Procedure if indicated.  
Administer injector antidotes if indicated.
- **Assess the first patient you encounter using the three objective criteria which can be remembered by RPM.**  
**R: Respiratory** (Respiratory rates are difficult to measure quickly, use work of breathing and respiratory distress)  
**P: Perfusion** (Capillary refill can be altered by many factors including skin temperature – use age appropriate heart rates)  
**M: Mental Status** (Motor component of GCS score is important indicator – ability to follow commands)
- If your patient falls into the RED TAG category, stop, place RED TAG and move on to next patient. Attempt only to correct airway problems, treat uncontrolled bleeding, or administer an antidote before moving to next patient.
- **Treatment:**  
Once casualties are triaged, a focus on treatment can begin. You may need to move patients to treatment areas.  
RED TAGs are moved/ treated first, followed by YELLOW TAGs. BLACK TAGs should remain in place.  
You may also indicate deceased patients by pulling their shirt/ clothing over their head.  
As more help arrives, then the triage/ treatment process may proceed simultaneously.
- **Lightning strike (Reverse Triage):**  
Lightning strike victims are amenable to airway, breathing, cardiac compressions as well as early defibrillation.  
Use concept of reverse triage with multiple casualties. Resuscitate lightning strikes as the priority.  
Lightning strike victims found alive do not often deteriorate quickly.
- **SMART triage tag system is utilized in NC.**



# Abdominal Pain Vomiting and Diarrhea

## History

- Age
- Time of last meal
- Last bowel movement/emesis
- Improvement or worsening with food or activity
- Duration of problem
- Other sick contacts
- Past medical history
- Past surgical history
- Medications
- Menstrual history (pregnancy)
- Travel history
- Bloody emesis / diarrhea

## Signs and Symptoms


- Pain
- Character of pain (constant, intermittent, sharp, dull, etc.)
- Distention
- Constipation
- Diarrhea
- Anorexia
- Radiation

### Associated symptoms:

Fever, headache, blurred vision, weakness, malaise, myalgias, cough, headache, dysuria, mental status changes, rash

## Differential

- CNS (increased pressure, headache, stroke, CNS lesions, trauma or hemorrhage, vestibular)
- Myocardial infarction
- Drugs (NSAID's, antibiotics, narcotics, chemotherapy)
- GI or Renal disorders
- Diabetic ketoacidosis
- OB-Gyn disease (ovarian cyst, PID, Pregnancy)
- Infections (pneumonia, influenza)
- Electrolyte abnormalities
- Food or toxin induced
- Medication or Substance abuse
- Psychological



### Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60

Ages  $\geq$  1 month: SBP < 70

Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90

Ages  $\geq$  65: SBP < 110

All ages Shock Index:  
HR > SBP



# Abdominal Pain Vomiting and Diarrhea

## Universal Protocol Section

### Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro
- **Abdominal/ back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and/ or lower extremity pain or diminished pulses, especially in patients over 50 and/ or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 35, diabetics and/ or women, especially with upper abdominal complaints.**
- **Heart Rate:** Tachycardia is one of the first clinical signs of dehydration and volume depletion and typically increases as dehydration becomes more severe.
- **Nausea without vomiting should be treated like vomiting. Patient will benefit from symptom control with antiemetic even if not actively vomiting.**
- **Promethazine (Phenergan):**  
May cause sedative effects in pediatric patients and in ages  $\geq 65$ , and the debilitated, etc.)  
When giving promethazine IV, dilute with 10 mL of normal saline and administer slowly as it can also harm the veins.  
Preferred route is IM.
- Isolated vomiting in children is common but can be a sign of more serious pathology. Pyloric stenosis, bowel obstruction, and CNS processes (bleeding, tumors, or increased CSF pressures) all often present with vomiting.
- Vomiting and diarrhea are common symptoms, but can be the symptoms of uncommon and serious pathology such as stroke, CO poisoning, acute MI, new onset diabetes, diabetic ketoacidosis (DKA), and organophosphate poisoning. Maintain a high index of suspicion for serious pathology.

**UP 3**



# Altered Mental Status

## History

- Known diabetic, medic alert tag
- Drugs, drug paraphernalia
- Report of illicit drug use or toxic ingestion
- Past medical history
- Medications
- History of trauma
- Change in condition
- Changes in feeding or sleep habits

## Signs and Symptoms

- Decreased mental status or lethargy
- Change in baseline mental status
- Bizarre behavior
- Hypoglycemia (cool, diaphoretic skin)
- Hyperglycemia (warm, dry skin; fruity breath; Kussmaul respirations; signs of dehydration)
- Irritability

## Differential

- Head trauma
- CNS (stroke, tumor, seizure, infection)
- Cardiac (MI, CHF)
- Hypothermia
- Infection (CNS and other)
- Thyroid (hyper / hypo)
- Shock (septic, metabolic, traumatic)
- Diabetes (hyper / hypoglycemia)
- Toxicological or Ingestion
- Acidosis / Alkalosis
- Environmental exposure
- Pulmonary (Hypoxia)
- Electrolyte abnormality
- Psychiatric disorder

Age Appropriate


Airway Protocol(s) AR 1, 2, 3, 5, 6  
*if indicated*

Blood Glucose Analysis Procedure

**B** 12 Lead ECG Procedure

IV or IO Access Protocol UP 6

Age Appropriate Diabetic Protocol(s) AM 2/ PM 2  
*if indicated*





# Altered Mental Status

## Pearls

- **Recommended Exam:** Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- **AMS may present as a sign of an environmental toxin or Haz-Mat exposure, protect personal safety.**
- **General:**
  - The patient with AMS poses one of the most significant challenges.
  - A careful assessment of the patient, the scene, and the circumstances should be undertaken.
  - Assume the patient has a life threatening cause of their AMS until proven otherwise.
  - Pay careful attention to the head exam for signs of bruising or other injury.
  - Information found at the scene must be communicated to the receiving facility.
  - Patients not able to communicate with you coherently require a complete secondary survey (head-to-toe) exam to assess for trauma, infection, or signs of maltreatment/ abuse, or neglect.
  - Acute Stroke should be considered in all patients with acute AMS when < 24 hours from onset.
- **Substance misuse:**
  - Patients ingesting substances can pose a great challenge.
  - DO NOT assume recreational drug use and/ or alcohol are the sole reasons for AMS.
  - Misuse of alcohol/ recreational drugs may lead to hypoglycemia or occult trauma.
  - More serious underlying medical and trauma conditions may be the cause.
- **Behavioral health:**
  - The behavioral health patient may present a great challenge in forming a differential.
  - DO NOT assume AMS is the result solely of an underlying psychiatric etiology.
  - Often an underlying medical or trauma condition precipitates a deterioration of a patients underlying disease.
- **Spinal Motion Restriction/ Trauma:**
  - Only utilize spinal immobilization if the situation warrants.
  - The patient with AMS may worsen with increased agitation when immobilized.
- **It is safer to assume hypoglycemia than hyperglycemia if doubt exists. Recheck blood glucose after Dextrose or Glucagon**
- Consider Restraints if necessary for patient's and/ or personnel's protection per USP 5 Restraints: Physical procedure.



# Back Pain

## History


- Age
- Past medical history
- Past surgical history
- Medications
- Onset of pain / injury
- Previous back injury
- Traumatic mechanism
- Location of pain
- Fever
- Improvement or worsening with activity

## Signs and Symptoms

- Pain (paraspinous, spinous process)
- Swelling
- Pain with range of motion
- Extremity weakness
- Extremity numbness
- Shooting pain into an extremity
- Bowel / bladder dysfunction

## Differential

- Muscle spasm / strain
- Herniated disc with nerve compression
- Sciatica
- Spine fracture
- Kidney stone
- Pyelonephritis
- Aneurysm
- Pneumonia
- Spinal Epidural Abscess
- Metastatic Cancer
- AAA





# Back Pain

## Pearls

- **Recommended Exam: Mental Status, Heart, Lungs, Abdomen, Neuro, Lower extremity perfusion, Back**
- Back pain is one of the most common complaints in medicine and affects more than 90% of adults at some point in their life. Back pain is also common in the pediatric population. Most often it is a benign process but in some circumstances can be life or limb threatening.
- Consider pregnancy or ectopic pregnancy with abdominal or back pain in women of childbearing age.
- Consider abdominal aortic aneurysm with abdominal pain especially in patients over 50 and/ or patients with shock/ poor perfusion. Patients may have abdominal pain and/ or lower extremity pain with diminished pulses. Notify receiving facility early with suspected abdominal aneurysm.
- Consider cardiac etiology in patients > 35, diabetics and/ or women especially with upper abdominal complaints.
- **Red Flags which may signal a more serious process associated with back pain:**
  - Age > 50 or < 18
  - Neurological deficit (leg weakness, urinary retention, or bowel incontinence)
  - IV Drug use
  - Fever
  - History of cancer, either current or remote
  - Night time pain in pediatric patients
- **Cauda equina syndrome is where the terminal nerves of spinal cord are being compressed (Symptoms include):**
  - Saddle anesthesia (numbness between the genitalia and rectum)
  - Recent onset of bladder and bowel dysfunction. (Urine retention and bowel incontinence)
  - Severe or progressive neurological deficit in the lower extremity.
  - Motor weakness of thigh muscles or foot drop
- **Back pain associated with infection:**
  - Fever/ chills.
  - IV Drug user (consider spinal infection)
  - Recent bacterial infection like pneumonia.
  - Immune suppression such as HIV or patients on chronic steroids like prednisone.
  - Meningitis.
- **Spinal motion restriction in patients with underlying spinal deformity should be maintained in their functional position.**
- Kidney stones typically present with an acute onset of flank pain which radiates around to the groin area.



# IV or IO Access

## Universal Protocol Section

### History


- Chronic medical conditions requiring recurrent need for IV access for medication, hydration, or blood sampling.
- Medical condition requiring administration of IV medications at home.
- End-stage renal disease requiring hemodialysis.
- Chronic medical condition requiring IV nutrition.

### Signs and Symptoms

- Fever
- Bleeding
- Hypotension
- Redness, swelling, and/or pain at IV catheter site
- Shortness of breath
- Chest pain
- IV catheter patency

### Differential

- Infection or sepsis
- Infection of catheter
- Clotted IV catheter
- Air embolism
- Pneumothorax
- Overdose of home medication
- Shock



### Pearls

- Frequent encounter of patients with IV access devices and confusion as to which device can be accessed and used by EMS providers are common.
- If unclear about device use, always ask "Is this device used for dialysis?"
- When accessing central catheter, always ensure sterility of catheter connection point by cleaning port with alcohol, or similar disinfectant, 2 – 3 times prior to access.
- Central line catheters placed for administration of chemotherapy, medications, electrolytes, antibiotics, and blood are available to EMS providers for access and administration of fluids, medications, antibiotics, and blood products.
- Central line catheters placed for hemodialysis are NOT available for access by EMS providers unless the patient is in cardiac arrest.
- Long term IV access is frequently needed for a variety of indications:
  - Medication administration such as antibiotics, pain relief, or chemotherapy.
  - Administration of IV nutrition or feeding.
  - Need for multiple IV line access or recurrent blood sampling.
  - Poor vasculature requiring repeated attempts at IV access.
  - End-stage renal disease requiring hemodialysis.
- Common complications of central access devices:
 

Infection	Loss of patency due to clogging or clotting
Damage to vasculature	Pneumothorax
Air embolism	



# IV or IO Access

## Types of IV catheters:

### Port-a-Cath® :


Surgically implanted device allowing easy access to venous system. The port and the catheter are all placed beneath the skin.

Requires a special kit and a specific needle to access.

**Paramedic does NOT routinely access this device.**

**Paramedic may utilize if already accessed with needle/ extension.**

**Paramedic may access if trained on procedure with access to proper equipment.**



### Dialysis Catheter:

Surgically implanted device used to access the vasculature for hemodialysis.

May be tunneled under the skin with access on outside of skin surface or may be non-tunneled with greater portion of catheter on outside of skin surface.

Catheter has a RED port indicating use for dialysis:


Most catheters have a RED port and a BLUE port.

Some catheters have a RED port and a WHITE port.

Dialysis catheters may be used for both short and long-term dialysis and should not be accessed or used for delivery of fluids, medications, antibiotics, or blood products as it increases risk of infection, which then requires removal and subsequent loss of dialysis access.

**Paramedic and AEMT do NOT routinely access this device.**

**Paramedic and AEMT MAY access during cardiac arrest only (Only if IV or IO access cannot be established.)**



### PICC (Peripherally Inserted Central Catheters):


Long catheter inserted into a vein in arm or leg (less common) with the tip of the catheter positioned into the central circulation.

Used for long-term IV fluids, medication administration, blood administration or blood draws.

May have 1 or 2 ports (possibly more, but less common.)

Port ends usually white, blue, or purple. (May be red, less common and is not used for dialysis.)

**Paramedic and AEMT may access and utilize following clean technique.**



### Central Lines:


Catheter placed in large vein in the neck, under the clavicle, or in the groin.

Used for long-term IV fluids, medication administration, blood administration or blood draws.

May have 1 - 4 ports (possibly more, but less common.)

Port ends usually white, blue, or purple. (May be red, less common and is not used for dialysis.)

**Paramedic and AEMT may access and utilize following sterile technique.**





# Dental Problems

## History

- Age
- Past medical history
- Medications
- Onset of pain / injury
- Trauma with "knocked out" tooth
- Location of tooth
- Whole vs. partial tooth injury


## Signs and Symptoms

- Bleeding
- Pain
- Fever
- Swelling
- Tooth missing or fractured

## Differential

- Decay
- Infection
- Fracture
- Avulsion
- Abscess
- Facial cellulitis
- Impacted tooth (wisdom)
- TMJ syndrome
- Myocardial infarction

## Universal Protocol Section



## Pearls

- **Recommended Exam: Mental Status, HEENT, Neck, Chest, Lungs, Neuro**
- Significant soft tissue swelling to the face or oral cavity can represent a cellulitis or abscess.
- Scene and transport times should be minimized in complete tooth avulsions. Reimplantation is possible within 4 hours if the tooth is properly cared for, but unlikely when > 1 hour from time of injury.
- Cardiac chest pain may radiate to the jaw and teeth mimicking dental pain.
- **Avulsed tooth:**  
Handle tooth by the crown, do not touch the root.  
Rinse tooth if soiled but do not scrub, as this can damage the ligaments vital for possible reimplantation.  
Rinse with mild, commercial tooth solution, normal saline or lactated ringers, or the patient's own saliva if dry.  
Transport tooth in milk, commercial solution, patient's own saliva, or IV solution in a container to protect.
- **TXA Use in Dental Bleeding:**  
May be used topically if approved by agency Medical Director.  
Procedure should be created with specific guidance on how to administer TXA for dental bleeding.  
TXA offers modest benefit as TXA instilled gauze combined with direct pressure.



# Emergencies Involving Indwelling Central Lines

## History


- Central Venous Catheter Type
  - Tunneled Catheter (Broviac / Hickman)
- PICC (peripherally inserted central catheter)
- Implanted catheter (Mediport / Hickman)
- Occlusion of line
- Complete or partial dislodge
- Complete or partial disruption

## Signs and Symptoms

- External catheter dislodgement
- Complete catheter dislodgement
- Damaged catheter
- Bleeding at catheter site
- Internal bleeding
- Blood clot
- Air embolus
- Erythema, warmth or drainage about catheter site indicating infection

## Differential

- Fever
- Hemorrhage
- Reactions from home nutrient or medication
- Respiratory distress
- Shock



## Pearls

- Always talk to family / caregivers as they have specific knowledge and skills.
- Use strict sterile technique when accessing / manipulating an indwelling catheter.
- Cardiac arrest: May access central catheter and utilize if functioning properly.
- Do not attempt to force catheter open if occlusion evident.
- Some infusions may be detrimental to stop. Ask family or caregiver if it is appropriate to stop or change infusion.
- Hyperalimentation infusions (IV nutrition): If stopped for any reason monitor for hypoglycemia.



# Epistaxis

## History


- Age
- Past medical history
- Medications (HTN, anticoagulants, aspirin, NSAIDs)
- Previous episodes of epistaxis
- Trauma
- Duration of bleeding
- Quantity of bleeding

## Signs and Symptoms

- Bleeding from nasal passage
- Pain
- Nausea
- Vomiting

## Differential

- Trauma
- Infection (viral URI or Sinusitis)
- Allergic rhinitis
- Lesions (polyps, ulcers)
- Hypertension



## Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60  
 Ages  $\geq$  1 month: SBP < 70  
 Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90  
 Ages  $\geq$  65: SBP < 110

All ages Shock Index: HR > SBP

## Universal Protocol Section

## Pearls

- **Recommended Exam:** Mental Status, HEENT, Heart, Lungs, Neuro
- **TXA Use in Epistaxis:**
  - May be used topically if approved by agency Medical Director.
  - Procedure should be created with specific guidance on how to administer TXA for epistaxis.
  - No clear evidence that TXA provides benefit over conventional vasoconstrictors and sustained direct pressure.
- It is very difficult to quantify the amount of blood loss with epistaxis.
- Bleeding may also be occurring posteriorly. Evaluate for posterior blood loss by examining the posterior pharynx.
- Anticoagulants include warfarin (Coumadin), Apixaban (Eliquis), heparin, enoxaparin (Lovenox), dabigatran (Pradaxa), rivaroxaban (Xarelto), and many over the counter headache relief powders.
- Anti-platelet agents like aspirin, clopidogrel (Plavix), aspirin/ dipyridamole (Aggrenox), and ticlopidine (Ticlid) can contribute to bleeding.



# Fever/ Infection Control

## History

- Age
- Duration of fever
- Severity of fever
- Past medical history
- Medications
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Environmental exposure
- Last acetaminophen or ibuprofen

## Signs and Symptoms


- Warm
- Flushed
- Sweaty
- Chills/Rigors

## Associated Symptoms (Helpful to localize source)

- Myalgias, cough, chest pain, headache, dysuria, abdominal pain, mental status changes, rash

## Differential

- Infections / Sepsis
- Cancer / Tumors / Lymphomas
- Medication or drug reaction
- Connective tissue disease
- Arthritis
- Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis



## Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro
- Febrile seizures are more likely in children with a history of febrile seizures and with a rapid elevation in temperature.
- Patients with a history of liver failure should not receive acetaminophen.
- **Droplet precautions** include standard PPE plus a standard surgical mask for providers who accompany patients in the back of the ambulance and a surgical mask or NRB O2 mask for the patient. This level of precaution should be utilized when influenza, meningitis, mumps, streptococcal pharyngitis, and other illnesses spread via large particle droplets are suspected. A patient with a potentially infectious rash should be treated with droplet precautions.
- **Airborne precautions** include standard PPE plus utilization of a gown, change of gloves after every patient contact, and strict hand washing precautions. This level of precaution is utilized when multi-drug resistant organisms (e.g. MRSA), scabies, or zoster (shingles), or other illnesses spread by contact are suspected.
- **All-hazards precautions** include standard PPE plus airborne precautions plus contact precautions. This level of precaution is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. SARS, SARS-CoV-2, COVID-19, MERS, Monkeypox).
- Rehydration with fluids increases the patient's ability to sweat and improves heat loss.
- Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen. Do not give to patients who have renal disease or renal transplant.
- NSAIDs should not be used in the setting of environmental heat emergencies.
- **Do not** give aspirin to a child, age ≤ 15 years.
- Agency Medical Director may require contact of medical control prior to EMT/ EMR administering any medication.



# Pain Control

## History

- Age
- Location
- Duration
- Severity (1 - 10)
- If child use Wong-Baker faces scale
- Past medical history
- Medications
- Drug allergies


## Signs and Symptoms

- Severity (pain scale)
- Quality (sharp, dull, etc.)
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

## Differential

- Per the specific protocol
- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural / Respiratory
- Neurogenic
- Renal (colic)

## Specific Complaint Protocol





# Pain Control

## Pearls

- **Recommended Exam:** Mental Status, Area of Pain, Neuro
- **Pain severity (0-10)** is a vital sign to be recorded before and after PO, IV, IO or IM medication delivery and at patient hand off. Monitor BP closely as sedative and pain control agents may cause hypotension.
- **Ketamine:**
  - Ketamine may be used in patients who are outside a Pediatric Medication/Skill Resuscitation System product.
  - Ketamine may be used in patients who fit within a Pediatric Medication/Skill Resuscitation System product only with DIRECT ONLINE MEDICAL ORDER, by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR.
- **Ketamine: appropriate indications for pain control:**
  - Patients who have developed opioid-tolerance. Sickle cell crisis patients with opioid-tolerance.
  - Patients who have obstructive sleep apnea.
  - May use in combination with opioids to limit total amount of opioid administration.
- **Ketamine: caution when using for pain control:**
  - Slow infusion or IV push over 10 minutes is associated with less side effects. Do not administer by rapid IV push.
  - Avoid in patients who have cardiac disease or uncontrolled hypertension.
  - Avoid in patients with increased intraocular pressure such as glaucoma.
  - Avoid use in combination with benzodiazepines due to decreased respiratory effort.
- **Both arms of the treatment may be used in concert. For patients in Moderate pain for instance, you may use the combination of an oral medication and parenteral if no contraindications are present.**
- **Pediatrics:**
  - For children use Wong-Baker faces scale or the FLACC score (see Assessment Pain Procedure)
  - Use Numeric (> 9 yrs), Wong-Baker faces (4-16yrs) or FLACC scale (0-7 yrs) as needed to assess pain
- **Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.**
- All patients who receive IM or IV medications must be observed 15 minutes for drug reaction in the event no transport occurs.
- Do not administer **Acetaminophen** to patients with a history of liver disease.
- Burn patients may require higher than usual opioid doses to titrate adequate pain control.
- Consider agency-specific anti-emetic(s) for nausea and/or vomiting.



# Police Custody

## History


- Traumatic Injury
- Drug Abuse
- Cardiac History
- History of Asthma
- Psychiatric History

## Signs and Symptoms

- External signs of trauma
- Palpitations
- Shortness of breath
- Wheezing
- Altered Mental Status
- Intoxication/Substance Abuse

## Differential

- Agitated Delirium Secondary to Psychiatric Illness
- Agitated Delirium Secondary to Substance Abuse
- Traumatic Injury
- Closed Head Injury
- Asthma Exacerbation
- Cardiac Dysrhythmia





# Police Custody

Patients should not be handcuffed to the stretcher.

## Pearls

- **Patient does not have to be in police custody or under arrest to utilize this protocol.**
- **Local EMS agencies should formulate a policy with local law enforcement agencies concerning patients requiring EMS and Law Enforcement services simultaneously.**
- **Agencies should work together to formulate a disposition in the best interest of the patient.**
- Patients restrained by law enforcement devices must be transported and accompanied by a law enforcement officer in the patient compartment who is capable of removing the devices. However, when rescuers have utilized restraints in accordance with Restraint Procedure, the law enforcement agent may follow the ambulance during transport.
- **All patients who receive either physical and chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.**
- The responsibility for patient care rests with the highest authorized medical provider on scene per North Carolina law.
- If an asthmatic patient is exposed to irritant/ pepper spray and released to law enforcement, all parties should be advised to immediately contact EMS if wheezing/ difficulty breathing occurs.
- All patients with decision-making capacity in police custody retain the right to participate in decision-making regarding their care and may request care or refuse care of EMS.
- If extremity/ chemical/ law enforcement restraints are applied, follow USP 5 Restraints: Physical.
- **Consider Haldol or Droperidol for patients with history of psychosis or a benzodiazepine for patients with presumed substance misuse.**
- **Haldol is acceptable treatment in pediatric patients  $\geq 12$  years old. Safety and efficacy is not established in younger ages. Contact Medical Control for advice as needed.**
- **Hyperactive Delirium with Severe Agitation:**  
Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent/ bizarre behavior, insensitivity to pain, hyperthermia and increased strength.  
Potentially life-threatening and associated with use of physical control measures, including physical restraints and Tasers.  
Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents. Alcohol withdrawal or head trauma may also contribute to the condition.  
**If patient suspected of Hyperactive Delirium with Severe Agitation suffers cardiac arrest, consider a fluid bolus, administration of calcium gluconate (or chloride), and sodium bicarbonate early.**
- Do not position or transport any restrained patient in such a way that could impact the patients respiratory or circulatory status.
- Patients exposed to chemical spray, with or without history of respiratory disease, may develop respiratory complaints up to 20 minutes post exposure.



# Seizure

## History

- Reported / witnessed seizure activity
- Previous seizure history
- Medical alert tag information
- Seizure medications
- History of trauma
- History of diabetes
- History of pregnancy
- Time of seizure onset
- Document number of seizures
- Alcohol use, abuse or abrupt cessation
- Fever


## Signs and Symptoms

- Decreased mental status
- Sleepiness
- Incontinence
- Observed seizure activity
- Evidence of trauma
- Unconscious

## Differential

- CNS (Head) trauma
- Tumor
- Metabolic, Hepatic, or Renal failure
- Hypoxia
- Electrolyte abnormality (Na, Ca, Mg)
- Drugs, Medications, Non-compliance
- Infection / Fever
- Alcohol withdrawal
- Eclampsia
- Stroke
- Hyperthermia
- Hypoglycemia

Age Appropriate	
	Airway Protocol(s) AR 1, 2, 3, 5, 6 <i>as indicated</i>
Altered Mental Status Protocol UP 4 <i>if indicated</i>	
	Childbirth/ Labor Protocol AO 1 Obstetrical Emergency Protocol AO 3 <i>if indicated</i>
Behavioral Protocol UP 17, 18, 19 <i>if indicated</i>	
Loosen any constrictive clothing Protect patient	



<b>P</b>	<b>Midazolam 2 – 2.5 mg IV / IO</b> May repeat every 3 to 5 minutes as needed <b>Maximum 10 mg</b>
<b>P</b>	<b>Peds: Midazolam 0.2 mg/kg IV / IO</b> <b>Maximum single dose 2.5 mg</b> May repeat every 3 to 5 minutes as needed <b>Maximum 10 mg</b>



# Seizure

## Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care.**
- **Brief seizure-like activity can be seen following ventricular fibrillation or ventricular tachycardia associated cardiac arrest.**
- **Status epilepticus is defined by seizure activity lasting > 5 minutes or multiple seizures without return to baseline.**
- **Most seizure activity is brief, lasting only 1 – 2 minutes, and is associated with transient hypoventilation.**
- **Be prepared for airway problems and continued seizures.**
- **Seizure activity may be a marker of closed head injury, especially in the very young, examine for trauma.**
- **Adult:**
  - Midazolam 10 mg IM is effective in termination of seizures.
  - Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.
- **Pediatrics:**
  - Midazolam 0.2 mg/kg (Maximum 5 mg) IM is effective in termination of seizures.
  - Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.
- **Do not delay administration of anti-epileptic drugs to check for blood glucose.**
- **Grand mal seizures (generalized) are associated with loss of consciousness, incontinence, and tongue trauma.**
- **Focal seizures affect only a part of the body and are not usually associated with a loss of consciousness, but can propagate to generalized seizures with loss of consciousness.**
- **Be prepared to assist ventilations especially if diazepam or midazolam is used.**
- **For any seizure in a pregnant patient, follow the AO 3 Obstetrical Emergencies Protocol.**
- **Midazolam (Versed) is shown to be as effective with IM route as Lorazepam (Ativan) is via the IV or IO route.**
- **Lorazepam (Ativan) is not as effective when administered IM. IV or IO route is preferred.**
- **Diazepam (Valium) is not effective when administered IM. Give IV or Rectally.**
- **Optimal conditions for patients refusing transport following a seizure:**

Known history of seizures/epilepsy	Seizure not associated with drugs or alcohol
Full recovery to baseline mental status	Only 1 seizure episode in the past hour
No injuries requiring treatment or evaluation	Seizure not associated with pregnancy
Adequate supervision	



# Suspected Stroke

## History


- Previous CVA, TIA's
- Previous cardiac / vascular surgery
- Associated diseases: diabetes, hypertension, CAD
- Atrial fibrillation
- Medications (blood thinners)
- History of trauma
- Sickle Cell Disease
- Immune disorders
- Congenital heart defects
- Maternal infection / hypertension

## Signs and Symptoms

- Altered mental status
- Weakness / Paralysis
- Blindness or other sensory loss
- Aphasia / Dysarthria
- Syncope
- Vertigo / Dizziness
- Vomiting
- Headache
- Seizures
- Respiratory pattern change
- Hypertension / hypotension

## Differential

- See Altered Mental Status
- TIA (Transient ischemic attack)
- Seizure
- Todd's Paralysis
- Hypoglycemia
- Stroke
  - Thrombotic or Embolic (~85%)
  - Hemorrhagic (~15%)
- Tumor
- Trauma
- Dialysis / Renal Failure





# Suspected Stroke

Destination decision is a critical component of the prehospital management of stroke and this decision has increased in complexity with the availability of endovascular therapy for patients with large vessel occlusion. Optimizing patient and system outcomes requires a balance between over- and under-triage of these patients to specialty centers capable of providing endovascular therapy. If a patient has a score equal to or greater than that listed below according to the scale used, consider direct transport to a Comprehensive Stroke Center that can provide endovascular therapy.

## Pearls

- **Recommended Exam:** Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro
- **Items in Red Text** are key performance measures used in the EMS Acute Stroke Care Toolkit.
- **Acute Stroke care is evolving rapidly. Time of Onset/ Last Seen Normal may be changed at any time depending on the capabilities and resources of your regional hospital(s).**
- **Refer to your Stroke: EMS Triage and Destination Plan which should be updated when community resources change.**
- **Time of Onset or Last Seen Normal:**  
One of the most important items the pre-hospital provider can obtain, of which all treatment decisions are based.  
Be very precise in gathering data to establish the time of onset and report as an actual time (i.e. 13:47 NOT “about 45 minutes ago.”)  
Without this information patient may not be able to receive thrombolytics at facility.  
Wake up stroke: Time starts when patient last awake or symptom free.
- **Time of Symptom Discovery:**  
Time when symptoms of stroke are first noticed by patient, bystanders, witnesses, or family/ caregivers.
- **Sources of information pertaining to Last Known Well Time or Symptoms Onset:**  
You are often in the best position to determine the actual Time of Onset while you have family, friends or caretakers available.  
Often these sources of information may arrive well after you have delivered the patient to the hospital.  
Delays in decisions due to lack of information may negatively impact patient care.  
Obtain contact information (phone number and name) of best witnesses and give to hospital providers.
- The **Reperfusion Checklist** should be completed for any suspected stroke patient as time allows.
- If possible place 2 IV sites, preferably above the wrists, and if possible both in the left upper extremity.
- **Blood Draw:**  
Many stroke centers utilize EMS venous blood samples. Follow your local policy and procedures.
- The differential listed in the UP 4 Altered Mental Status Protocol should also be considered.
- Be alert for airway problems (swallowing difficulty, vomiting/aspiration).
- Hypoglycemia can present as a localized neurologic deficit, especially in the elderly.
- Document the EMS Stroke Screen, Stroke Severity Score, and Stroke Alert notification time in the ePCR or PCR.
- Agencies may use validated pre-hospital stroke screen of choice.
- **Pediatrics:**  
Strokes do occur in children, they are slightly more common in ages < 2, in boys, and in African-Americans.  
Newborn and infant symptoms consist of seizures, extreme sleepiness, and using only one side of the body.  
Children and teenagers symptoms may consist of severe headaches, vomiting, sleepiness, dizziness, and/or loss of balance or coordination.



# Suspected Sepsis

## History

- Duration and severity of fever
- Past medical history
- Medications/ Recent antibiotics
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Indwelling medical device
- Last acetaminophen or ibuprofen
- Recent Hospital/ healthcare facility
- Bedridden or immobile
- Elderly and very young – at risk
- Prosthetic device / indwelling device

## Signs and Symptoms

- Warm
- Flushed
- Sweaty
- Chills/ Rigors
- Delayed cap refill
- Mental status changes

### Associated Symptoms (Helpful to localize source)

- myalgias, cough, chest pain, headache, dysuria, abdominal pain, rash

## Differential

- Infections: UTI, Pneumonia, skin/ wound
- Cancer/ Tumors/ Lymphomas
- Medication or drug reaction
- Connective tissue disease: Arthritis, Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis
- Hypoglycemia/hypothermia
- MI/ CVA

Consider: Contact, Droplet, and Airborne Precautions	
Temperature Measurement Procedure <i>if available</i>	
	<b>Fever/ Infection Control Protocol UP 10</b> <i>if needed</i>
	<b>Altered Mental Status Protocol UP 4</b> <i>if needed</i>
<b>B</b>	12 Lead ECG Procedure
	<b>IV or IO Access Protocol UP 6</b> <i>If indicated</i>
<b>P</b>	Cardiac Monitor

### Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60

Ages  $\geq$  1 month: SBP < 70


Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90

Ages  $\geq$  65: SBP < 110

All ages Shock Index:  
HR > SBP

Exit to  
Age Appropriate  
Condition Appropriate  
Protocol(s)



### MAP (Mean Arterial Pressure)

$$\text{SBP} + 2(\text{DBP})$$

3

Monitor usually calculates this value on screen

<b>Adult SIRS Criteria</b>
Temperature $\geq 100.4^{\circ} \text{ F}$ ( $38^{\circ} \text{ C}$ ) Or $\leq 96.8^{\circ} \text{ F}$ ( $36^{\circ} \text{ C}$ )
<b>AND</b>
Any 1 of the following: HR > 90 RR > 20 EtCO $< 25 \text{ mmHg}$
<b>Adult qSOFA Criteria</b>
SBP $\leq 100 \text{ mmHg}$ RR $\geq 22$ AMS or new mental status change
<b>Pediatrics SIRS Criteria</b>
Temperature Same as adult
<b>AND</b>
Heart Rate 1 month – 1 year > 180 2 – 5 years > 140 6 – 12 years > 130 13 – 18 years > 120

<b>A</b>	Venous Access Blood Draw <i>if applicable</i>
<b>P</b>	<b>Normal Saline 500 mL Bolus</b> Repeat as needed Titrate SPB $\geq 90 \text{ mmHg}$ MAP $> 65 \text{ mmHg}$ <b>Maximum 2 L</b>
	<b>Peds: 20 mL/kg IV / IO</b> Repeat to titrate Age Appropriate SBP $\geq 70 + 2 \times \text{Age}$ <b>Maximum 60 mL/kg</b>
	<b>This Space Left Blank Intentionally</b>

Age Appropriate  
 Hypotension/ Shock  
Protocol AM 5/ PM 3

	<b>Notify Destination or Contact Medical Control</b>	
--	--	--

**UP 15**



# Suspected Sepsis

Abdominal pain is not a typical presentation of sepsis, but some patients do have nausea and vomiting. Septic shock is the result of a massive systemic inflammatory response to infection by gram-negative or gram-positive aerobes, anaerobes, fungi, or viruses. Gram-negative organisms appear to be the primary cause of sepsis, especially hospitalized patients. More patients are remaining at home and have medical devices inserted, making the patients prone to infection. Many of these patients have compromised immune systems, putting them at even greater risk for sepsis. The basis of septic shock and systemic inflammatory response syndrome (SIRS) is a complex process of inflammatory response and multisystem organ failure. Two or more of the following criteria must be met for a diagnosis of SIRS:

- Temp. >100F (38C) or <97F (36C)
- Pulse rate > 90 beats/min
- Resp. rate >20 bpm or PaCO<sub>2</sub> <32 mmHg
- WBC >12,000/mm<sup>3</sup>, <4,000/mm<sup>3</sup>, or >10% band neutrophilia

The Robson Prehospital Severe Sepsis Screening tool has a 75% success rate in identifying sepsis. It is key that prehospital providers relay the potential for sepsis to hospital personnel.

- Temp. >100.9F (38.3C) or <96.8F (36.0C)
- Pulse rate > 90 bpm
- Resp. rate > 20 bpm
- Acutely altered mental status
- FSBS > 119 mg/dL – unless diabetic

If these findings are present in a patient with a history that is suggestive of infection, sepsis should be considered.

## Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Recommended Exam Pediatrics: In childhood, physical assessment reveals important clues for sepsis. Look for mental status abnormalities such as anxiety, restlessness, agitation, irritability, confusion, or lethargy. Cardiovascular findings to look for include cool distal extremities, capillary refill >3 seconds, or mottled skin.**
- **Sepsis is a life threatening condition where the body's immune response to infection injures its own tissues and organs.**
- **Severe sepsis is a suspected infection with 2 or more SIRS criteria (or qSOFA) along with organ dysfunction, such as AMS, hypotension, or hypoxia.**
- **Septic shock is severe sepsis and poor perfusion unimproved after fluid bolus.**
- Agencies administering antibiotics should inquire about drug allergies specific to antibiotics or family of antibiotics.
- Following each fluid bolus, assess for pulmonary edema. Consider administration of agency specific vasopressor.
- Supplemental oxygen should be given and titrated to oxygenation saturation  $\geq 92\%$ .
- EKG should be obtained with suspected sepsis, but should not delay care in order to obtain.
- Abnormally low temperatures increase mortality and are found often in geriatric patients.
- Quantitative waveform capnography can be a reliable surrogate for lactate monitoring in detecting metabolic distress in sepsis patients. EtCO<sub>2</sub> < 25 mm Hg are associated with serum lactate levels  $> 4$  mmol/L.
- Patients with a history of liver failure should not receive acetaminophen.
- **Droplet precautions** include standard PPE plus a standard surgical mask for providers who accompany patients in the back of the ambulance and a surgical mask or NRB O<sub>2</sub> mask for the patient. This level of precaution should be utilized when influenza, meningitis, mumps, streptococcal pharyngitis, and other illnesses spread via large particle droplets are suspected. A patient with a potentially infectious rash should be treated with droplet precautions.
- **Airborne precautions** include standard PPE plus utilization of a gown, change of gloves after every patient contact, and strict hand washing precautions. This level of precaution is utilized when multi-drug resistant organisms (e.g. MRSA), scabies, or zoster (shingles), or other illnesses spread by contact are suspected.
- **All-hazards precautions** include standard PPE plus airborne precautions plus contact precautions. This level of precaution is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. **SARS, SARS-CoV-2, COVID-19, MERS, Monkeypox**).
- Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen.
- Agency Medical Director may require contact of medical control prior to EMT / MR administering any medication.
- **Sepsis Screen:**  
Agencies may use Adult / Pediatric Systemic Inflammatory Response Syndrome (SIRS) criteria or quickSOFA (qSOFA) criteria.  
Receiving facility should be involved in determining Sepsis Screen utilized by EMS.



# Syncope

## History

- Cardiac history, stroke, seizure
- Occult blood loss (GI, ectopic)
- Females: LMP, vaginal bleeding
- Fluid loss: nausea, vomiting, diarrhea
- Past medical history
- Medications

## Signs and Symptoms

- Loss of consciousness with recovery
- Lightheadedness, dizziness
- Palpitations, slow or rapid pulse
- Pulse irregularity
- Decreased blood pressure

## Differential

- Vasovagal
- Orthostatic hypotension
- Cardiac syncope
- Micturition / Defecation syncope
- Psychiatric
- Stroke
- Hypoglycemia
- Seizure
- Shock (see Shock Protocol)
- Toxicological (Alcohol)
- Medication effect (hypertension)
- PE
- AAA

	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 <b><i>if indicated</i></b>
	Blood Glucose Analysis Procedure
<b>B</b>	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
<b>P</b>	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <b><i>if indicated</i></b>
	Age Appropriate Cardiac Protocol(s) <b><i>if indicated</i></b>
	Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 <b><i>if indicated</i></b>
	Multiple Trauma Protocol TB 6 Spinal Motion Restriction Procedure/ Protocol TB 8 <b><i>if indicated</i></b>

## Age Specific Blood Pressure indicating possible shock

**Age 0 – 28 days: SBP < 60**  
**Ages  $\geq$  1 month: SBP < 70**  
**Age 1 – 9: SBP < 70 + (2x Age)**

**Ages 10 – 64: SBP < 90**  
**Ages  $\geq$  65: SBP < 110**

**All ages Shock Index:  
HR > SBP**

Serious Signs/ Symptoms  
Hypotension, poor perfusion, shock

YES

	IV or IO Access Protocol UP 6 Consider 2 Large Bore sites
<b>A</b>	<b>Normal Saline 500 mL Bolus</b> Repeat as needed Titrate SPB $\geq$ 90 mmHg <b>Maximum 2 L</b>  <b>Peds: 20 mL/kg IV / IO</b> Repeat as needed Titrate to Age Appropriate SBP $\geq$ 70 + 2 x Age <b>Maximum 60 mL/kg</b>

Exit to

Age Appropriate  
Condition Appropriate  
Protocol(s)

Notify Destination or Contact Medical Control



# Syncope

## Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro
- **Syncope is both loss of consciousness and loss of postural/ muscle tone with collapse. Symptoms preceding the event are important in determining etiology.**
- **Syncope often is due to a benign process but can be an indication of serious underlying disease in both the adult and pediatric patient.**
- **Often patients with syncope are found normal on EMS evaluation. In general patients experiencing syncope require cardiac monitoring and emergency department evaluation.**
- **Differential should remain wide and include:**

Cardiac arrhythmia	Neurological problem	Choking	Pulmonary embolism
Hemorrhage	Stroke	Respiratory	Hypo or Hyperglycemia
GI Hemorrhage	Seizure	Sepsis	
- **High-risk patients:**

Age $\geq$ 60	Syncope with exertion
History of CHF	Syncope with chest pain
Abnormal ECG	Syncope with dyspnea
- **Abdominal/ back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and/ or lower extremity pain or diminished pulses, especially in patients over 50 and/ or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients  $> 35$ , diabetics, and/ or women especially with upper abdominal complaints.**
- **Heart Rate: Tachycardia is one of the first clinical signs of dehydration, typically increases as dehydration becomes more severe.**
- **Syncope with no preceding symptoms or event may be associated with an arrhythmia.**
- **Assess for signs and symptoms of trauma if associated or questionable fall with syncope.**
- **Consider dysrhythmias, GI bleed, ectopic pregnancy, and seizure as possible causes of syncope.**
- **In general these patients should be transported: Patients who experience syncope associated with headache, neck pain, chest pain, abdominal pain, back pain, dyspnea, or dyspnea on exertion need prompt medical evaluation.**
- **More than 25% of geriatric syncope is cardiac dysrhythmia based.**



# Behavioral Health Crisis

## History

- Situational crisis
- Psychiatric illness/medications
- Injury to self or threats to others
- Medic alert tag
- Substance abuse / overdose
- Diabetes

## Signs and Symptoms

- Anxiety, agitation, confusion
- Affect change, hallucinations
- Delusional thoughts, bizarre behavior
- Combative violent
- Expression of suicidal / homicidal thoughts

## Differential

- Altered Mental Status
- Alcohol Intoxication
- Toxin / Substance abuse
- Medication effect / overdose / withdrawal
- Depression
- Bipolar (manic-depressive)
- Schizophrenia
- Anxiety disorders

Call for help

Call for additional resources

Stage prior to arrival

or

Withdraw from scene until safe

Screen patient for weapons  
Screen for scene safety

Assess for underlying medical or traumatic condition causing behavioral disturbance

Age Appropriate Protocol(s)

### Establish rapport

- Genuine respect for feelings/ circumstances
- Active listening
- Eye contact and at meet at eye level

### Create a quiet and safe environment

- Only 1 provider talks to patient to limit stimuli
- Decrease unnecessary stimuli

### Identify major problem or crisis

- "What happened to upset you?"
- "How are you feeling right now?"

### Assess for suicidal and/or homicidal thoughts

### Identify major problem or crisis

- "What happened to upset you?"
- "How are you feeling right now?"

### BARS

1 – Difficult or unable to wake

2 – Asleep, but responds normally to verbal or physical stimuli

3 – Drowsy, appears sedated

4 – Quiet and awake (normal activity)

5 – Overt activity (physical or verbal)  
Agitated but not disruptive

6 – Extremely or continuously active,  
Agitated, disruptive, but not violent

7 – Violent, requires restraint  
Agitated and violent

Exit to  
Behavioral

Agitation/ Sedation Guide  
UP 18

Hyperactive Delirium with  
Severe Agitation  
UP 19

Age Appropriate  
Protocol(s)



Assess and score: BARS  
Behavioral Activity Rating Scale

P Evaluation and Screening  
Mental Health and Substance Use Protocol  
CIT Paramedic Only  
if available  
Page 3

Triage and Alternative Destination  
Mental Health / Substance Abuse  
if available

Page 3

T Notify Destination or  
Contact Medical Control

## Universal Protocol Section



# Behavioral Health Crisis


## Pearls

- **Recommended Exam:** Mental Status, Skin, Heart, Lungs, Neurologic status
- **Crew/ responders safety is the main priority. Call for assistance, stage, or withdraw from scene if necessary.**
- **Law Enforcement:**
  - Any patient who is handcuffed or restrained by Law Enforcement and transported by EMS, must be accompanied by law enforcement during transport.
  - Patient should not be transported with upper extremities hand-cuffed behind back as this prevents proper assessment and could lead to injury.
  - Consider multidisciplinary coordination with law enforcement to approach verbal de-escalation, restraint, and/ or USP 6 Restraints: Therapeutic Take-down Procedure.
- **Maintain high-index of suspicion for underlying medical or traumatic disorder causing or contributing to behavioral disturbance. Medical causes more likely in ages < 12 or > 40.**
- **General communications techniques**
  - Ask Open-ended questions (questions that cannot be answered with a yes/no)
    - “Tell me how we can help you?” “What caused you to call 911 today?”*
  - Active listening (stay engaged, be able to summarize patient’s story, use your body language to convey listening)
    - Eye contact, nodding your head, periodically repeating back part of patient’s story*
  - Encouraging (remain positive, convey interest in patient’s crisis)
    - “Tell me more about that...”*
  - Clarifying questions (ask patient to rephrase or repeat if you don’t understand)
    - “I’m not sure I understand, can you...?”*
  - Emotional labeling (naming emotions patient is demonstrating, validating emotions)
    - “You look upset.” “You seem angry.”*
  - Conversational pause (okay to allow a period of silence for patient to process information)
- **Behavioral health disturbance incidents are increasing and commonly involve the following:**

Substance misuse	Psychosis
Depression/ Anxiety/ Stress Reactions / Bipolar	Schizophrenia or schizophrenia-like illness
- **Restraints:**
  - All patients who receive either physical or chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.
  - Do not position or transport any restrained patient in such a way that could impact the patients respiratory or circulatory status (i.e. prone position)
- **Maintain high-index of suspicion for medical, trauma, abuse, or neglect causes:**
  - Hypoglycemia, hyperglycemia, overdose, substance abuse, hypoxia, head injury, shock, sepsis, stroke, etc.
  - Domestic violence, child or geriatric abuse/ neglect.
- **Extrapyramidal reactions:**
  - Condition causing involuntary muscle movements or spasms typically of the face, neck and upper extremities. May present with contorted neck and trunk with difficult motor movements. Typically an adverse reaction to antipsychotic drugs like Haloperidol and may occur with your administration. When recognized, give **Diphenhydramine 50 mg IV / IO / IM / PO** in adults or **1 mg/kg IV / IO / IM / PO** in pediatrics, **Maximum 50 mg**.
- **May add page 3 to protocol for specific for local mental health and / or substance misuse resources or destinations.**




# Behavioral CIT Paramedic (Optional)






# Behavioral Agitation/ Sedation Guide





# Behavioral Hyperactive Delirium With Severe Agitation

## Universal Protocol Section





# Behavioral Hyperactive Delirium With Severe Agitation

## Pearls

- **Ketamine for sedation purposes:**

**Ketamine may be used in pediatric patients who fit within a Pediatric Medication/ Skill Resuscitation System product, ≤ 15 years of age, or ≤ 49 kg) with DIRECT ONLINE MEDICAL ORDER by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR only.**

- **Hyperactive Delirium with Severe Agitation:**

Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent/ bizarre behavior, insensitivity to pain, hyperthermia and increased strength.

Potentially life-threatening and associated with use of physical control measures, including physical restraints.

Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents.

Alcohol or substance withdrawal as well as head trauma may also contribute to the condition.

- **Restraint use:**

Physical restraints are not contraindicated in agitated or excited delirium, but you must use caution.

Once sedated, prevent patient from continued struggle, which can worsen metabolic condition.

Prevent patient from assuming a prone position for prolonged period, move to supine position as quickly as possible.

**Team approach for sedation and Restraint Therapeutic Take Down Procedure USP-6:**

- 1 provider for each limb.

- 1 provider to lead restraint, maintain airway and control head.

- 1 Provider to administer medication.

Do not position prone or prone with restraints, as this can impede respiration and ventilation.

- **Hyperthermia: Assess for and treat hyperthermia.**



# Sickle Cell Crisis

## History

- Past medical history
- Medications
- Recent illness
- Prior pain crisis location
- Pain regimen at home

## Signs and Symptoms

- Pain
- One sided paralysis / weakness
- Difficulty walking / speaking
- Sudden vision changes
- Unexplained numbness
- Severe headache
- Fever
- SOB
- Chest Pain
- Abdominal Pain
- Pallor

## Differential

- Sickle Cell Pain Crisis
- Aplastic Crisis
- Acute Chest Syndrome
- Alcohol / drug use
- Toxic ingestion
- Seizure
- Stroke
- Altered baseline mental status
- Sepsis
- Pneumonia

Apply Hot Packs to affected areas: especially joints and areas of increased pain

Provide emotional support  
Calm and continual reassurance

Fluids indicated if signs of shock or hypotensive

<b>B</b>	Blood Glucose Analysis as needed Assess Pain Severity 12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
<b>P</b>	Cardiac Monitor  Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Manage Airway as appropriate Protocol UP 13 <i>if indicated</i>
	Suspected Stroke Protocol AM 7 <i>if indicated</i>
	Suspected Stroke Protocol AR 1 & AR 5 <i>if indicated</i>
	Hypotension/ Shock Protocol AM 5 & PM 3 <i>if indicated</i>

**Sickle Cell Crisis**  
- Acute Chest Syndrome  
- Abdominal Crisis  
- Joint Crisis

YES

Pain Control  
Protocol UP 11

Notify Destination or  
Contact Medical Control



# Sickle Cell Crisis

## PEARLS

Patterns of an acute sickle cell crisis are now recognizable. They are based on the part of the body where the crisis occurs.

Is this their typical pain crisis? If not, what is different about it?

Any fever, SOB, pleuritic chest pain?

### ***Acute chest syndrome:***

Sudden acute chest pain with coughing up of blood can occur. Low-grade fevers can be present. The person is usually short of breath. If a cough is present, it often is nonproductive. Acute chest syndrome is common in a young person with sickle cell disease. Chronic (long-term) sickle cell lung disease develops over time because the acute and subacute lung crisis leads to scarred lungs as well as other problems.

### ***Abdominal crisis:***

The pain associated with the abdominal crisis of sickle cell disease is constant and sudden. It becomes unrelenting. The pain may or may not be localized to any one area of the abdomen. Nausea, vomiting, and diarrhea may or may not occur.

### ***Joint crisis:***

Acute and painful joint crisis may develop without a significant traumatic history. Its focus is either in a single joint or in multiple joints. Often the connecting bony parts of the joint are painful. Range of motion is often restricted because of the pain. Avascular necrosis of the hips can occur, causing permanent damage.

### ***Stroke:***

Many sickle cell patients can present with strokes at a younger age than average. Ensure hospital pre-notification indicating it is a sickle cell patient as the standard of care for a Sickle Cell Stroke can involve exchange therapy as opposed to other treatments such as tPA.

# General Medical EMS Triage and Destination Plan




## General Medical Patient

- ❖ A patient with signs and/or symptoms that cannot be categorized into STEMI, Stroke, Pediatric or Trauma and Burn Triage and Destination Plans.

The purpose of this plan is to:

- ❖ Identify general medical patients who call 911 or present to EMS
- ❖ Minimize field decision making on transport destinations
- ❖ Identify those patients that should be transported to the geographically closest facility based on complaint and facility capabilities
- ❖ Identify those patients that may be transported to a geographically distant facility based on complaint and facility capabilities
- ❖ Maximize EMS system resource utilization
- ❖ Provide quality EMS service and patient care to the EMS Systems citizens
- ❖ Continuously evaluate the EMS System based on North Carolinas EMS performance measures.



## General Medical EMS Triage and Destination Plan

### Pearls and Definitions

- ❖ All General Medical patients should be triaged and transported using this plan. This plan is in effect 24/7/365
- ❖ Appropriate Facility = a hospital which provides emergency care 24/7/365 that is determined, based on complaint, to be capable of addressing the primary patient condition.
- ❖ Non-Critical Injury = Injuries that may be considered time sensitive but not life threatening, i.e. fracture with loss of PMS
- ❖ Critical Illness = Any potentially life-threatening complaint that does not meet the criteria for STEMI, Stroke, Pediatric or Trauma and Burn Triage and Destination Plans.
- ❖ Non-Critical Illness = Any non life-threatening complaint (illness) that meets the guidelines for medical necessity of ambulance transport.
- ❖ Individual insurance plan requirements should not be a factor in determining the appropriate facility. Consideration of insurance plan requirements may be construed as treatment based on the ability to pay and such practices are prohibited.

# Pediatric EMS Triage and Destination Plan




## Pediatric Patient

- Any patient less than 16 years of age with a life-threatening illness (Not Trauma)
- Life Threatening Illness**
- Decreased mental Status (GCS<13)
- Non-responsive respiratory Distress
- Intubation
- Post Cardiac Arrest
- Non-responsive Hypotension (shock)
- Severe Hypothermia or Hyperthermia
- Status Epilepticus
- Potential Dangerous Envenomation
- Life Threatening Ingestion/Chemical Exposure
- Children with Special Healthcare Needs (and destination choice based on parental request)

## The purpose of this plan is to:

- Rapidly identify acute Pediatric patients who call 911 or present to EMS with a life-threatening illness
- Minimize the time from EMS contact to definitive care
- Quickly diagnose patients with pediatric life-threatening illness for EMS treatment and stabilization
- Rapidly identify the best hospital destination based on symptom onset time, vital signs, response to treatment and predicted transport time
- Early activation/notification to the hospital prior to the patient arrival.
- Minimize scene time with a 'load and go' approach
- Provide quality EMS service and patient care to the EMS Community
- Continuously evaluate the EMS System based on North Carolinas EMS performance measures



## Pediatric EMS Triage and Destination Plan

### Pearls and Definitions

- All Pediatric patients with a life threatening illness must be triaged and transported using this plan. This plan is in effect 24/7/365
- The Trauma and Burn Triage and Destination Plan should be used for all injured patients regardless of age
- All patient care is based on EMS Suspected Stroke Protocol
- Pediatric Capable Hospital = a hospital which provides emergency and pediatric intensive care capability including but not limited to:
  - Emergency Department staffed 24 hours per day with board certified Emergency Physicians.
  - An inpatient Pediatric Intensive Care Unit (with a physician pediatric intensivist available in-house or on call 24/7/365)
  - Accepts all patients regardless of bed availability
  - Provides outcome and performance measure feedback to the EMS including case review
- Community Hospital = a local hospital within the EMS Systems service area which provides emergency care but does not meet the criteria of a Pediatric Capable Hospital
- Pediatric Specialty Care Transport Program = an air or ground based specialty care transport program that has specific pediatric training and equipment addressing the needs of a pediatric patient that can assume care of a pediatric patient from EMS or a Community hospital and transport to a Pediatric Capable Hospital.

# STEMI

## EMS Triage and Destination Plan




**STEMI Patient**  
**(ST Elevation Myocardial Infarction)**

- ❖ Cardiac Symptoms greater than 15 minutes and less than 12 hours  
**And**
- ❖ 12 lead ECG criteria of 1mm ST elevation in 2 or more contiguous leads.  
**or**
- ❖ Left Bundle Branch Block  
**NOT KNOWN to be present in the past**

**The purpose of this plan is to:**

- ❖ Rapidly identify STEMI patients who call 911 or present to EMS
- ❖ Minimize the time from onset of STEMI symptoms to coronary reperfusion
- ❖ Quickly diagnose a STEMI by 12 lead ECG
- ❖ Complete a reperfusion checklist (unless being transported directly to a PCI hospital) to determine thrombolytic eligibility
- ❖ Rapidly identify the best hospital destination based on a symptom onset time, reperfusion checklist and predicted transport time
- ❖ Early activation/notification to the hospital prior to the patient arrival
- ❖ Minimize scene time to 15 minutes or less (including a 12 lead ECG)
- ❖ Provide quality EMS service and patient care to the EMS Systems citizens.
- ❖ Continuously evaluate the EMS System based on North Carolinas EMS performance measures



**Pearls and Definitions**

- ❖ All STEMI patients must be triaged and transported using this plan. This plan is in effect 24/7/365
- ❖ All patient care is based on EMS Chest Pain and STEMI Protocol
- ❖ Consider implementing a prehospital thrombolytic program if a STEMI patient cannot reach a hospital within 90 minutes using air or ground EMS transport
- ❖ **PCI (Percutaneous Coronary Intervention) Capable Hospital** = a hospital with an emergency interventional cardiac catheterization laboratory capable of providing the following services to acute STEMI patients. Free standing emergency departments and satellite facilities are not considered part of the PCI hospital.
  - ❖ 24/7 PCI capability within 30 minutes of notification (interventional cardiologist present at the start of the case)
  - ❖ Single Call Activation number for use by EMS
  - ❖ Accepts all patients regardless of bed availability
  - ❖ Provides outcome and performance measure feedback to the EMS including case review
- ❖ **Non-PCI Hospital** = a local hospital within the EMS Systems service area which provides emergency care including thrombolytic administration to an Acute STEMI patient but does not provide PCI services
- ❖ **Specialty Care Transport Program** = an air or ground based specialty care transport program which can assume care of an acute STEMI patient from EMS or a Non-PCI hospital and transport the patient to a PCI capable hospital

**Rowan County EMS System**

This plan has been altered from the original NCCEP Protocol by the local EMS Medical Director

**2022**

**STEMI EMS Triage and Destination Plan**



# STROKE and LVO Stroke

## EMS Triage and Destination Plan

### Stroke Patient

- Signs and symptoms of an acute Stroke identified on EMS Stroke Screen Assessment.
- **Last Known Well (LKW)**
- Refer to UP 14 Suspected Stroke Protocol

### The Purpose of this plan:

- Use plan in conjunction with UP 14 Suspected Stroke Protocol
- Rapidly identify acute Stroke patients presenting to EMS system and minimize the time from Stroke onset to definitive care
- Rapidly identify most appropriate facility destination in region
- Provide quality EMS service and patient care to the EMS system's citizens
- Maintain performance improvement of the EMS system based on NC Stroke Performance measures

CPSS  
LAMS  
RACE - Preferred  
VAN

LVO Suspected Score:  
CPSS  $\geq$  2  
LAMS  $\geq$  3  
RACE  $\geq$  4 - Preferred  
Positive VAN

RACE  
Facial palsy  
Arm/leg motor function  
Head and gaze deviation

Positive Stroke Screen Tool  
and/or  
Positive Stroke Screen/  
Stroke Severity Tool

Last Known Well  
Greater than  
24 Hours

Transport to Destination A or B:  
Nearest facility or certified stroke center  
honoring patient/family preference

Absolute contraindications for fibrinolytic:  
Prior intracranial hemorrhage  
Ischemic stroke within 3 months  
Known cerebrovascular abnormality  
Known malignant intracranial tumor  
Significant closed-head trauma (recent)  
Significant facial trauma (recent)

LVO Suspected  
Stroke Severity Score  
Greater than/equal to  
.

Contraindications  
to fibrinolysis?  
Reperfusion Checklist  
Or  
See box →

Last Known Well  
Greater than  
4.5 hours

Transport to Destination B

May transport to Destination A  
• If estimated transport time to Destination B  
is greater than 45 minutes

NO

NO

Transport to Destination A

May transport to Destination B  
• If estimated transport time to Destination A  
is greater than na minutes  
And  
• If total time from LKW and transport time will  
not exceed 4 hours by facility arrival

**DESTINATION A**  
Rapid / Early Notification of receiving facility  
Activation of Stroke Team  
Nearest Certified Stroke Center and/or  
patient/family preference

Novant Health - Rowan Medical Center  
Atrium Northeast  
Lake Norman Regional Medical Center  
Iredell Memorial Hospital  
Novant Health - Forsyth Medical Center  
Atrium Wake Forest

**DESTINATION B**  
Rapid / Early Notification of receiving facility  
Activation of Stroke Team  
Thrombectomy Capable Stroke Center (TSC)  
Comprehensive Stroke Center (CSC)

Novant Health - Forsyth Medical Center  
Atrium Wake Forest  
Novant Health Presbyterian  
Atrium - Main



# STROKE

## EMS Triage and Destination Plan

### Pearls

- Use the AHA resource document for assistance on transport decision-making:  
<https://www.ahajournals.org/doi/10.1161/STROKEAHA.120.033228>
- Agencies may reconfigure this document to align with EMS and regional stroke care resources.
- If unstable airway or unstable hemodynamic condition may divert transport to closest appropriate facility.
- All Stroke patients should be triaged and transported using this plan.
- Expectation: EMS agency will collaborate with their regional stroke resources to establish point-to-point and inter-facility transport workflows for patient requiring higher level of acute care in consideration of potential EMS system impact and regional approach to stroke care.
- **Stroke Severity/Large Vessel Occlusion (LVO) Tool and Score:**  
Score severity and LVO score level should be set based on collaboration with all stroke centers where EMS agency routinely transports in the region. Majority of strokes are NOT large vessel occlusion strokes and inappropriately low severity scores can result in an over-triage of patients to TSC / CSC negatively impacting both the EMS and healthcare system.
- **EMS Transport Times in Destination Decisions:**  
EMS Transport times should be set based on collaboration with all stroke centers where EMS agency routinely transports in the region.
- **Reperfusion Checklist and contraindications to fibrinolysis in acute stroke patients:**  
Systems may use the Reperfusion Checklist or may establish regionally agreed upon absolute contraindications.
- Many EMS systems have a variety of stroke certified medical facilities within similar transport time parameters.
- Destination choices should use regional stroke system of care plans and patient/family preferences in choosing most medically appropriate facility.
- **Modality of transport in acute stroke depends on multiple factors, but safest and fastest should be considered, whether ground EMS, air medical EMS, or specialty/critical care ground transport.**  
Consider air medical transport options when no Comprehensive or Thrombectomy Capable Stroke Centers are within a 60 minute total transport time.
- **Acute Stroke-Ready Hospital Components:**  
Director of stroke care, written emergency stroke care protocols and transfer agreements with a neurosurgical capable hospital, 24-hour CT capability, and ability to administer thrombolytics.  
Facility may have Telemedicine / Telestroke capability for consultation with neurologic specialist.
- **Primary Stroke Center:**  
Has same capabilities as Acute Stroke-Ready Hospital.  
Accredited and certified by the Joint Commission.
- **Thrombectomy-Capable Stroke Center:**  
Has same capabilities as Primary Stroke Center.  
Capable of providing mechanical thrombectomy with no day or hour limitation.
- **Comprehensive Stroke Center:**  
Has same capabilities as a Primary Stroke Center.  
Capable of offering full spectrum, state-of-the art Stroke care with no day or hour limitation.  
Ability to treat stroke patients with catheter-based procedures to remove or dissolve blood clots.  
Accredited and certified by the Joint Commission.
- **Guidelines only for prioritization of hospital choices based on capabilities:**  
Prioritize rural hospitals that have formal agreements with Comprehensive Stroke Center or Thrombectomy-Capable Stroke Center with access to expert stroke consultation.  
Prioritize rural hospitals with stroke center certification and/or those actively engaged in stroke center certification and who track their performance on evidenced-based stroke care.  
Prioritize Primary Stroke Centers over Acute Stroke Ready Hospitals when total transport time is < 30 minutes difference.  
Prioritize Comprehensive Stroke Center over Thrombectomy-Capable Stroke Center when total transport time is < 30 minutes difference.

# Trauma and Burn


## EMS Triage and Destination Plan



**Trauma or Burn Patient** = Any patient (regardless of age) with significant injury or burn

### The purpose of this plan is to:

- ❖ Rapidly identify injured or burned patients who call 911 or present to EMS
- ❖ Minimize the time from injury to definitive care for critical injuries or burns
- ❖ Quickly identify life or limb threatening injuries for EMS treatment and stabilization
- ❖ Rapidly identify the best hospital destination based on time of injury and predicted transport time
- ❖ Early activation/notification to the hospital of a critically injured or burned patient prior to patient arrival
- ❖ Minimize scene time to 10 minutes or less from patient extrication with a 'load and go' approach
- ❖ Provide quality EMS service and patient care to the EMS Systems citizens.
- ❖ Continuously evaluate the EMS System based on North Carolinas EMS performance measures.



### Trauma and Burn EMS Triage and Destination Plan

### Pearls and Definitions

- ❖ **All Injury and Burn patients must be triaged and transported using this plan. This plan is in effect 24/7/365**
- ❖ All patient care is based on EMS Trauma Protocols
- ❖ **Designated Trauma Center** = a hospital that is currently designated as a Trauma Center by the North Carolina Office of Emergency Medical Services. Trauma Centers are designated as Level 1, 2, or 3 with Level 1 being the highest possible designation. Free standing emergency departments and satellite facilities are not considered part of the Trauma Center
- ❖ **Burn Center** = an ABA verified burn Center co-located with a designated Trauma Center
- ❖ **Community Hospital** = a local hospital within the EMS Systems service area which provides emergency care but has not been designated as a Trauma Center
- ❖ **Specialty Care Transport Program** = an air or ground based specialty care transport program which can assume care of an acutely injured patient from EMS or a Community Hospital and transport the patient to a Designated Trauma Center



# Drug List



**ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.**

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Acetaminophen (Tylenol)</b>  <b>NCCEP Protocol:</b> * 7-Pain Control-Adult * 46-Pain Control-Pediatric * 72-Fever  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Indicated for pain and fever control</li><li>Avoid in patients with severe liver disease</li></ul>	<ul style="list-style-type: none"> <li>1000 mg po</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>15 mg/kg po</li> </ul>
<b>Adenosine (Adenocard)</b>  <b>NCCEP Protocol:</b> * 16-Adult Tachycardia Narrow Complex * 17-Adult Tachycardia Wide Complex * 52-Pediatric Tachycardia  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Specifically for treatment or diagnosis of Supraventricular Tachycardia</li></ul>	<ul style="list-style-type: none"> <li>6 mg IV push over 1-3 seconds. If no effect after 1-2 minutes,</li> <li>Repeat with 12 mg IV push over 1-3 seconds.</li> <li>Repeat once if necessary</li> <li>(use stopcock and 20 ml Normal Saline flush with each dose)</li> </ul>	<ul style="list-style-type: none"> <li>0.1 mg/kg IV (Max 6 mg) push over 1-3 seconds. If no effect after 1-2 minutes,</li> <li>Repeat with 0.2 mg/kg IV (Max 12 mg) push over 1-3 seconds.</li> <li>Repeat once if necessary</li> <li>(use stopcock and Normal Saline flush with each dose)</li> </ul>
<b>Albuterol Beta-Agonist</b>  <b>NCCEP Protocol:</b> * 24-Allergic Reaction Anaphylaxis * 26-COPD Asthma * 56-Pediatric Allergic Reaction * 61-Pediatric Respiratory Distress  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Beta-Agonist nebulized treatment for use in respiratory distress with bronchospasm</li></ul>	<ul style="list-style-type: none"> <li>2.5-5.0 mg (3cc) in nebulizer continuously x 3 doses. See local protocol for relative contraindications and/or indications to contact medical control for use of this drug.</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>2.5mg (3cc) in nebulizer continuously x 3 doses. See local protocol for relative contraindications and/or indications to contact medical control for use of this drug.</li> </ul>

## Drug List A (Page 1 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Amiodarone (Cordarone)</b>  <b>NCCEP Protocol:</b> * 17-Adult Tachycardia Wide Complex * 18-VF Pulseless VT * 52-Pediatric Tachycardia * 53-Pediatric VF Pulseless VT * 54-Pediatric Post Resuscitation  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Antiarrhythmic used mainly in wide complex tachycardia and ventricular fibrillation.</li><li>Avoid in patients with heart block or profound bradycardia.</li><li>Contraindicated in patients with iodine hypersensitivity</li></ul>	<p><u>V-fib / pulseless V-tach</u></p> <ul style="list-style-type: none"> <li>300 mg IV push</li> <li>Repeat dose of 150 mg IV push for recurrent episodes</li> </ul> <p><u>V-tach with a pulse</u></p> <ul style="list-style-type: none"> <li>150 mg in 100cc D5W over 10 min</li> </ul>	<p><u>V-fib / pulseless V-tach</u></p> <ul style="list-style-type: none"> <li>5 mg/kg IV push over 5 minutes</li> <li>May repeat up to 15mg/kg IV</li> </ul> <p><u>V-tach with a pulse</u></p> <ul style="list-style-type: none"> <li>5 mg/kg IV push over 5 minutes</li> <li>May repeat up to 15mg/kg IV</li> <li>Avoid in Length Tape Color <b>Pink</b></li> </ul>
<b>Aspirin</b>  <b>NCCEP Protocol:</b> * 7-Pain Control Adult * 14-Chest Pain and STEMI  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>An antiplatelet drug for use in cardiac chest pain</li></ul>	<ul style="list-style-type: none"> <li>81 mg chewable (baby) Aspirin Give 4 tablets to equal usual adult dose.</li> </ul>	Ø
<b>Atropine</b>  <b>NCCEP Protocol:</b> * 12-Bradycardia Pulse Present * 49-Pediatric Bradycardia * 84-WMD Nerve Agent  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Anticholinergic drug used in bradycardias.</li><li>(For Endotracheal Tube use of this drug, double the dose)</li><li>In Organophosphate toxicity, large doses may be required (&gt;10 mg)</li></ul>	<p><u>Bradycardia</u></p> <ul style="list-style-type: none"> <li>0.5 - 1.0 mg IV every 3 – 5 minutes up to 3 mg. (If endotracheal -- max 6 mg)</li> </ul> <p><u>Organophosphate</u></p> <ul style="list-style-type: none"> <li>1-2 mg IM or IV otherwise as per medical control</li> </ul>	<p><b>See Color Coded List</b></p> <p><u>Bradycardia</u></p> <ul style="list-style-type: none"> <li>0.02 mg/kg IV, IO (Max 0.5 mg per dose, Max total dose 1mg IV)</li> <li>(Min 0.1 mg) per dose</li> <li>May repeat in 3 - 5 minutes</li> </ul> <p><u>Organophosphate</u></p> <ul style="list-style-type: none"> <li>0.05 mg/kg IV or IO otherwise as per medical control</li> </ul>

## Drug List A (Page 2 of 17)

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b><u>Atropine and Pralidoxime Auto-Injector Nerve Agent Kit</u></b>  <b>NCCEP Protocol:</b> * 84-WMD Nerve Agent  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>• Antidote for Nerve Agents or Organophosphate Overdose</li></ul>	<ul style="list-style-type: none"> <li>• One auto-injector then per medical control</li> </ul>	<ul style="list-style-type: none"> <li>• <b>See Color Coded List</b></li> <li>• One pediatric auto-injector then as per medical control</li> </ul>
<b><u>Calcium Chloride</u></b>  <b>NCCEP Protocol:</b> * 28-Dialysis Renal Failure * 31-Overdose Toxic Ingestion * 60-Ped OD Toxic Ingestion * 83-Marine Envenomations * 88-Crush Syndrome  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>• Indicated for severe hyperkalemia</li></ul>	<ul style="list-style-type: none"> <li>• 1 gm IV / IO</li> <li>• Avoid use if pt is taking digoxin</li> </ul>	<ul style="list-style-type: none"> <li>• <b>See Color Coded List</b></li> <li>• 20 mg/kg IV or IO slowly</li> </ul>
<b><u>Dextrose 10%, 25%, 50% Glucose solutions</u></b>  <b>NCCEP Protocol:</b> * Multiple  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>• Use in altered mental status or hypoglycemic states</li></ul>	See local protocol for concentration and dosing	<ul style="list-style-type: none"> <li>• <b>See Color Coded List</b></li> </ul> <p>See local protocol for concentration and dosing</p>

## Drug List A (Page 3 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<p><b>Diazepam</b> <b>(Valium)</b> <b>Benzodiazepene</b></p> <p><b>NCCEP Protocol:</b> * 32-Seizure * 39-Obstetrical Emergency * 62-Pediatric Seizure</p> <p><b>Indications/Contraindications:</b></p> <ul style="list-style-type: none"> <li>Seizure control</li> <li>Mild Sedation</li> </ul>	<ul style="list-style-type: none"> <li>4 mg IV / IO initially then 2 mg IV / IO every 3 - 5 minutes up to 10 mg max unless med control dictates</li> <li>Do not administer IM. The drug is not absorbed.</li> <li>10 mg Rectally if unable to obtain an IV.</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>0.1 - 0.3 mg/kg IV/IO</li> <li>(Max dose 4 mg IV, IO)</li> <li>0.5 mg/kg rectally (Dia-Stat)</li> <li>(Max dose 10 mg rectally)</li> <li>Repeat as directed by local protocol</li> </ul>
<p><b>Diltiazem</b> <b>(Cardizem)</b> <b>Calcium Channel Blocker</b></p> <p><b>NCCEP Protocol:</b> * 16-Adult Tachycardia Narrow Complex</p> <p><b>Indications/Contraindications:</b></p> <ul style="list-style-type: none"> <li>Calcium channel blocker used to treat narrow complex SVT</li> <li>Contraindicated in patients with heart block, ventricular tachycardia, and/or acute MI</li> </ul>	See local protocol for dosing	Ø
<p><b>Diphenhydramine</b> <b>(Benadryl)</b></p> <p><b>NCCEP Protocol:</b> * 24-Allergic Reaction Anaphylaxis * 56-Pediatric Allergic Reaction</p> <p><b>Indications/Contraindications:</b></p> <ul style="list-style-type: none"> <li>Antihistamine for control of allergic reactions</li> </ul>	<ul style="list-style-type: none"> <li>25-50 mg IV/IO/IM/PO</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>1 mg/kg IV/IO/IM/PO</li> <li>Do not give in infants &lt; 3 mo</li> </ul>

## Drug List A (Page 4 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Dopamine</b>	<ul style="list-style-type: none"> <li>2 - 20 micrograms/kg/min IV or IO, titrate to BP systolic of 90 mmHg</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>2 - 20 micrograms/kg/min IV or IO, titrate to BP systolic appropriate for age</li> </ul>
<b>Epinephrine 1:1,000</b>	<ul style="list-style-type: none"> <li>0.3 mg IM</li> <li>See local protocol for relative contraindications and/or indications to contact medical control for use of this drug.</li> </ul> <p><b>Nebulized Epinephrine</b></p> <ul style="list-style-type: none"> <li>1 mg mixed with 2 ml of Normal Saline</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>0.01 mg/kg IM</li> <li>(Max dose 0.3 mg)</li> </ul> <p><b>Nebulized Epinephrine</b></p> <ul style="list-style-type: none"> <li>1 mg mixed with 2 ml of Normal Saline</li> </ul>
<b>Epinephrine 1:10,000</b>	<ul style="list-style-type: none"> <li>1.0 mg IV / IO</li> <li>Repeat every 3 - 5 minutes until observe response</li> <li>(May be given by Endotracheal tube in double the IV dose)</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>0.01 mg/kg IV or IO</li> <li>(Max dose 1 mg)</li> <li>Repeat every 3 - 5 minutes per protocol</li> <li>(May be given by Endotracheal tube in double the IV dose)</li> </ul>
<b>Etomidate (Amidate)</b>	<ul style="list-style-type: none"> <li>0.3 mg/kg IV / IO</li> <li>Usual adult dose = 20 mg</li> </ul>	Ø

## Drug List A (Page 5 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Fentanyl (Sublimaze) Narcotic Analgesic</b>  <b>NCCEP Protocol:</b> <span style="color: red;">*</span> Multiple	<ul style="list-style-type: none"> <li>50-75 mcg IM/IV/IO bolus then 25 mcg IM/IV/IO every 20 minutes until a maximum of 200 mcg or clinical improvement</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>1 mcg/kg IM/IN/IV/IO May repeat 0.5 mcg/kg every 5 minutes Maximum dose 2 mcg/kg</li> </ul>
<b>Eurosemide (Lasix)</b>  <b>NCCEP Protocol:</b> <span style="color: red;">*</span> 15-CHF Pulmonary Edema <span style="color: red;">*</span> 50-Pediatric CHF Pulmonary Edema	<ul style="list-style-type: none"> <li>See local protocol for dosing guidelines</li> </ul>	<ul style="list-style-type: none"> <li>See local protocol for dosing guidelines</li> </ul>
<b>Glucagon</b>  <b>NCCEP Protocol:</b> <span style="color: red;">*</span> 27-Diabetic; Adult <span style="color: red;">*</span> 31-Overdose Toxic Ingestion <span style="color: red;">*</span> 58-Pediatric Diabetic <span style="color: red;">*</span> 60-Ped OD Toxic Ingestion	<ul style="list-style-type: none"> <li>1 - 2 mg IM</li> <li>Repeat blood glucose measurement in 15 minutes, if <math>\leq</math> 69 mg / dl repeat dose.</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>0.1 mg/kg IM, Maximum 1 mg</li> <li>Repeat blood glucose measurement in 15 minutes, if <math>\leq</math> 69 mg / dl repeat dose.</li> </ul>

## Drug List A (Page 6 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Glucose Oral</b> <b>Glucose Solutions</b>  <b>NCCEP Protocol:</b> <span style="color: red;">*</span> 27-Diabetic; Adult <span style="color: red;">*</span> 58-Pediatric Diabetic  <u>Indications/Contraindications:</u> <ul style="list-style-type: none"> <li>• Use in conscious hypoglycemic states</li> </ul>	<ul style="list-style-type: none"> <li>• One tube or packet</li> <li>• Repeat based on blood glucose results, per protocol</li> </ul>	<ul style="list-style-type: none"> <li>• See Color Coded List</li> <li>• One Tube or packet</li> <li>• Repeat based on blood glucose results, per protocol</li> <li>• Consider patient's ability to swallow and follow directions based on age</li> </ul>
<b>Haloperidol</b> <b>(Haldol)</b> <b>Phenothiazine</b> <b>Preperation</b>  <b>NCCEP Protocol:</b> <span style="color: red;">*</span> 6-Behavioral  <u>Indications/Contraindications:</u> <ul style="list-style-type: none"> <li>• Medication to assist with sedation of agitated patients</li> </ul>	<ul style="list-style-type: none"> <li>• 2.5-10 mg IV/IM, per local protocol</li> <li>• See local protocol for relative contraindications and/or indications to contact medical control for use of this drug.</li> </ul>	Ø

## Drug List A (Page 7 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<p><b>Ibuprofen</b> <b>(Motrin)</b> <b>Non-steroidal Anti-inflammatory Drug</b></p> <p><b>NCCEP Protocol:</b> * 7-Pain Control Adult * 46-Pediatric Pain Control * 72-Fever</p> <p><b>Indications/Contraindications:</b></p> <ul style="list-style-type: none"> <li>Avoid NSAIDS in women who are pregnant or could be pregnant.</li> <li>A nonsteroidal anti-inflammatory drug (NSAID) used for pain and fever control.</li> <li>Not to be used in patients with history of GI Bleeding (ulcers) or renal insufficiency.</li> <li>Not to be used in patients with allergies to aspirin or other NSAID drugs</li> <li>Avoid in patients currently taking anticoagulants, such as coumadin.</li> </ul>	<ul style="list-style-type: none"> <li>400-800 mg po</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>10 mg/kg po</li> <li>Do not use in patients 6 months of age or younger</li> </ul>
<p><b>Ipratropium</b> <b>(Atrovent)</b></p> <p><b>NCCEP Protocol:</b> * 24-Allergic Reaction Anaphylaxis * 26-COPD Asthma * 56-Pediatric Allergic Reaction * 61-Pediatric Respiratory Distress</p> <p><b>Indications/Contraindications:</b></p> <ul style="list-style-type: none"> <li>Medication used in addition to albuterol to assist in patients with asthma and COPD</li> </ul>	<ul style="list-style-type: none"> <li>2 puffs per dose of MDI (18 mcg/spray) --- OR ---</li> <li>0.5 mg per nebulizer treatment</li> </ul>	<ul style="list-style-type: none"> <li>Use in Pediatrics as a combined Therapy with a Beta Agonist such as Albuterol</li> <li>2 puffs per dose of MDI (18 mcg/spray) --- OR ---</li> <li>0.5 mg per nebulizer treatment</li> </ul>

## Drug List A (Page 8 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Ketorolac (Toradol) Non-steroidal Anti-inflammatory Drug</b>  <b>NCCEP Protocol:</b> <span style="color: orange;">*</span> 7-Pain Control Adult <span style="color: orange;">*</span> 46-Pediatric Pain Control  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"> <li>Avoid NSAIDS in women who are pregnant or could be pregnant</li> <li>A nonsteroidal anti-inflammatory drug used for pain control.</li> <li>Not to be used in patients with history of GI bleeding (ulcers), renal insufficiency, or in patients who may need immediate surgical intervention (i.e. obvious fractures).</li> <li>Not to be used in patients with allergies to aspirin or other NSAID drugs such as motrin</li> <li>Avoid in patients currently taking anticoagulants such as coumadin</li> </ul>	<ul style="list-style-type: none"> <li>30 mg IV / IO or 60 mg IM</li> </ul>	<ul style="list-style-type: none"> <li>0.5 mg/kg IV / IO / IM Maximum 30 mg</li> </ul>
<b>Lactated Ringer's Solution</b>  <b>NCCEP Protocol:</b> <span style="color: orange;">*</span> 40-Adult Thermal Burn <span style="color: orange;">*</span> 67-Pediatric Thermal Burn  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"> <li>Crystalloid solution preferred for fluid resuscitation and preferred in burn care.</li> </ul>	<ul style="list-style-type: none"> <li>Dosing per protocol, similar to Normal Saline</li> </ul>	<ul style="list-style-type: none"> <li>Dosing per protocol, similar to Normal Saline</li> </ul>

## Drug List A (Page 9 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Lidocaine</b>	<ul style="list-style-type: none"> <li>1.5 mg/kg IV / IO bolus (ETT dose = 2 x IV dose) up to 3mg/kg max bolus dose</li> <li>See local protocol for specific dosing algorithm</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>1 mg/kg IV / IO Maximum 100 mg Repeat 0.5 mg/kg Maximum 3 mg/kg total</li> </ul>
<b>Magnesium Sulfate</b>	<p><b>Respiratory Distress:</b></p> <ul style="list-style-type: none"> <li>2 g IV / IO over 10 minutes</li> <li>Repeat dosing per local protocol</li> </ul> <p><b>Obstetrical Seizure:</b></p> <ul style="list-style-type: none"> <li>2 g IV / IO over 2-3 minutes</li> <li>Dose may be repeated once, or as per local protocol</li> </ul>	<ul style="list-style-type: none"> <li>40 mg/kg IV / IO over 20 minutes (Max 2 gms)</li> <li>Repeat dosing per local protocol</li> </ul>
<b>Methylprednisolone (Solu-medrol) Steroid Preparation</b>	<ul style="list-style-type: none"> <li>125 mg IV / IO</li> <li>IM dosing only if indicated by local protocol</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>2 mg/kg IV / IO (Max 125 mg)</li> <li>IM dosing only if indicated by local protocol</li> </ul>

## Drug List A (Page 10 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Midazolam</b> <b>(Versed)</b> <b>Benzodiazepine</b>  <b>NCCEP Protocol:</b> <span style="color: red;">*</span> Multiple  <u>Indications/Contraindications:</u> <ul style="list-style-type: none"> <li>• Benzodiazepine used to control seizures and sedation</li> <li>• Quick acting Benzodiazepine</li> <li>• Preferred over Valium for IM use</li> <li>• Use with caution if BP &lt; 110</li> </ul>	<ul style="list-style-type: none"> <li>• See individual protocols for dosing</li> <li>• Usual total dose: 2.5-5 mg IV / IO / IM</li> </ul>	<ul style="list-style-type: none"> <li>• <b>See Color Coded List</b></li> <li>• See individual protocols for dosing</li> <li>• Usual total dose 0.1-0.2 mg/kg IV / IO / IM / IN</li> </ul>
<b>Morphine Sulfate</b> <b>Narcotic Analgesic</b>  <b>NCCEP Protocol:</b> <span style="color: red;">*</span> Multiple  <u>Indications/Contraindications:</u> <ul style="list-style-type: none"> <li>• Narcotic pain relief</li> <li>• Possible beneficial effect in pulmonary edema</li> <li>• Avoid use if BP &lt; 110</li> </ul>	<ul style="list-style-type: none"> <li>• 4 mg IM/IV/IO bolus then 2 mg IM/IV/IO every 5-10 minutes until a maximum of 10 mg or clinical improvement</li> </ul>	<ul style="list-style-type: none"> <li>• <b>See Color Coded List</b></li> <li>• 0.1 mg/kg IV / IO / IM May repeat every 5 minutes Maximum single dose 5 mg Maximum dose 10 mg</li> </ul>
<b>Naloxone</b> <b>(Narcan)</b> <b>Narcotic Antagonist</b>  <b>NCCEP Protocol:</b> <span style="color: red;">*</span> 31-Overdose Toxic Ingestion <span style="color: red;">*</span> 60-Ped OD Toxic Ingestion  <u>Indications/Contraindications:</u> <ul style="list-style-type: none"> <li>• Narcotic antagonist</li> </ul>	<ul style="list-style-type: none"> <li>• 0.4 - 2 mg IV / IO / IM / IN / ETT bolus titrated to patient's respiratory response</li> </ul>	<ul style="list-style-type: none"> <li>• <b>See Color Coded List</b></li> <li>• 0.1 mg/kg IV / IO / IN / IM / ETT (Max 2 mg)</li> <li>• Repeat as per protocol</li> </ul>

## Drug List A (Page 11 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Normal Saline</b> <b>Crystalloid Solutions</b>  <b>NCCEP Protocol:</b> * Multiple  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>IV fluid for IV access or volume infusion</li></ul>	<ul style="list-style-type: none"><li>See individual protocol for bolus dosing and/or infusion rate</li></ul>	<ul style="list-style-type: none"><li><b>See Color Coded List</b></li><li>See individual protocol for bolus dosing and/or infusion rate</li><li>Usual initial bolus 20 mL / kg IV / IO</li></ul>
<b>Nitroglycerin</b>  <b>NCCEP Protocol:</b> * 14-Chest Pain and STEMI * 15-CHF Pulmonary Edema  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Vasodilator used in anginal syndromes and CHF.</li></ul>	<ul style="list-style-type: none"><li>0.3 / 0.4 mg SL every 5 minutes until painfree</li><li>See Chest Pain Protocol for paste dosing</li></ul>	Ø
<b>Nitrous Oxide</b>  <b>NCCEP Protocol:</b> * 7-Pain Control Adult  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Medication used to assist with control of pain</li></ul>	<ul style="list-style-type: none"><li>Inhaled gas to effect per local protocol</li></ul>	<ul style="list-style-type: none"><li>Inhaled gas to effect per local protocol</li></ul>

## Drug List A (Page 12 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Ondansetron (Zofran) Anti-emetic</b>  <b>NCCEP Protocol:</b> * 23-Abdominal Pain Protocol * 35-Vomiting and Diarrhea * 63-Pediatric Vomiting and Diarrhea  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Anti-Emetic used to control Nausea and/or Vomiting</li><li>Ondansetron (Zofran) is the recommended Anti-emetic for EMS use since it is associated with significantly less side effects and sedation.</li></ul>	<ul style="list-style-type: none"><li>4 mg IV / IO / IM / PO / ODT</li><li>Repeat only as per local protocol</li></ul>	<ul style="list-style-type: none"><li>0.15 mg/kg IV / IO / IM (Max 4 mg)</li><li>0.2 mg/kg PO / ODT (Max 4 mg)</li><li>Repeat only as per local protocol</li></ul>
<b>Oxygen</b>  <b>NCCEP Protocol:</b> * Multiple  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Indicated in any condition with increased cardiac work load, respiratory distress, or illness or injury resulting in altered ventilation and/or perfusion. Goal oxygen saturation 94-99%.</li><li>Indicated for pre-oxygenation whenever possible prior to endotracheal intubation. Goal oxygen saturation 100%.</li></ul>	<ul style="list-style-type: none"><li>1-4 liters/min via nasal cannula</li><li>6-15 liters/min via NRB mask</li><li>15 liters via BVM / ETT / BIAD</li></ul>	<ul style="list-style-type: none"><li>1-4 liters/min via nasal cannula</li><li>6-15 liters/min via NRB mask</li><li>15 liters via BVM / ETT / BIAD</li></ul>

## Drug List A (Page 13 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Oxymetazoline (Afrin or Otrivin) Nasal Decongestant Spray</b>  NCCEP Protocol: * 71-Epistaxis  <u>Indications/Contraindications:</u> <ul style="list-style-type: none"><li>• Vasoconstrictor used with nasal intubation and epistaxis</li><li>• Relative Contraindication is significant hypertension</li></ul>	<ul style="list-style-type: none"><li>• 2 sprays in affected nostril</li><li>• Usual concentration is 0.05% by volume</li></ul>	<ul style="list-style-type: none"><li>• <b>See Color Coded List</b></li><li>• 1-2 sprays in affected nostril</li><li>• Usual concentration is 0.05% by volume</li></ul>
<b>Pralidoxime (2-PAM)</b>  NCCEP Protocol: * 84-WMD Nerve Agent  <u>Indications/Contraindications:</u> <ul style="list-style-type: none"><li>• Antidote for Nerve Agents or Organophosphate Overdose</li><li>• Administered with Atropine</li></ul>	<ul style="list-style-type: none"><li>• 600 mg IV / IO / IM over 30 minutes for minor symptoms</li><li>• 1800 mg IV / IO / IM over 30 minutes for major symptoms</li><li>• See local protocol for minor versus major indications</li></ul>	<ul style="list-style-type: none"><li>• 15 – 25 mg/kg IV / IM / IO over 30 minutes</li><li>• See local protocol for specific pediatric dosing recommendations</li></ul>

## Drug List A (Page 14 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Promethazine (Phenergan) Anti-emetic</b>  <b>NCCEP Protocol:</b> * 23-Abdominal Pain * 35-Vomiting and Diarrhea  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>IV Promethazine (Phenergan) should be given IV only with great caution. Extravasation of this drug can result in significant local tissue damage.</li><li>Anti-Emetic used to control Nausea and/or Vomiting</li><li>Ondansetron (Zofran) is the recommended Anti-emetic for EMS use since it is associated with significantly less side effects and sedation.</li></ul>	<ul style="list-style-type: none"> <li>12.5 mg IV / IO / IM</li> <li>May repeat as per local protocol</li> </ul>	Ø
<b>Rocuronium</b>  <b>NCCEP Protocol:</b> * 4-Airway Rapid Sequence Intubation  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Non-depolarizing paralytic agent used as a component of drug assisted intubation (Rapid Sequence Intubation), when succinylcholine is contraindicated.</li><li>Onset of action is longer than succinylcholine, up to 3 minutes, patient will NOT defasciculate.</li></ul>	<ul style="list-style-type: none"> <li>1 mg / kg IV / IO</li> <li>Only may repeat x1 per RSI protocol</li> </ul>	Ø

## Drug List A (Page 15 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Sodium Bicarbonate</b>	<ul style="list-style-type: none"> <li>Initial bolus 50 mEq IV / IO</li> <li>See individual protocol for specific dosing algorithm.</li> </ul>	<ul style="list-style-type: none"> <li><b>See Color Coded List</b></li> <li>Initial bolus 1 mEq / kg IV / IO</li> <li>Maximum 50 mEq</li> <li>See individual protocol for specific dosing algorithm.</li> </ul>
<b>Succinylcholine</b> <b>Paralytic Agent</b>	<ul style="list-style-type: none"> <li>1.5 mg/kg IV / IO</li> <li>Only may repeat x1 per RSI protocol</li> </ul>	∅

## Drug List A (Page 16 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Drug List



ONLY some medications that are included by name and dose in the 2024 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference for the 2024 NCCEP Protocols.

For a full list of medications approved for use by EMS professionals, please refer to the North Carolina Medical Board document entitled: Approved Medications for Credentialed EMS Personnel. Individual EMS Systems may or may not utilize these or other approved drugs. Individual EMS Systems are strongly encouraged to maintain a system-specific drug list for use with system-specific protocols for daily operations and training. See the Pediatric Color Coded Drug List for pediatric dosages

Medication	Adult Dosing	Pediatric Dosing
<b>Vasopressin (Pitressin)</b>  <b>NCCEP Protocol:</b> * 11-Asystole Pulseless Electrical Activity * 18-VF Pulseless VT  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Medication used in place of and/or in addition to epinephrine in the setting of cardiac arrest</li></ul>	<ul style="list-style-type: none"><li>40 units IV / IO, may replace first or second dose of epinephrine</li></ul>	Ø
<b>Vecuronium Paralytic Agent</b>  <b>NCCEP Protocol:</b> * 4-Airway Rapid Sequence Intubation * 19-Post Resuscitation * 20-Induced Hypothermia  <b>Indications/Contraindications:</b> <ul style="list-style-type: none"><li>Long-acting non-depolarizing paralytic agent</li><li>Avoid in patients with chronic neuromuscular disease (e.g., muscular dystrophy).</li></ul>	<ul style="list-style-type: none"><li>0.1 mg/kg IV / IO or 10 mg IV / IO, as per individual protocol</li><li>Only may repeat dosing as per individual protocol</li></ul>	Ø

## Drug List A (Page 17 of 17 )

This formulary is provided as a reference only. It does not contain all of the contraindications and potential adverse reactions for each listed drug. It is the responsibility of each EMS System, Agency, and Medical Director to assure that each EMS professional is knowledgeable about the use each drug in this formulary.



# Pediatric Color Coded Drug List



Length < 59.5 cm		Weight 3-5 Kg (Avg 4.0 Kg)				Gray (0-3 months)	
<b>Vital Signs</b>							
Heart Rate 120-150		Acetaminophen 64 mg				Epinephrine 1:10,000 0.04 mg	
Respirations 24-48		Adenosine 1 <sup>st</sup> Dose- 0.3 mg				Epinephrine 1:1000 Nebulized 2.0 mg	
BP Systolic 70 (+/-25)		Repeat Dose- 0.6 mg				Epinephrine 1:1000 IM 0.05 mg	
<b>Equipment</b>		Afrin Nasal Spray HOLD				Fentanyl 8.0 mcg	
ET Tube 2.5 - 3.5		Albuterol 2.5mg				Glucagon 0.5 mg	
Blade Size 0 - 1		Amiodarone 20 mg				Ibuprofen N/A	
<b>Defibrillation</b>		Atropine 0.10 mg				Ipratropium 500 mcg	
Defibrillation 8 J, 15 J		Calcium Chloride 80 mg				Leva buterol 0.31 mg	
Cardioversion 2 J, 4 J		Charcoal N/A				Lidocaine 4 mg	
<b>Normal Saline</b> 80 ml		Dextrose 10% 20 ml				Lorazepam 0.2 mg	
		Diazepam (IV) 0.8 mg				Magnesium Sulfate 200 mg	
		(Rectal) 2.0 mg				Methylprednisolone 6.25 mg	
		Dilauidid HOLD				Midazolam 0.5 mg	
		Diphenhydramine 6.5 mg				Morphine Sulfate 0.4 mg	
		Dopamine (800 mg in 500 cc) 2 mcg/kg/min 0.3 ml/hr				Naloxone 0.4 mg	
		5 mcg/kg/min 0.9 ml/hr				Ondansetron 0.6 mg	
		10 mcg/kg/min 1.7 ml/hr				Prednisone 4.0 mg	
		20 mcg/kg/min 3.3 ml/hr				Sodium Bicarbonate 4 mEq	

Length 59.5-66.5 cm		Weight 6-7 Kg (Avg 6.5 Kg)				Pink (3-6 Months)	
<b>Vital Signs</b>		Acetaminophen 96 mg				Epinephrine 1:10,000 0.06 mg	
Heart Rate 120-125		Adenosine 1 <sup>st</sup> Dose- 0.6 mg				Epinephrine 1:1000 Nebulized 2.0 mg	
Respirations 24-48		Repeat Dose- 1.2 mg				Epinephrine 1:1000 IM 0.06 mg	
BP Systolic 85 (+/-25)		Afrin Nasal Spray HOLD				Fentanyl 13.0 mcg	
<b>Equipment</b>		Albuterol 2.5 mg				Glucagon 0.5 mg	
ET Tube 3.5		Atropine 0.13 mg				Ibuprofen N/A	
Blade Size 1		Amiodarone 30 mg				Ipratropium 500 mcg	
<b>Defibrillation</b>		Calcium Chloride 130 mg				Leva buterol 0.31 mg	
Defibrillation 10 J, 20 J		Charcoal HOLD				Lidocaine 6 mg	
Cardioversion 2 J, 5 J		Dextrose 10% 35 ml				Lorazepam 0.33 mg	
<b>Normal Saline</b> 130 ml		Diazepam (IV) 1.3 mg				Magnesium Sulfate 300 mg	
		(Rectal) 3.2 mg				Methylprednisolone 12.5 mg	
		Dilauidid HOLD				Midazolam 0.5 mg	
		Diphenhydramine 5 mg				Morphine Sulfate 0.6 mg	
		Dopamine (800 mg in 500 cc) 2 mcg/kg/min 0.5 ml/hr				Naloxone 0.6 mg	
		5 mcg/kg/min 1.3 ml/hr				Ondansetron 1.0 mg	
		10 mcg/kg/min 2.5 ml/hr				Prednisone 6.5 mg	
		20 mcg/kg/min 5.0 ml/hr				Sodium Bicarbonate 6 mEq	

Length 66.5-74 cm		Weight 8-9 Kg (Avg 8.5 Kg)				Red (7-10 Months)	
<b>Vital Signs</b>		Acetaminophen 128 mg				Epinephrine 1:10,000 0.08 mg	
Heart Rate 120		Adenosine 1 <sup>st</sup> Dose- 0.9 mg				Epinephrine 1:1000 Nebulized 2.0 mg	
Respirations 24-32		Repeat Dose- 1.8 mg				Epinephrine 1:1000 IM 0.08 mg	
BP Systolic 92 (+/-30)		Afrin Nasal Spray HOLD				Fentanyl 17.0 mcg	
<b>Equipment</b>		Albuterol 2.5 mg				Glucagon 0.5 mg	
ET Tube 3.5-4.0		Atropine 0.17 mg				Ibuprofen 4.0 ml	
Blade Size 1		Amiodarone 40 mg				Ipratropium 500 mcg	
<b>Defibrillation</b>		Calcium Chloride 170 mg				Leva buterol 0.31 mg	
Defibrillation 20 J, 40 J		Charcoal HOLD				Lidocaine 8 mg	
Cardioversion 5 J, 9 J		Dextrose 10% 43 ml				Lorazepam 0.43 mg	
<b>Normal Saline</b> 170 ml		Diazepam (IV) 1.7 mg				Magnesium Sulfate 400 mg	
		(Rectal) 4.25 mg				Methylprednisolone 12.5 mg	
		Dilauidid HOLD				Midazolam 0.85 mg	
		Diphenhydramine 10 mg				Morphine Sulfate 0.8 mg	
		Dopamine (800 mg in 500 cc) 2 mcg/kg/min 0.7 ml/hr				Naloxone 0.8 mg	
		5 mcg/kg/min 1.6 ml/hr				Ondansetron 1.2 mg	
		10 mcg/kg/min 3.2 ml/hr				Prednisone 8.5 mg	
		20 mcg/kg/min 6.5 ml/hr				Sodium Bicarbonate 8 mEq	



# Pediatric Color Coded Drug List



Length 74-84.5 cm

<b>Vital Signs</b>	
Heart Rate	115-120
Respirations	22-30
BP Systolic	96 (+/-30)
<b>Equipment</b>	
ET Tube	4.0
Blade Size	1
<b>Defibrillation</b>	
Defibrillation	20 J, 40 J
Cardioversion	5 J, 10 J
<b>Normal Saline</b>	
210 ml	

## Weight 10-11 Kg (Avg 10.5 Kg)

Acetaminophen	160 mg	Epinephrine 1:10,000	0.1 mg
Adenosine	1 <sup>st</sup> Dose- Repeat Dose-	0.9 mg 1.8 mg	Epinephrine 1:1000 IM Epinephrine 1:1000 Nebulized
Afrin Nasal Spray	HOLD	2.0 mg	2.0 mg
Albuterol	2.5 mg	Fentanyl	21.0 mcg
Atropine	0.2 mg	Glucagon	1.0 mg
Amiodarone	50 mg	Ibuprofen	5.0 ml
Calcium Chloride	210 mg	Ipratropium	500 mcg
Charcoal	HOLD	Leva buterol	0.63 mg
Dextrose 10%	50 ml	Lidocaine	10 mg
Diazepam	(IV) 1.0 mg (Rectal) 5.0 mg	Lorazepam	0.53 mg
Dilauidid	HOLD	Magnesium Sulfate	500 mg
Diphenhydramine	10 mg	Methylprednisolone	18.75 mg
Dopamine	(800 mg in 500 ml Normal Saline)	Midazolam	1.0 mg
	2 mcg/kg/min 0.8 ml/hr	Morphine Sulfate	1.0 mg
	5 mcg/kg/min 2.0 ml/hr	Naloxone	1.0 mg
	10 mcg/kg/min 4.0 ml/hr	Ondansetron	1.6 mg
	20 mcg/kg/min 8.0 ml/hr	Prednisone	10.5 mg
		Sodium Bicarbonate	10 mEq

Purple (11-18 Months)

Length 84.5-97.5 cm

<b>Vital Signs</b>	
Heart Rate	110-115
Respirations	20-28
BP Systolic	100(+/-30)
<b>Equipment</b>	
ET Tube	4.5
Blade Size	2
<b>Defibrillation</b>	
Defibrillation	30 J, 50 J
Cardioversion	6 J, 15 J
<b>Normal Saline</b>	
260 ml	

## Weight 12-14 Kg (Avg 13 Kg)

Acetaminophen	192 mg	Epinephrine 1:10,000	0.10 mg
Adenosine	1 <sup>st</sup> Dose- Repeat Dose-	1.2 mg 2.4 mg	Epinephrine 1:1000 IM Epinephrine 1:1000 Nebulized
Afrin Nasal Spray	1 spray	2.0 mg	2.0 mg
Albuterol	2.5 mg	Fentanyl	26.0 mcg
Atropine	0.26 mg	Glucagon	0.5 mg
Amiodarone	65 mg	Ibuprofen	6.5 ml
Calcium Chloride	260 mg	Ipratropium	500 mcg
Charcoal	15 gms	Leva buterol	0.63 mg
Dextrose 10%	60-80 ml	Lidocaine	14 mg
Diazepam	(IV) 2.6 mg (Rectal) 6.5 mg	Lorazepam	0.65 mg
Dilauidid	HOLD	Magnesium Sulfate	650 mg
Diphenhydramine	10 mg	Methylprednisolone	25.0 mg
Dopamine	(800 mg in 500 ml Normal Saline)	Midazolam	1 mg
	2 mcg/kg/min 0.8 ml/hr	Morphine Sulfate	1.0 mg
	5 mcg/kg/min 2.5 ml/hr	Naloxone	1.3 mg
	10 mcg/kg/min 5.0 ml/hr	Ondansetron	2.0 mg
	20 mcg/kg/min 10 ml/hr	Prednisone	13.0 mg
		Sodium Bicarbonate	13 mEq

Yellow (19-35 Months)

Length 97.5-110 cm

<b>Vital Signs</b>	
Heart Rate	100-15
Respirations	20-26
BP Systolic	100(+/-20)
<b>Equipment</b>	
ET Tube	5.0
Blade Size	2
<b>Defibrillation</b>	
Defibrillation	30 J, 70 J
Cardioversion	8 J, 15 J
<b>Normal Saline</b>	
330 ml	

## Weight 15-18 Kg (Avg 16.5 Kg)

Acetaminophen	240 mg	Epinephrine 1:10,000	0.16 mg
Adenosine	1 <sup>st</sup> Dose- Repeat Dose-	1.8 mg 3.6 mg	Epinephrine 1:1000 IM Epinephrine 1:1000 Nebulized
Afrin Nasal Spray	1 spray	2.0 mg	2.0 mg
Albuterol	2.5 mg	Fentanyl	33.0 mcg
Atropine	0.32 mg	Glucagon	0.5 mg
Amiodarone	80 mg	Ibuprofen	8.0 ml
Calcium Chloride	330 mg	Ipratropium	500 mcg
Charcoal	15-30 gms	Leva buterol	0.63 mg
Dextrose 10%	80 ml	Lidocaine	15 mg
Diazepam	(IV) 3.3 mg (Rectal) 8.25 mg	Lorazepam	0.83 mg
Dilauidid	HOLD	Magnesium Sulfate	800 mg
Diphenhydramine	15 mg	Methylprednisolone	31.25 mg
Dopamine	(800 mg in 500 ml Normal Saline)	Midazolam	1.5 mg
	2 mcg/kg/min 1.2 ml/hr	Morphine Sulfate	1.0 mg
	5 mcg/kg/min 3.0 ml/hr	Naloxone	1.6 mg
	10 mcg/kg/min 6.0 ml/hr	Ondansetron	2.4 mg
	20 mcg/kg/min 12 ml/hr	Prednisone	16.5 mg
		Sodium Bicarbonate	16 mEq

White (3-4 yrs)



# Pediatric Color Coded Drug List



## Weight 19-22 Kg (Avg 20.75 Kg)

<b>Vital Signs</b> Heart Rate 100 Respirations 20-24 BP Systolic 100(+/-15)	Acetaminophen 288 mg Adenosine 1 <sup>st</sup> Dose- 2.1 mg Repeat Dose- 4.1 mg Afrin Nasal Spray 1 spray Albuterol 2.5 mg Atropine 0.4 mg Amiodarone 100 mg Calcium Chloride 420 mg Charcoal 20-40 gms Dextrose 10% 100 ml Diazepam (IV) 4.0 mg (Rectal) 10.0 mg Dilaudid 0.31 mg Diphenhydramine 25.0 mg Dopamine (800 mg in 500 ml Normal Saline) 2 mcg/kg/min 1.6 ml/hr 5 mcg/kg/min 3.9 ml/hr 10 mcg/kg/min 7.8 ml/hr 20 mcg/kg/min 16 ml/hr	Epinephrine 1:10,000 0.2 mg Epinephrine 1:1000 IM 0.2 mg Epinephrine 1:1000 Nebulized 2.0 mg Fentanyl 40.0 mcg Glucagon 1.0 mg Ibuprofen 10.0 ml Ipratropium 500 mcg Levalbuterol 0.63 mg Lidocaine 20 mg Lorazepam 1.0 mg Magnesium Sulfate 1000 mg Methylprednisolone 37.5 mg Midazolam 2.0 mg Morphine Sulfate 2.0 mg Naloxone 2.0 mg Ondansetron 3.0 mg Prednisone 20.0 mg Sodium Bicarbonate 20 mEq
<b>Equipment</b> ET Tube 5.5 Blade Size 2		
<b>Defibrillation</b> Defibrillation 40 J, 85 J Cardioversion 10 J, 20 J		
<b>Normal Saline</b> 410 ml		

Blue (5-6 yrs)

## Weight 24-30 Kg (Avg 27 Kg)

<b>Vital Signs</b> Heart Rate 90 Respirations 18-22 BP Systolic 105(+/-15)	Acetaminophen 384 mg Adenosine 1 <sup>st</sup> Dose- 2.7 mg Repeat Dose- 5.4 mg Afrin Nasal Spray 1 spray Albuterol 2.5 mg Atropine 0.5 mg Amiodarone 135 mg Calcium Chloride 540 mg Charcoal 25-50 gms Dextrose 10% 135 ml Diazepam (IV) 4.0 mg (Rectal) 10.0 mg Dilaudid 0.4 mg Diphenhydramine 25 mg Dopamine (800 mg in 500 ml Normal Saline) 2 mcg/kg/min 2 ml/hr 5 mcg/kg/min 5 ml/hr 10 mcg/kg/min 10 ml/hr 20 mcg/kg/min 20 ml/hr	Epinephrine 1:10,000 0.27 mg Epinephrine 1:1000 IM 0.3 mg Epinephrine 1:1000 Nebulized 2.0 mg Fentanyl 54.0 mcg Glucagon 1.0 mg Ibuprofen 13 ml Ipratropium 500 mcg Levalbuterol 0.63 mg Lidocaine 20 mg Lorazepam 1.35 mg Magnesium Sulfate 1350 mg Methylprednisolone 54.0 mg Midazolam 2.0 mg Morphine Sulfate 2.0 mg Naloxone 2.0 mg Ondansetron 4.0 mg Prednisone 27.0 mg Sodium Bicarbonate 27 mEq
<b>Equipment</b> ET Tube 6.0 Blade Size 2-3		
<b>Defibrillation</b> Defibrillation 50 J, 100 J Cardioversion 15 J, 30 J		
<b>Normal Saline</b> 540 ml		

Orange (7-9 yrs)

## Weight 32-40 Kg (Avg 36 Kg)

<b>Vital Signs</b> Heart Rate 85-90 Respirations 16-22 BP Systolic 115(+/-20)	Acetaminophen 544 mg Adenosine 1 <sup>st</sup> Dose- 3.6 mg Repeat Dose- 7.2 mg Afrin Nasal Spray 2 spray Albuterol 2.5 mg Atropine 0.5 mg Amiodarone 180 mg Calcium Chloride 700 mg Charcoal 25-50 gms Dextrose 10% 180 ml Diazepam (IV) 4.0 mg (Rectal) 10.0 mg Dilaudid 0.54 mg Diphenhydramine 35 mg Dopamine (800 mg in 500 ml Normal Saline) 2 mcg/kg/min 2.7 ml/hr 5 mcg/kg/min 7.0 ml/hr 10 mcg/kg/min 14.0 ml/hr 20 mcg/kg/min 28.0 ml/hr	Epinephrine 1:10,000 0.3 mg Epinephrine 1:1000 IM 0.3 mg Epinephrine 1:1000 Nebulized 2.0 mg Fentanyl 62.0 mcg Glucagon 1.0 mg Ibuprofen 18 ml Ipratropium 500 mcg Levalbuterol 0.63 mg Lidocaine 36 mg Lorazepam 1.8 mg Magnesium Sulfate 1800 mg Methylprednisolone 62.5 mg Midazolam 2.0 mg Morphine Sulfate 3.0 mg Naloxone 2.0 mg Ondansetron 4.0 mg Prednisone 36.0 mg Sodium Bicarbonate 36 mEq
<b>Equipment</b> ET Tube 6.5 Blade Size 3		
<b>Defibrillation</b> Defibrillation 60 J, 150 J Cardioversion 15 J, 30 J		
<b>Normal Saline</b> 720 ml		

Green (10-12 yrs)